

MILWAUKEE PUBLIC MUSEUM

---

# Contributions

in  
BIOLOGY  
and  
GEOLOGY

---

Number 51

November 29, 1982

A Compendium of Fossil Marine Families

J. John Sepkoski, Jr.

MILWAUKEE PUBLIC MUSEUM

---

# Contributions

in  
BIOLOGY  
and  
GEOLOGY

---

Number 51

November 29, 1982

## A COMPENDIUM OF FOSSIL MARINE FAMILIES

**J. JOHN SEPKOSKI, JR.**  
*Department of the Geophysical Sciences  
University of Chicago*

---

**REVIEWERS FOR THIS PUBLICATION:**

Robert Gernant, University of Wisconsin-Milwaukee

David M. Raup, Field Museum of Natural History

Frederick R. Schram, San Diego Natural History Museum

Peter M. Sheehan, Milwaukee Public Museum

---

---

**ISBN 0-893260-081-9**

---

Milwaukee Public Museum Press

Published by the Order of the Board of Trustees

## CONTENTS

Abstract -----	2
Introduction-----	2
Compendium -----	6
Protozoa -----	6
Porifera-----	9
Archaeocyatha-----	14
Coeleenterata-----	17
Platyhelminthes-----	24
Rhynchocoela -----	24
Priapulida -----	24
Nematoda -----	24
Mollusca -----	24
Sipunculida -----	46
Echiurida -----	46
Annelida -----	46
Incertae sedis -----	49
Arthropoda -----	49
Phoronida -----	63
Bryozoa -----	63
Brachiopoda -----	67
Chaetognatha -----	73
Echinodermata -----	73
Pogonophora -----	86
Conodonts -----	87
Hemichordata -----	88
Chordata -----	89
Bibliography -----	102

## ABSTRACT

A comprehensive listing of the stratigraphic ranges of all established fossil marine animal families is presented. This listing summarizes previous compilations and provides updated taxonomic classifications and enhanced stratigraphic resolution. Other special features are coverage of both invertebrates and vertebrates, inclusion of problematica, and incorporation of previously unavailable stage-level data for the Vendian and Cambrian Systems.

## INTRODUCTION

This compendium lists approximately 3300 animal families and their known stratigraphic ranges in the marine fossil record. It summarizes and informally updates information available in the *Treatise on Invertebrate Paleontology* (Moore et al. 1953-1981), *The Fossil Record* (Harland et al 1967), and *Vertebrate Paleontology* (Romer 1966), providing a comprehensive listing at a consistent taxonomic level of all animals known through all time within a single large-scale ecosystem, the world ocean. This information was originally compiled for my studies of the evolution of taxonomic diversity through the Phanerozoic (Sepkoski 1978, 1979, 1981a, b). By publishing this compendium, I hope to make these data widely available for other studies of diversity and evolutionary rates as well as provide a concise reference on the classification and ages of marine animals and a starting point for further revision of data from the fossil record.

There are several important features of this compendium that warrant its publication at this time:

1. The classification employed is both comprehensive and consistent. An attempt has been made to compile all currently accepted families of marine protozoans, invertebrates, and vertebrates that have documented fossil records. Thus, this compendium includes information on several taxonomic groups not covered in other summaries, including coeno- and opisthobranch gastropods and coleoid cephalopods. Also, many problematical taxa (e.g. Burgess Shale soft-bodied organisms and Tommotian phosphatic shells) are listed in their most probable phyla.

2. The classification is up-to-date. Revised classifications of fossil classes published since the *Treatise* have been substituted wherever feasible. Groups affected include the Sclerospongia (Stromatoporoidea), Priapulida, Monoplacophora, Rostroconchia, Hyolitha, Polychaeta, Edrioasteroidea, Eocrinoidea, and Placodermi. In addition, numerous recently erected families and orders have been included in this compendium.

3. The stratigraphic information is of the highest resolution presently available. Nearly 85% of the data on first and last occurrences of families are listed in terms of 84 internationally recognizable stratigraphic stages (Table 1); these include 10 stage-level subdivisions of the Cambrian and 4 of the Vendian which have not been available in previous compilations. Most of the remaining data are resolved to one of 30 stratigraphic series. Much of the enhanced stratigraphic resolution of this

compendium is based on monographs and secondary sources published subsequent to the previous summaries (see Bibliography).

The particular qualities of this compendium must, of course, be balanced against the shortcomings characteristic of this and other such compilations. Despite all attempts to be current, some of the data presented here are undoubtedly obsolescent. A number of class and ordinal classifications are in need of major revision, including, most notably, those for the Porifera, Trilobita, Bryozoa, and Osteichthyes. I have attempted to incorporate partial revisions of these and other groups wherever possible but, I am sure, have missed many pertinent publications and have misunderstood taxonomic subtleties in others. Clearly, the classifications presented here should in no way be considered authoritative. The stratigraphic data, too, must be considered an approximation of the truth at best. New discoveries extending the stratigraphic ranges of established families and altering understanding of intra- and inter-continental correlations are being published continuously, keeping data on the fossil record in constant flux. (In one year alone while preparing this compilation, I needed to change my information on nearly 10% of the families as a result of new publications.)

The format used to present the data is rather simple. With few exceptions, the families are arranged in an abbreviated classification with only phyla, classes, and orders listed. The ordering of these higher taxa generally follows that used by Kukalová-Peck (1973). The principal sources of information on the classification and stratigraphic ranges of the families are given, along with miscellaneous notes, under each class. More specific sources of taxonomic and/or stratigraphic information are indicated for individual families by numbered codes in parentheses which refer to numbered entries in the bibliography.

Families are listed in alphabetic (and not "evolutionary") arrangement beneath each taxonomic order. Only families with marine fossil records are listed, and nonmarine families and approximately 1000 extant families without documented fossil occurrences are excluded. (The approximate number of these extant-only families is indicated for most classes.) Occasionally, distinctive genera that probably merit familial status or very early genera of uncertain familial relationships are listed in addition to the established families.

A pair of stratigraphic codes indicating the documented times of first and last appearance in the marine fossil record are given with each family. These codes consist generally of a single letter indicating geologic system followed, when known, either by a four-symbol code for stage or by a one- or two-letter code for series, both enclosed in parentheses. A complete listing of these stratigraphic codes is given in Table 1.

			Stage	Series	System
QUATERNARY		Recent (= Holocene) Pleistocene	R Q (Plei)		
TERTIARY	Pliocene		T (Plio)		
	Miocene	upper (= Messinian) middle (= Tortonian) lower (= Aquitanian)	T (Mi-u) T (Mi-m) T (Mi-l)		T (Mi)
	Oligocene	upper (= Chattian) lower (= Rupelian)	T (Ol-u) T (Ol-l)		T (Ol)
	Eocene	upper (= Bartonian) middle (= Lutetian) lower (= Ypresian)	T (Eo-u) T (Eo-m) T (Eo-l)		T (Eo)
	Paleocene	"Paleocene" (= Thanetian) Danian	T (Pale) T (Dan)		
	Upper Seno- mian	Maestrichtian Campanian Santonian Coniacian Turonian Cenomanian	K (Maes) K (Camp) K (Sant) K (Coni) K (Turo) 	K (Seno)	K (u)
CRETACEOUS	Lower Neoco- mian	Albian Aptian Barremian Hauterivian Valangian Berriasian	K (Albi) K (Apti) K (Barr) K (Haut) K (Vala) 		K (l)
	Upper	Tithonian (= Portlandian) Kimmeridgian Oxfordian	J (Tith) J (Kimm) J (Oxfo)		J (u)
	Middle	Callovian Bathonian Bajocian (+ Aalenian)	J (Call) J (Bath) J (Bajo)		J (m)
	Lower	Toarcian Pliensbachian Sinemurian Hettangian	J (Toar) J (Plie) J (Sine) J (Hett)		J (l)
JURASSIC	Upper	Rhaetian Norian Carnian	Tr (Rhae) Tr (Nori) Tr (Carn)		Tr (u)
	Middle	Ladinian Anisian	Tr' (Ladi) Tr (Anis)		Tr (m)
	Lower	Olenekian Induan	Tr (Olen) Tr (Indu)		Tr (l)
TRIASSIC	Upper	Dzhulfian (= Tartarian, Ochoan) Guadelupian (= Kazanian)	P (Dzhu) P (Guad)		P (u)
	Middle				Tr
	Lower	Leonardian (= Artinskian) Sakmarian (= u. Wolfcampian) Asselian (= l. Wolfcampian)	P (Leon) P (Sakm) P (Asse)		P (l)
PERMIAN					P

			Stage	Series	System
CARBON- FEROUS	Pennsylvanian	Stephanian	(= Uralian)	C (Step)	C
		Moscovian	(= Westphalian C & D)	C (Mosc)	
		Bashkirian	(= Westphalian A & B)	C (Bash)	
		Namurian		C (Nam)	
		Visean		C (Vise)	
	Mississippian	Tournaisean		C (Tour)	C
		Upper	Famennian Frasnian	D (Fame) D (Fras)	
		Middle	Givetian Eifelian	D (Give) D (Eife)	
		Lower	Emsian Siegenian Gedinnian	D (Emsi) D (Sieg) D (Gedi)	
		Upper	= Ludlovian + Pridolian	S (u)	
DEVONIAN	Upper	Middle	= Wenlockian	S (m)	S
		Lower	= Llandoveryan	S (l)	
		Upper	Ashgillian	Θ (Ashg)	
	Middle	Middle	Caradocian	Θ (Cara)	Θ
		Lower	Llandeilian	Θ (Llde)	
		Upper	Llanvirnian	Θ (Llv)	
	Lower	Middle	Arenigian	Θ (Aren)	Θ
		Lower	Tremadocian	Θ (Trem)	
		Upper	U3 (= Trempealeauan) U2 (= Franconian) U1 (= Dresbachian)	€ (Trep) € (Fran) € (Dres)	
CAMBRIAN	Upper	Middle	M3 M2 (= Templetonian) M1 (= Ordian, = Elankian)	€ (uMid) € (mMid) € (lMid)	€
		Lower	L3 (= "Botomian" or Lenian s.s.) L2 (= Attabanian) L1 (= Tommotian)   L1b (= Kenyadian)   L1a (= Sunnagin)	€ Boto € (Atda) € (uTom) € (!Tom)	
		Upper	= Nemakit-Daldyn	V (N-Da) V (u) V (m) V (l)	
		Middle			
		Lower			
VENDIAN	upper { up. middle lower		upper	u Riph	V
RIPHEAN					

**Table 1.** Stratigraphic subdivisions and codes used in this compendium. Each code listed in the three right hand columns signifies a specific stage, series, or system as listed in the equivalent, mirror-image column on the left. The stratigraphic subdivisions of the Ordovician through Quaternary Systems are essentially the same as those used by Harland et al. (1967) in *The Fossil Record*; the subdivisions of the Vendian and Cambrian Systems are explained in Sepkoski (1979, Fig. 2 and Appendix 1).

## COMPENDIUM

## PROTOZOA

## Cl. Rhizopodea

[Classification and stratigraphic ranges from the *Treatise*, Pt. C, and *The Fossil Record*. Approximately 13 extant free-living marine families (including all in the Amoebida) without known fossil representatives are not listed.]

Or. Gromiida			
Amphitrematidae	Q (Plei)	— R	
Gromiidae	Q (Plei)	— R	
Or. Foraminiferida			
Acervulinidae	T (Eo)	— R	
Allogromiidae	Є (Fran)	— R	
Alveolinidae	K (Albi)	— R	
Ammodiscidae	Θ (Aren)	— R	
Amphisteginidae	T (Pale)	— R	
Annulopatellinidae	T (Mi-l)	— R	
Anomalinidae	Tr (Rhae)	— R	
Archaediscidae	C (Tour)	— P (Dzhu)	
Asterigerinidae	K (Camp)	— R	
Astrorhizidae	Θ (Aren)	— R	
Ataxophragmiidae	C (Step)	— R	
Barkerinidae	K (Albi)	— K (Turo)	
Biseriamminidae	C (Tour)	— P (Dzhu)	
Bolivinitidae	J (Plie)	— R	
Buliminidae	T (Dani)	— R	
Calcarinidae	K (Turo)	— R	
Caligellidae	D (Fras)	— C (Tour)	
Cassidulinidae	K (Camp)	— R	
Caucasinidae	K (Turo)	— R	
Ceratobuliminidae	Tr (Carn)	— R	
Cibicididae	K (Haut)	— R	
Colaniellidae	D (Fras)	— P (Dzhu)	
Cymbaloporidae	Tr (Ladi)	— R	
Dicyclinidae	J (Sine)	— T (Eo-m)	
Discocyclinidae	T (Pale)	— T (Eo-u)	
Discorbidae	Tr (Ladi)	— R	
Elphidiidae	K (Maes)	— R	
Endothyridae	D (Give)	— P (Dzhu)	
Eostaffellidae	C (Vise)	— C (Step)	
Eouvigerinidae	K (Albi)	— R	
Epistomariidae	K (Sant)	— R	
Eponididae	T (Pale)	— R	
Fischerinidae	S (u)	— R	
Fusulinidae	C (Namu)	— P (Guad)	

Glabratellidae	T (Eo-u)	— R
Glandulinidae	J (Hett)	— R
Globigerinidae	K (Maes)	— R
Globorotaliidae	T (Pale)	— R
Globotruncanidae	K (Turo)	— K (Maes)
Hantkeninidae	T (Pale)	— R
Heterohelicidae	J (Bajo)	— T (Ol)
Homotrematidae	K (Sant)	— R
Hormosinidae	C (Tour)	— R
Involutinidae	Tr (Ladi)	— K (Turo)
Islandiellidae	T (Pale)	— R
Lasiodiscidae	C (Vise)	— P (Dzhu)
Lepidocylinidae	T (Eo-m)	— T (Mi-m)
Lituolidae	C (Tour)	— R
Loxostomidae	K (Coni)	— Q (Plei)
Miliolidae	J (Hett)	— R
Miogypsinidae	T (Ol-l)	— T (Mi-m)
Moravamminidae	Θ	— C (Mosc)
Nodosariidae	P (Guad)	— R
Nodosinellidae	S (l)	— P (Dzhu)
Nonionidae	J (Bajo)	— R
Nubeculariidae	C (Namu)	— R
Nummulitidae	K (Maes)	— R
Orbitoididae	K (Camp)	— T (Pale)
Orbitolinidae	K (Barr)	— T (Eo-u)
Osangulariidae	K (Barr)	— R
Ozawainellidae	P (l)	— P (Dzhu)
Palaeotextulariidae	C (Tour)	— P (Dzhu)
Parathuramminidae	D (Give)	— C (Tour)
Pavonitinidae	J (Bath)	— R
Planomaliniidae	K (Apti)	— T (Dani)
Planorbulinidae	T (Pale)	— R
Pleurostomellidae	K (Albi)	— R
Polymorphinidae	Tr (Carn)	— R
Pseudorbitoididae	K (Ceno)	— K (Maes)
Ptychocladidae	K (Fras)	— P (Sakm)
Robertinidae	K (Camp)	— R
Rotaliidae	K (Coni)	— R
Rotaliporidae	K (Haut)	— K (Maes)
Rzehakinidae	K (Berr)	— R
Saccamminidae	Θ (Aren)	— R
Schackoinidae	K (Apti)	— K (Maes)
Schizamminidae	Tr (Rhae)	— R
Schubertellidae	C (Namu)	— P (Dzhu)
Schwagerinidae	C (Step)	— P (Guad)
Semitemilaridiidae	D (Give)	— D (Fras)
Siphoninidae	T (Pale)	— R

(361a)

Soritidae	Tr (Rhae)	— R
Sphaeroidinidae	K (Camp)	— R
Spirillinidae	C (Vise)	— R
Staffellidae	C (Bash)	— P (Dzhu)
Tetrataxidae	C (Tour)	— Tr (Rhae)
Textulariidae	C (Tour)	— R
Tournayellidae	D (Give)	— P (Dzhu)
Trochamminidae	S (m)	— R
Turrilinidae	J (Bajo)	— R
Uvigerinidae	K (Turo)	— R
Verbeekinidae	P (Sakm)	— P (Dzhu)

## Cl. Radiolaria

[Classification and stratigraphic ranges from Kling (1978), the *Treatise*, Pt. D, and *The Fossil Record*; 18 extant marine families without known fossil representatives are not listed.]

Or. Tripylea		
Challengeriidae	T (Mi-m)	— R
Or. Spumellaria		
Actinommidae	Tr	— R
Coccodiscidae	K (Maes)	— T (Ol-l)
Collosphaeridae	T (Mi-l)	— R
Entactinidae	Є (m)	— C (Namu) (217)
Litheliidae	C (Bash)	— R
Orosphaeridae	T (Eo)	— R
Phacodiscidae	C (Vise)	— R
Pseudoaulophacidae	K (Coni)	— K (Camp)
Pyloniidae	T (Eo)	— R
Spongodiscidae	D (Give)	— R
Tholoniidae	T (Plio)	— R
Or. Nassellaria		
Acanthodesmiidae	T (Pale)	— R
Amphipyndaciidae	K (Apti)	— K (Maes)
Artostrobiidae	K (Camp)	— R
Cannabotrydidae	T (Eo-u)	— R
Carpocaniidae	T (Eo-u)	— R
Plagoniidae	K (Sant)	— R
Pterocorythidae	T (Eo-m)	— R
Rotaformidae	K	
Theoperidae	Tr (Anis)	— R
Or. Incertae sedis		
Albaillellidae	S	— C (Namu)
Palaeoscenidiidae	D (Fame)	— R

## Cl. Acantharia

[Classification and stratigraphic ranges from the *Treatise*, Pt. D; 13 extant marine families without known fossil representatives are not listed.]

Acanthochiasmataidae	T (Eo)	— R	
Astrolonchidae	T (Mi)	— R	

## Cl. Ciliata

[Classification from Tappan and Loeblich (1968) and the *Treatise*, Pt. D; stratigraphic ranges from *The Fossil Record*. More than 27 extant marine families without known fossil representatives are not listed.]

Or. Spirotrichia (including Tintinnina)			
Codonellidae	Θ (Ashg)	— R	(181)
Codonellopsidae	J (Tith)	— R	
Coxliellidae	K (Berr)	— R	
Heterotrichina	T (Dani)	— R	(181)
Hypotrichina	T (Eo)	— R	(181)
Xystonellidae	J (u)	— R	

## Cl. Incertae sedis

Or. Calpionellida			
Calpionellidae	J (Tith)	— K (Vala)	(258)
Colomiellidae	K (Apti)	— K (Albi)	(258)

?Or. Chitinozoa			
Ancyrochitinidae	Θ (Aren)	— D (Fame)	(344)
Conochitinidae	Θ (Aren)	— D (Fame)	(344)
Desmochitinidae	Θ (Aren)	— C (Tour)	(344)
Hoegispharidae	Θ (Aren)	— D (Fras)	(344)
Lagenochitinidae	Θ (Aren)	— D (Fame)	(344)
?incertae sedis	u Riph		(28)

?Or. Reitlingerillida			
Reitlingellidae	€ (l)	— Θ (Cara)	(208)

## PORIFERA

[Taxa listed below are based principally on the somewhat antiquated but comprehensive classification of DeLaubenfels in the *Treatise*, Pt. E., with numerous additions and emendations.]

## Cl. Demospongia

[Classification and stratigraphic ranges from the *Treatise*, Pt. E., with emendations from *The Fossil Record* and other sources, as listed; approximately 13 extant families without known fossil representatives are not listed.]

Or. Monaxonida			
Acarniidae	K	— R	
Amphilectidae	Q (Plei)	— R	

Choiidae	Є (mMid)	— C (Mosc)	(272)
Cladorhizidae	Q (Plei)	— R	
Clionidae	S	— R	
Corralioiidae	Є		
Desmacidontidae	Q (Plei)	— R	
Dysideidae	J	— R	
Haliclonidae	Θ (l)	— R	
Hamptoniidae	Є (mMid)		
Hazeliiidae	Є (mMid)		
Heliospongidae	C (u)	— P (Leon)	(94)
Myxillidae	K	— R	(229)
Piraniidae	Є (mMid)	— S	
Sollasellidae	Θ (m)	— R	
Spirastrellidae	Q (Plei)	— R	
Spongiidae	C	— T (Eo)	
Suberitidae	K (l)	— R	
Takakkawiidae	Є (mMid)		
Tedaniidae	Q (Plei)	— R	
Wapkiidae	Є (mMid)		

## Or. Choristida (= Astrophorida)

Ancorinidae	J	— R
Craniellidae	K (u)	— R
Geodiidae	C (Vise)	— R
Halinidae	C (u)	— R
Ophiraphiditidae	J (u)	— R
Scolioraphididae	J	— K (u)

## Or. Lithistida

?Anomoclonellidae	Θ (m)	— S (m)
Anthaspidellidae	Θ (Aren)	— P (Dzhu)
Anthracosyconidae	P (Leon)	— P (Guad) (94)
Astroboliidae	J	— T (Mi)
Astrocladiidae	P (Sakm)	— T
Astylospongiidae	Θ (Cara)	— P (l) (267)
Aulaxiniidae	K (l)	— K (u)
Chenendoporidae	J (m)	— K (Seno) (213)
Chiastoclonellidae	S (m)	— P (Guad)
Chonellidae	J (u)	— T (Mi)
Cnemidiastridae	C (l)	— J (u)
Cylindrophymatidae	J (Bajo)	— J (Kimm) (132)
Discodermiidae	K (l)	— R
Dorydermatidae	C (l)	— K (Seno)
Gignouxiidae	J (Oxfo)	— R
Hallirhoiidae	J (m)	— T
Haplistiidae	Θ (m)	— P (Leon) (273)
Helobrachiidae	K (u)	
Hindiidae	Θ (Llde)	— P (Dzhu) (95)

Isoraphiniidae	J	— K (Maes)	
Jereicidae	J (u)	— R	
Jereidae	P (Leon)	— K (u)	(94)
Kaliapsidae	K (Turo)	— R	(142)
Leiodorellidae	J (Oxfo)	— R	(213)
Microspongiidae	Θ (Cara)	— P (Dzhu)	
Neopeltidae	K (u)	— R	
Phymaraphiniidae	K (Ceno)	— T	
Pleromidae	J (u)	— R	
Plinthodermatiidae	K (u)		
Plinthosellidae	K (Ceno)	— K (Maes)	
Pseudoverruculinidae	K (u)		
Roanespongiidae	P (Leon)		(273)
?Saccospongiidae	Θ (Cara)	— S (u)	
Scleritodermatidae	K	— R	
Scytraliidae	J	— K (Camp)	(142)
?Vauxiidae	Є (uMid)		(270)
Vetulinidae	J (Oxfo)	— R	(94)

## Cl. Calcarea (= Calcispongia)

[Classification and stratigraphic ranges principally from the *Treatise*, Pt. E, and *The Fossil Record*; 9 extant families without known fossil representatives are not listed.]

## Or. Heteractinida

Astraeospongiidae	Θ (Cara)	— D (Give)	(269)
Wewokellidae	C (Tour)	— P (Leon)	(269)

## Or. Calcaronea (= Lebetida, Syconosa)

Grantiidae	J (u)	— R	
Leuconiidae	C (Vise)	— R	

## Or. Pharetronida

Bactronellidae	J (u)	— T (Eo)	
Discocoeliidae	Tr (m)	— K (Maes)	(133)
Elasmocoeliidae	J (Oxfo)	— K (l)	(142)
Elasmostomatidae	P (Leon)	— K (Ceno)	(101)
Lelapiidae	P (Sakm)	— R	
Pharetrospongiidae	K (l)	— K (Maes)	
Porosphaeridae	K (l)	— R	
Sestrostomellidae	Tr (m)	— R	
Stellispongiidae	P (Sakm)	— K (Ceno)	(96)

## Or. Sphinctozoa (= Thalamida)

Celyphiidae	Θ (Ashg)	— Tr (Carn)	(271)
Cryptocoeliidae	Tr (Anis)	— K (Maes)	(133)
Cystothalamiidae	Θ (Ashg)	— Tr (Carn)	(96,112,271)
Fissispongiidae	C (Mosc)	— P (Leon)	(96)
Guadalupiidae	P (Sakm)	— P (Guad)	(95,133)

Maendrostiidae	C (Mosc)	— C (Step)	(96)
Phragmocoeliidae	Tr (Carn)		
Polytholosiidae	Tr (Anis)	— Tr (Carn)	(112,133)
Sebargasiidae	Θ (Ashg)	— Tr (u)	(271)
Sphaerocoeliidae	J (u)	— K (Ceno)	(133)
Thaumastocoeliidae	C (u)	— Tr (Carn)	(133)
Verticillitidae	P (Asse)	— K (Maes)	(133)

## Cl. Hexactinellida (= Triaxonida, Hyalospongea)

[Classification and stratigraphic ranges based on the *Treatise*, Pt. E, with emendations from *The Fossil Record* and other sources, as listed. Two extant families without known fossil representatives are not listed.]

## Or. Hexactinosida (= Dictyida)

Aphrocallistidae	K (Coni)	— R	(142)
Botryosellidae	J (u)	— K (Sant)	(229,256)
Craticulariidae	K (Apti)	— R	(142)
Dactylocalycidiae	K (Sant)	— R	(229)
Emploidae	J (m)		
Eubrochididae	K (Albi)	— K (Maes)	(256)
Euretidae	J (u)	— R	
Hexactinellidae	J	— R	
Leptophragmatidae	J (u)	— T (Mi)	
Myliusiidae	J (m)	— R	
Pleurostomatidae	Tr (Anis)	— T (Eo)	
Polystigmatiidae	K (u)		
Polythyrididae	K (u)		
Porospongiidae	J (u)	— K (u)	
Staurodermatidae	Tr (Anis)	— T (Mi)	
Wapkiosidae	K (u)	— R	

## Or. Lychniscosida (= Lychniskida)

Becksiidae	J	— K (Maes)	(256)
Callodictyidae	K (Albi)	— K (Maes)	(256)
Calyptrellidae	J	— K (Ceno)	(256)
Camerospongidae	J (u)	— R	
Coeloptychiidae	K (Coni)	— K (Maes)	(142,229)
Coeloscyphiidae	K (u)		
Coscinoporidae	K (Sant)	— K (Maes)	(256)
Cypelliidae	J (u)	— R	(229,256)
Oncotoechidae	K (u)		
Polyblastidiidae	J	— K (u)	
Sporadosciniidae	K (Camp)		(142)
Ventriculitidae	J (u)	— T (Mi)	(142)
incertae sedis	Tr (Carn)		

## Or. Lyssacinosida (= Lyssakida)

Brachiospongidae	Θ (Cara)	— S (u)	(269)
------------------	----------	---------	-------

Caulophacidae	T (Eo)	— R
Dictyospongiidae	S (I)	— P (Leon)
?Dierespongiidae	€ (uMid)	— Θ (m) (269)
Docodermatidae	C (Mosc)	— J (u) (133)
Euplectellidae	J (I)	— R (133)
Malumispongidae	S	— (133)
Mattaspongiidae	D (Fras)	(269)
Multivasculatidae	€ (Fran)	— € (Trep) (223)
Pelicaspongiidae	Θ (Ashg)	— D (Fras) (267,268)
Pileolitidae	P (Sakm)	— P (Leon) (96)
Protospongiidae	€ (ITom)	— S (I) (269,395)
Rossellidae	T (Eo)	— R
Stauractinellidae	J (Call)	— J (Oxfo)
Stereodictyiidae	C (Step)	— P (Leon) (94)
Stiodermatidae	C (Vise)	— P (Guad) (94)
Stromatidiidae	P (Guad)	(94,96)
Teganiidae	€ (uMid)	— P (Leon) (269)
Titusvilliidae	D (Fras)	— C (Tour)

## Or. Amphidiscophora

Hyalonematidae	C (I)	— R
Pheronematidae	K (Camp)	— R

## Cl. Sclerospongiae

## ?Or. Stromatoporoidea

[Classification based on Stearn (1980) with additions from the *Treatise*, Pt. F; stratigraphic ranges from Stearn (1980), Bogoyavlenskaya (1974), and the *Treatise*.]

Actinostromatidae	S (I)	— D (Fame)
Amphiporidae	D (Sieg)	— D (Fras)
Aulaceridae	Θ (Cara)	— S (I)
Clathrodictyidae	Θ (Llde)	— D (Fame)
Cliefdenellidae	Θ (Cara)	— Θ (Ashg) (370)
Densastromatidae	S (I)	— D (Gedi)
Diplostromatidae	S (u)	(187)
?Disjectoporidae	P (Guad)	— Tr (Carn) (112)
Ecclimadictyidae	Θ (Cara)	— D (Fame) (370)
Hermatostromatidae	S (u)	— D (Fras)
Labechiidae	Θ (Cara)	— D (Fame)
Lophiostromatidae	Θ	— S
?Milleporellidae	J (u)	— K (Maes)
?Milleporidiidae	J (u)	— K (Maes)
Pseudolabechiidae	S (I)	— S (u)
Rosenellidae	Θ (m)	— D (Fame)
Stachyoditidae	D (Eife)	— D (Fras)
Stictostromatidae	Θ (Cara)	— D (Fame)

Stromatoporidae	S (m)	— K (u)	
Syringostromatidae	S (u)	— J (Tith)	
Syringostromellidae	S (m)	— D (Fras)	
Tienodictyidae	S (l)	— D (Fras)	(187)
?incertae sedis	€ (Atda)		(310)
Or. Ceratoporellida			
Ceratoporellidae	P	— R	(133)
Or. Tabulospongida			
Acanthochaetidae	J (Oxfo)	— T (Eo-m)	(208)
Chaetetidae	Θ (Ashg)	— Tr (Carn)	(208)
?Cryptolichenariidae	Θ (Trem)	— Θ (Ashg)	(208)
?Desmidoporidae	Θ (Ashg)	— D (Eife)	(208)
?Lamottiidae	Θ (Llde)		(208)
Tabulospongidae	J (Oxfo)	— R	(133)
?Tiverinidae	S (m)	— D (l)	(208)
Or. Muranida			
Muranidae	K (Haut)		(161)

## ARACHAEOCYATHA

[Classification and stratigraphic ranges based largely on the *Treatise*, Pt. El (revised).]

## Cl. Regulares

Or. Monocyathida			
Capsolynthidae	€ (Atda)	— € (Boto)	
Capsulocyathidae	€ (uTom)	— € (Boto)	(288)
Cryptoporocyathidae	€ (lTom)	— € (Atda)	(288)
Ethmolynthidae	€ (Atda)	— € (Boto)	
Fransuasaecyathidae	€ (uTom)	— € (Boto)	(178,288)
Globosocyathidae	€ (Boto)		
Monocyathidae	€ (lTom)	— € (lMid)	(288)
Rhabdocyathellidae	€ (Atda)	— € (mMid)	(288,362)
Tumuliolynthidae	€ (uTom)	— € (Boto)	(288)
Uralocyathellidae	€ (Boto)		(288)
Uralocyathidae	€ (uTom)	— € (Boto)	
Or. Putapacyathida			
Aptocyathidae	€ (Atda)	— € (Boto)	(288)
Gerbicanicyathidae	€ (Boto)		
Putapacyathidae	€ (Atda)		
Or. Ajacicyathida			
Acanthinocyathidae	€ (Boto)		
Agyrekocyathidae	€ (lMid)		
Ajacicyathidae	€ (lTom)	— € (lMid)	(76,288)
Alataucyathidae	€ (Atda)	— € (Boto)	(288)

Anaptyctocyathidae	€ (Atda)	— € (Boto)	
Annulocyathidae	€ (Boto)	— € (lMid)	(288)
Boscecuclyathidae	€ (Boto)	— € (Boto)	(178,288)
Botomocyathidae	€ (Atda)	— € (Boto)	(288)
Bronchocyathidae	€ (uTom)	— € (Boto)	(288)
Calyptocosciniidae	€ (Atda)		
Carinacyathidae	€ (uTom)	— € (Boto)	(288)
Chankacyathidae	€ (Boto)		
Clathricosciniidae	€ (Atda)	— € (lMid)	(288)
Compositocyathidae	€ (uTom)	— € (Boto)	
Coscinocyathellidae	€ (Boto)		(288)
Coscinocyathidae	€ (uTom)	— € (Boto)	(76,178,288)
Cyclocyathellidae	€ (Atda)	— € (Boto)	(76,288)
Dokidocyathidae	€ (lTom)	— € (Boto)	
Erbocyathidae	€ (Atda)	— € (lMid)	(178,288)
Ethmocoscinidae	€ (Atda)	— € (Boto)	
Ethmocyathidae	€ (Boto)		
Ethmopectinidae	€ (Boto)		
Ethmophyllidae	€ (Atda)	— € (lMid)	(288)
Fallocyathidae	€ (Atda)		
Fansicyathidae	€ (Atda)		
Formosocyathidae	€ (uTom)	— € (Boto)	
Geocyathidae	€ (uTom)	— € (Boto)	
Gloriosocyathidae	€ (Atda)	— € (Boto)	
Hupecyathellidae	€ (Boto)		(288)
Hyptocyathidae	€ (Boto)		(180)
Kaltatocyathidae	€ (uTom)	— € (Boto)	(288)
Kasyricyathidae	€ (Atda)	— € (Boto)	(178,288)
Kidrjasocyathidae	€ (Atda)	— € (Boto)	(288)
Kijacyathidae	€ (Atda)	— € (Boto)	(288)
Kordecyathidae	€ (Atda)	— € (Boto)	(288)
Lenocyathidae	€ (Atda)	— € (Boto)	(288)
Mrassucyathidae	€ (Atda)	— € (Boto)	(288)
Nochoroicyathidae	€ (lTom)	— € (lMid)	
Peregriniccyathidae	€ (Boto)		(288)
Piamaecyathidae	€ (Boto)		(288)
Polycosciniidae	€ (Atda)	— € (Boto)	
Porocoscinidae	€ (Boto)		(180)
Porocyathidae	€ (Atda)	— € (Boto)	(288)
Pretiosocyathidae	€ (Atda)	— € (Boto)	(288)
Robertocyathidae	€ (Atda)	— € (Boto)	
Robustocyathidae	€ (lTom)	— € (Boto)	
Rozanovicyathidae	€ (Boto)		
Sajanocyathidae	€ (Boto)	— € (lMid)	
Sanarkocyathidae	€ (Boto)		
Schidertycyathidae	€ (Boto)	— € (lMid)	
Sigmocoscinidae	€ (Boto)		

Sigmocyathidae	€ (Boto)		
Soanicyathidae	€ (Boto)	— € (Boto)	(288)
Stillicidocyathidae	€ (Atda)	— € (Boto)	(288)
Tannuolacyathidae	€ (Boto)		
Tennericyathidae	€ (Atda)	— € (Boto)	(288)
Tercyathidae	€ (Boto)		(288)
Tumulocyathidae	€ (Atda)	— € (Boto)	(288)
Vologdinocyathidae	€ (Atda)	— € (lMid)	(288)
Wrighticyathidae	€ (Boto)		(180)

## Cl. Irregulares

Or. Thalassocyathida (= Rhizacyathida)			
Bacatocyathidae	€ (uTom)	— € (mMid)	(362)

Or. Archaeocyathida			
Anthomorphidae	€ (Atda)	— € (lMid)	
Archaeocyathidae	€ (Boto)	— € (mMid)	
Archaeofungiidae	€ (Atda)	— € (Boto)	(288)
Archaeopharetridae	€ (l)		
Archaeosyconidae	€ (uTom)	— € (lMid)	
Bicyathidae	€ (uTom)	— € (Boto)	
Copleicyathidae	€ (l)		
Dictyocoscinidae	€ (Boto)		
Dictyocyathidae	€ (uTom)	— € (Boto)	(76)
Flindersicyathidae	€ (Boto)	— € (lMid)	
Metacoscinidae	€ (Boto)	— € (lMid)	
Metacyathidae	€ (uTom)	— € (lMid)	(288)
Prismocyathidae	€ (Boto)		
Protocyclocyathidae	€ (l)		
Protopharetridae	€ (uTom)	— € (lMid)	(76)
Pycnoidocoscinidae	€ (l)		
Sigmofungiidae	€ (Boto)		
Tabellaecyathidae	€ (Boto)	— € (lMid)	
Tabulacyathidae	€ (Atda)	— € (Boto)	

Or. Syringocnemidida			
Syringocnemididae	€ (Boto)		(288)
Syringocoscinidae	€ (l)		

## Cl. Incertae sedis

Or. Kazakhstanicyathida			
Kazakhstanicyathidae	€ (Boto)	— € (lMid)	

Or. Archaeophyllida			
Archaeophyllidae	€ (Atda)		

## Insertae sedis

Acanthopyrgidae	€ (Boto)		
Korovinellidae	€ (Boto)		

Manacyathidae	Є (Atda)
Mattewcyathidae	Є (m)

## COELENTERATA

## Cl. "Protomedusae" (= Brooksellida)

[Artificial taxon incorporating Vendian and lower Paleozoic medusoids of uncertain taxonomic affinities; includes approximately 22 "Ediacaran" genera (including all listed as "medusae of uncertain affinities" by Glaessner in the *Treatise*, Pt. A), as well as other medusae from the Cambrian, Ordovician, and Silurian.]

incertae sedis	V (m)	— S (l)	(132,155,231,327)
----------------	-------	---------	-------------------

## Cl. "Petalonamae"

[The "petal organisms" of Pflug (1970-1972), considered pinnatulaceans by Glaessner in various pre-1975 publications. Classification follows the *Treatise*, Pt. A.]

?Charniidae	V (m)	— V (u)	(162,328)
Erniettidae	V (u)		(114,241)
?Pteridiniidae	V (m)	— V (u)	(162,163,231,241)
?Rangeidae	V (m)	— V (u)	(122,124,241,328)
incertae sedis <i>(Arumberia, Baikalina, Inkrylovia, Namalia, Nasepia)</i>	V (m)	— V (N-Da)	(114,124,231,328)

## Cl. Hydrozoa

[Basic classification and stratigraphic ranges follow the *Treatise*, Pt. F, and *The Fossil Record*; approximately 25 extant marine families without known fossil representatives are not listed.]

## Or. Trachylinida

Cuninidae	J (Tith)	— R
?Kirklandiidae	J (Bajo)	— T (Pale)
?Velumbrella	Є (m)	(310)

## Or. Hydroida

Calyptoblastina	Є (mMid)	— R
Hydractiniidae	J (Tith)	— R
Oceaniidae	C (Step)	— R
Sertulariidae	K (Maes)	— R (360)

## Or. Siphonophorida

Chondroplidae	V (u)	(364,365)
Porpitidae	V (u)	— R (118,365)
Velellidae	D (Gedi)	— R (118,310)

## Or. Spongimorphida

Spongimorphidae	Tr (Olen)	— J (Tith)
-----------------	-----------	------------

Or. Milleporina			
Axoporidae	T (Eo)	— T (Ol)	
Milleporidae	T (Dani)	— R	
Or. Styelasterina			
Styelasteridae	T (Dani)	— R	
Or. Sphaeractinida			
Heterastridiidae	Tr (m)	— Tr (Nori)	
Sphaeractinidae	C (Step)	— K (Ceno)	
Incertae sedis			
<i>Beltanelliformis</i>	V (u)		(231)
<i>Nemiana</i>	V (u)	— V (N-Da)	(231)
<i>Tirasiaria</i>	V (u)		(231)

## Cl. Scyphozoa

[Basic classification and stratigraphic ranges follow the *Treatise*, Pt. F, and *The Fossil Record*; approximately 13 extant families (including all in the Stauro-medusida) without known fossil representatives are not listed.]

Or. Carybdeida			
Carybdeidae	C (Mosc)	— R	(157)
Or. Coronatida			
? <i>Byronia</i>	Є (mMid)		(310)
Collaspidae	Θ (Trem)	— R	
Periphyllidae	J (Tith)	— R	
Or. Semaeostomatida			
* <i>Eulithotidae</i>	J (Tith)		
* <i>Semaeostomidae</i>	J (Tith)		
incertae sedis	C (Mosc)		(105)
Or. Lithorhizostomatida			
Rhizostomidae	J (Tith)		
Or. Rhizostomatida			
Leptobrachiidae	J (Tith)	— R	
incertae sedis	C (Mosc)		(105)
incertae sedis	C (Step)		(230)
?Or. Conulariida			
<i>Circonulariidae</i>	S (l)	— S (u)	(25)
<i>Conchopeltidae</i>	V (m)	— Θ (Cara)	(118,329)
<i>Conulariellidae</i>	Є (mMid)	— Θ (Aren)	
<i>Conulariidae</i>	Є (u)	— Tr (Rhae)	
<i>Conulariopsidae</i>	Tr (Olen)		
? <i>Palaearacmaeidae</i>	Є (Trep)	— Θ (Trem)	(292,391a)
<i>Paleoconulariidae</i>	Є (Boto)		(395)

\*May be related to unlisted extant families.

Or. Septodaearia			
? <i>Septodaeum</i>	Θ (Aren)	— D (Sieg)	(24)
Incertae sedis			
? <i>Albumares</i>	V (u)		(231)
? <i>Brachina</i>	V (u)		(365)
<i>Hallidaya, Skinnera</i>	V (u)		(124,363)
? <i>Kimberella</i>	V (u)		(365)
Cl. Hydroconozoa			
3 families	Є (Boto)		(395)
Cl. Anthozoa			
[Classification and stratigraphic ranges based on the <i>Treatise</i> , Pt. F (including 1981 supplement), and <i>The Fossil Record</i> , except where noted otherwise; approximately 70 extant families (including all in the Ceriantaria, Telestacea, Zoanthiniaria, and Corallimorpharia) without known fossil representatives are not listed.]			
Or. Antipatharia			
* <i>Leiopathes</i>	T (Mi)		
Or. Stolonifera			
Clavulariidae	K (Berr)	— R	
Tubiporidae	T (Mi)	— R	(229)
Or. Alcyonacea			
Alcyoniidae	S (m)	— R	(17)
Nephtheidae	J (Hett)	— R	
Xeniidae	K (u)	— R	(229)
?incertae sedis	Є (Boto)		(38)
Or. Trachypsammiaeae			
Trachypsammidae	P (Guad)		
Or. Coenothecalia			
Helioporidae	K (Berr)	— R	
Or. Gorgonacea			
? <i>Pragnellia</i>	Θ (Ashg)		(310)
Coralliidae	K (Berr)	— R	
Ellisellidae	T (Dani)	— R	
Gorgoniidae	T (Eo-m)	— R	(116)
Isididae	K (Maes)	— R	(229)
Melithaeidae	T (Eo)	— R	(229)
Parisididae	T (Eo)	— R	(229)
Primnoidae	K (u)	— R	
Or. Pennatulacea			
Pennatulidae	Tr	— R	(229)

\*May be related to unlisted extant families.

Pteroeididae	T (Eo)	— R
Virgulariidae	K (Berr)	— R
Or. Tabulata		
Agetolitidae	Θ (Ashg)	— S (m)
Alveolitidae	S (l)	— D (Fras)
Aulocystidae	S (l)	— P (u)
Auloheliidae	P (Dzhu)	
Auloporidae	Θ (Aren)	— P (Dzhu) (310)
Bajgoliidae	Θ (m)	— Θ (Ashg)
Billingsariidae	Θ (Llde)	— S (l)
Chonostegitidae	D (Emsi)	— D (Eife)
Cleistoporidae	S (u)	— C (Bash)
Coccoserididae	Θ (Cara)	— Θ (Ashg) (310)
Coenitidae	S (l)	— D (Give)
Cyrtophyllidae	Θ (Cara)	— Θ (Ashg) (310)
Favositidae	Θ (Cara)	— P (Guad) (310)
Fletcheriellidae	Θ (Ashg)	— D (m)
Gorskytidae	C (Tour)	— P (l)
Halysitidae	Θ (Cara)	— S (u)
Heliolitidae	Θ (Llde)	— D (Give)
Khmeriidae	C (Bash)	— P
Kozlowskiocystiidae	D (Eife)	
Lichenariidae	Θ (Cara)	(310)
?Lipoporidae	€ (IMid)	— € (mMid) (153)
Michelinidae	S (u)	— P (Dzhu)
Multisoleniidae	Θ (Ashg)	— S (u)
Multithecoporidae	S (l)	— P (u)
Pachyporidae	S (l)	— P (Dzhu)
Palaeacidae	C (Tour)	— C (Vise)
Palaeoporitidae	Θ (Cara)	— Θ (Ashg) (185)
Paleoalveolitidae	Θ (m)	
Parastriatoporidae	Θ (Ashg)	— D (Eife)
Periphaceloporidae	D (Give)	
Plasmoporellidae	Θ (m)	— S (m)
Plasmoporidae	S (l)	— D (Eife)
Proheliolitidae	Θ (Cara)	— S (u)
Proporidae	Θ (Cara)	— S (u)
Pseudofavositidae	P (u)	
Pseudoplasmoporidae	Θ (Ashg)	— D (Gedi)
Pycnolithidae	S (l)	
Pyrigiidae	S (u)	— P (Dzhu) (185)
Roemeridae	D (Emsi)	— P (Sakm)
Romingeridae	S (m)	— C (Namu)
Sibiriolitidae	Θ (Ashg)	
Sinoporidae	S (l)	— P (Leon)
Stelliporellidae	Θ (Ashg)	— D (Eife)

Syringolitidae	S (l)	— D (l)	
Syringophyllidae	Θ (Llde)	— S (l)	(310)
Syringoporidae	Θ (Cara)	— P (Guad)	(222)
Taeniolitidae	Θ (Ashg)		
Tetradiidae	Θ (Livi)	— Θ (Ashg)	(310)
Tetraporellidae	Θ (m)	— P (Dzhu)	
Theciidae	S (l)	— D (Eife)	
Thecostegitidae	D (Gedi)	— C (Namu)	
Trachypsammidiidae	P (Asse)	— P (Dzhu)	
Vaughaniidae	C (Tour)		
 Or. Rugosa			
Acervulariidae	S (m)	— S (u)	
Acrophyllidae	D (Emsi)		(225)
Adamantanophyllidae	C (Tour)	— C (Namu)	
Amplexidae	S (l)	— C (Vise)	
Amsdenoididae	S (u)		
Anisophyllidae	S (u)		
Antiphyllidae	C (Vise)	— C (Mosc)	
Aphrophyllidae	C (Tour)	— C (Vise)	
Arachnophyllidae	S (l)	— D (Sieg)	
Aulophyllidae	C (Tour)	— P (l)	
Axophyllidae	D (Vise)	— C (Namu)	
Bethanyphyllidae	D (Give)		
Bothrophyllidae	C (Vise)	— P (l)	
Calostylidae	Θ (Cara)	— D (Gedi)	(225,310)
Centristelidae	D (Give)		
Columnariidae	D (Gedi)	— D (Give)	
Combophyllidae	D (Emsi)	— D (Eife)	(225)
?Cothoniidae	€ (lMid)		(153)
Cyathaxoniidae	D (Fame)	— P (Guad)	
Cyathophyllidae	D (Sieg)	— D (Fras)	(225)
Cyathopsidae	C (Tour)	— P (l)	
Cystiphyllidae	Θ (Cara)	— D (Give)	(225,310)
Digonophyllidae	D (Eife)	— D (Give)	
Disphyllidae	S (u)	— D (Fame)	
Ditoecholasmatidae	S (u)		
Durhaminidae	C (Mosc)	— P (Guad)	(222)
Edridophyllidae	S (u)	— D (Give)	(225)
Ektasophyllidae	C (Vise)		
Endamplexidae	P (Guad)		
Endophyllidae	S (m)	— D (Fras)	(225)
Endotheciidae	P (u)		
Entelophyllidae	S (l)	— D (Gedi)	
Expressophyllidae	S (m)	— S (u)	
Fasciphyllidae	D (Emsi)	— D (Give)	
Fletcheriidae	S (l)	— S (m)	

Geyerophyllidae	C (Namu)	— P (Leon)	
Goniophyllidae	S (l)	— D (Give)	(208)
Hadrophyllidae	D (Sieg)	— D (Eife)	(225)
Halliidae	D (Sieg)	— D (Give)	(225)
Hapsiphyllidae	D (Emsi)	— P (Dzhu)	(225)
Holmophyllidae	S (m)	— D (Eife)	
Ketophyllidae	S (l)	— S (u)	
Kielcephyllidae	D (Fame)		
Kiziliidae	C (Vise)		
Kodonophyllidae	S (l)	— D (Eife)	
Koninkocariniidae	C (Mosc)		
Kyphophyllidae	Θ (Ashg)	— D (Gedi)	
Laccophyllidae	S (l)	— P (Guad)	
Lambelasmatidae	Θ (Cara)	— S (l)	
Lindstroemiidae	D (m)		
Lithostrotionidae	C (Tour)	— P (Leon)	
Lophophyllidae	C (Tour)	— P (Dzhu)	(100)
Lykophyllidae	S (l)	— S (u)	
Metriophyllidae	D (l)	— D (Fras)	
Mucophyllidae	S (l)	— D (Emsi)	
Neocolumnariidae	D (Eife)		
?Numidiaphyllidae	P (Guad)		(100)
Palaeocyclidae	Θ (m)	— D (Eife)	
Palaeosmiliidae	C (Vise)	— C (Namu)	
Paliphyllidae	Θ (Cara)	— S (m)	(185)
Pentaphyllidae	D (Emsi)	— P (l)	(225)
Petalaxidae	C (Vise)	— P (l)	
Petraiidiae	Θ (m)	— D (Fame)	
Phillipsastreidae	D (Emsi)	— D (Fras)	(225)
Plerophyllidae	D (Emsi)	— P (u)	
Polycoeliidae	S (u)	— P (Dzhu)	(100,225)
Pseudopavonidae	C (Namu)		
Ptenophyllidae	S (m)	— D (Fras)	
Ptychophyllidae	S (l)	— S (u)	
Pycnostylidae	S (l)	— D (Give)	
Spongophyllidae	S (u)	— D (Give)	
Stauriidae	Θ (Cara)	— D (Give)	(208)
Stereolasmatidae	D (Give)	— C (Mosc)	
Streptelasmatidae	Θ (Cara)	— D (Give)	
Stringophyllidae	D (Gedi)	— D (Give)	(208,225)
Timorphyllidae	P (Dzhu)		
Tryplasmatidae	Θ (Cara)	— D (Give)	(225)
Uraliniidae	C (Tour)	— C (Vise)	(229)
Verbeekiellidae	C (Vise)	— P (Leon)	
Waagenophyllidae	C (Mosc)	— P (Dzhu)	(100)
Zaphrentidae	D (Emsi)	— D (Give)	
Zaphrentoididae	C (Vise)	— C (Bash)	

## Or. Heterocorallia

Heterophyllidae D (Eife) — C (Namu)

## Or. Scleractinia

[Classification from the *Treatise*, Pt. F; stratigraphic ranges based on *The Fossil Record*.]

Acroporidae	K (Turo)	— R
Actinacididae	J (Bath)	— T (Ol-l)
Agariciidae	K (Apti)	— R
Amphastreidae	J (Bath)	— K (Ceno)
Astrocoeniidae	Tr (Anis)	— R
Calamophyllidae	J (Plie)	— K (Ceno)
Caryophyllidae	J (Hett)	— R
Cyclolitidae	K (Haut)	— T (Mi-m)
Dendrophylliidae	K (Ceno)	— R
Faviidae	J (Bajo)	— R
Flabellidae	K (Albi)	— R
Fungiidae	K (Apti)	— R
Guyniidae	J (Oxfo)	— R
Haplaraeidae	J (Kimm)	— T (Eo-u)
Meandrinidae	K (Barr)	— R
Micrabaciidae	K (Barr)	— R
Microsolenidae	J (Hett)	— K (Sant)
Montlivaltidae	Tr (Anis)	— T (Eo-m)
Mussidae	J (Kimm)	— R
Oculinidae	K (Apti)	— R
Pectiniidae	T (Eo-m)	— R
Pocilloporidae	K (Turo)	— R
Poritidae	K (Ceno)	— R
Procyclolitidae	Tr (Anis)	— J (Bajo)
Rhipidogyridae	J (Kimm)	— K (Sant)
Rhizangiidae	K (Barr)	— R
Siderastreidae	K (Apti)	— R
Stylinidae	Tr (Anis)	— T (Eo-u)
Stylophyllidae	Tr (Anis)	— J (Bajo)
Synastreidae	J (Hett)	— K (Coni)
Thamnasteriidae	Tr (Anis)	— R
Incertae sedis		
<i>Tabulaconus</i>	€ (Boto)	(310)
Tjanshanophyllidae	Θ (u)	— S (l)
incertae sedis	€ (Atda)	(357a)
Or. Actiniaria		
* <i>Mackenzia</i>	€ (mMid)	(59)

\*May be related to unlisted extant families.

## Cl. Incertae sedis

<i>Escumasia</i>	C (Mosc)	(219a)
?Peytoidae	€ (mMid)	(62a)

## PLATYHELMINTHES

?Dickinsoniidae	V (u)	(122,231,369)
?Turbellaria	T (Mi) — R	(62)

RHYNCHOCOELA (Nemertinea) C (Namu) — R (304)

## PRIAPULIDA

[Classification and stratigraphy from Conway Morris (1977a).]

<i>Lecythioscopia</i>	€ (mMid)	
<i>Scolecofura</i>	€ (mMid)	
Ancalagonidae	€ (mMid)	
Fieldiidae	€ (mMid)	
Miskoiidae	€ (mMid)	
Ottoiidae	€ (mMid)	
Priapulidae	C (Mosc) — R	(54,302)
Selkirkidae	€ (mMid)	

NEMATODA (Rhabditida) C (Namu) — R (304,336)

## MOLLUSCA

## Cl. Stenothecoida

Cambridiiidae	€ (Atda) — € (uMid)	(252,275,314,387)
---------------	---------------------	-------------------

## Cl. Monoplacophora

[Classification and stratigraphic ranges from Runnegar and Jell (1976), with emendations as noted.]

## Or. Cyrtinellida

Cyrtolitidae	Θ (l) — D (Give)	
Helcionellidae	€ (ITom) — Θ (l)	(289)
Hypseloconidae	€ (uMid) — Θ (Lvi)	
?Multifariidae	€ (Atda) — Θ (Aren)	(19)
?Pelagiellidae	€ (ITom) — Θ (Trem)	(198,295,388)
Scenellidae	€ (uTom) — D (Sieg)	
Stenothecidae	€ (ITom) — € (uMid)	(198,289,292)
Yochelcionellidae	€ (Boto) — € (IMid)	
?Janospira	Θ (Aren)	(104)

## Or. Tryblidiida

Tryblidiidae	€ (IMid) — R	(19)
--------------	--------------	------

## Cl. Amphineura (= Polyplacophora)

[Classification based on the *Treatise*, Pt. I.]

Or. Palaeoloricata				
Chelodidae	Θ (Trem)	— K (u)		
Gotlandochitonidae	S (m)			
?Mattheviidae	Є (Fran)	— Є (Trep)	(204,293,390)	
Preacanthochitonidae	Є (Trep)	— Θ (Trem)	(132,294)	
Scanochitonidae	K (Camp)			(132)
Septemchitonidae	Θ (Cara)			

Or. Neoloricata

Acanthochitonidae	T (Ol-l)	— R		
Afossochitonidae	T (Mi-m)	— R		
Callistoplacidae	T (Mi-l)	— R		
Callochitonidae	T (Ol)	— R		
Chaetopleuridae	T (Mi)	— R		
Chitonidae	K (Camp)	— R	(317)	
Hanleyidae	Q (Plei)	— R		
Ischnochitonidae	T (Eo-u)	— R		
Lepidopleuridae	Θ (Ashg)	— R		
Llandeilochitonidae	Θ (Llde)			
Mopaliidae	T (Mi-l)	— R		
Schizochitonidae	K (Maes)	— R	(317)	

Cl. Gastropoda

[Classification based on the *Treatise*, Pt. I, and Taylor and Sohl (1962) with numerous sources for stratigraphic ranges, as listed. Only families with marine representatives are listed; 96 extant marine families (including all in the Soleolifera, Phelinoglossoidea, Acochlidoidea, and Parasita) without known fossil records are not listed.]

Or. Bellerophontida

Bellerophontidae	Є (Trep)	— Tr (Indu)	(19,291)	
"Sinuitidae"	Є (Trep)	— Tr (Olen)	(19,291,318)	

Or. Archaeogastropoda

Acmaeidae	Tr (Ladi)	— R	(132)	
Amberleyidae	Tr (Ladi)	— T (Ol)		
Anomphalidae	S (m)	— P (Guad)		
Ataphridae	Tr (Ladi)	— K (Maes)		
Catantostomatidae	D (Eife)	— D (Give)	(189)	
Cirridae	Tr (u)	— J (Bath)		
Clisospiridae	Θ (Aren)	— D (Give)	(136,189)	
Cocculinidae	T (Mi)	— R		
Codonochelidae	S (u)	— J (Bajo)		
Craspedostomatidae	Θ (Ashg)	— J (Bajo)	(229)	
Crossostomatidae	Tr (Ladi)	— J (Bajo)		
Cyclostrematidae	J (Oxfo)	— R		
Deianiridae	K (Camp)	— T (Dani)		

Elasmonematidae	S (u)	— C (Mosc)	(386)
Eotomariidae	€ (Trep)	— J (l)	(132)
Euomphalidae	Θ (Aren)	— K (Maes)	(136,318)
Euomphalopteridae	S (m)	— S (u)	
Fissurellidae	Tr (m)	— R	
Gosseletinidae	Θ (m)	— Tr (Nori)	(176)
Haliotidae	K (Maes)	— R	
Helicotomidae	Θ (Aren)	— J (Bajo)	(136)
Holopeidae	Θ (Aren)	— P (Leon)	
Kittlidiscidae	Tr (Ladi)		
Laubellidae	Tr (Ladi)		
Lepetellidae	T (Mi)	— R	(318)
Lepetidae	T (Mi)	— R	
Lophospiridae	Θ (Llde)	— Tr (m)	(136)
Luciellidae	Θ (Aren)	— Tr (u)	(8,194)
Machuritidae	Θ (Trem)	— Θ (Cara)	(194)
Metoptomatidae	S (m)	— P (Guad)	(8)
Microdomatidae	Θ (m)	— P (Guad)	
Murchisoniidae	Θ (Trem)	— Tr (Nori)	(132)
Neritidae	Tr (Anis)	— R	
Neritopsidae	D (Gedi)	— R	(189)
Nododelphinulidae	Tr (Nori)	— K (Seno)	
Omphalotrochidae	D (Fras)	— P (Guad)	(189)
Onychochilidae	€ (Dres)	— D (Sieg)	(189,291,315)
Oriostomatidae	S (m)	— D (Give)	(189)
Palaeotrochidae	D (Gedi)	— D (Fras)	(189)
Paraturbinidae	Tr (Ladi)	— K (Seno)	(229)
Patellidae	J (Sine)	— R	(176)
Phanerotrematidae	Θ (m)	— D (Sieg)	(189)
Phasianellidae	T (Pale)	— R	
Phenacolepadidae	T (Eo)	— R	
Phymatopleuridae	D (Eife)	— K (Camp)	(136,189)
Plagiothyridae	D (Eife)	— C (l)	(189)
Planitrochidae	Θ (Aren)	— D (Eife)	(189,194)
Platyacridae	Tr (Ladi)	— J (Kimm)	(136, 229)
Platyceratidae	Θ (Aren)	— P (Leon)	
Plethospiridae	Θ (Trem)	— Tr (Nori)	(132,194)
Pleurotomariidae	P (Guad)	— R	(136)
Polytremariidae	C (l)	— P (Leon)	
Porcelliidae	D (Give)	— J (Bajo)	(189)
Portlockiellidae	D (Sieg)	— P (Guad)	(7,189)
Pseudophoridae	S (m)	— P (Guad)	
Raphistomatidae	€ (Trep)	— J (l)	(132)
Raphischismatidae	C (l)		
Schizogoniidae	Tr (Ladi)	— Tr (Carn)	
Scissurellidae	T (Dani)	— R	
Sinuopeidae	€ (Trep)	— P (Leon)	(132)

Stomatellidae	Tr (Nori)	— R	
Symmetrocapulidae	Tr (Ladi)	— J (Kimm)	(71)
Temnotropidae	Tr (Ladi)	— Tr (Carn)	
Trochidae	Tr (Ladi)	— R	
Trochonematidae	Θ (Cara)	— P (Leon)	(136)
Trochotomidae	Tr (Nori)	— J (Tith)	(176)
Tubinidae	D (Sieg)	— Tr (Nori)	(189)
Turbinidae	Tr (Ladi)	— R	
Velainellidae	T (Eo)		
Weeksiidae	Tr	— K (Maes)	(132,318)
Zygitidae	Tr (Ladi)		

## Or. Mesogastropoda

Aciculidae	J (l)	— R	(243)
Aclididae	T (Dani)	— R	(75)
Aporrhaidae	J (Bajo)	— R	(71,320)
Architectonicidae	K (Ceno)	— R	(322)
Assimineidae	K (Maes)	— R	(75)
Atlantidae	K (u)	— R	(213)
Brachytremidae	J (Bajo)	— J (Tith)	(229)
Bursidae	K (Maes)	— R	(347)
Caecidae	T (Pale)	— R	(75)
Calyptaeidae	J (Oxfo)	— R	(213)
Capulidae	Tr	— R	(243)
Carinariidae	T (Eo)	— R	(213)
Cassididae	K (Coni)	— R	(243,323)
Cerithiidae	J (Kimm)	— R	(318,320)
Cerithiopsidae	K (l)	— R	(75)
Coelostylinidae	D	— J (Kimm)	(229)
Colombellinidae	J (u)	— K (Maes)	(138,347)
Cymatiidae	K (Albi)	— R	(75,320,347)
Cypraeidae	J (Tith)	— R	(229)
Diastomidae	K (u)	— R	(229)
Epitoniiidae	J (m)	— R	(75,229)
Eratoidae	T (Dani)	— R	(75)
Eulimidae	K (Ceno)	— R	(322)
Eustomidae	J (Bajo)	— J (Tith)	(229)
Ewekoroidae	T (Pale)		(1)
Ficidae	T (Dani)	— R	(347)
Fossaridae	P	— R	
Glauconiidae	J (Tith)	— K (Maes)	(229)
Hipponicidae	K (Maes)	— R	(229)
Iravadiidae	T (Eo)	— R	(229)
Janthinidae	T (Mi)	— R	(75)
Lacunidae	Tr (m)	— R	(75)
Lamellariidae	T (Pale)	— R	(206,213)
Littorinidae	Tr	— R	(229)

Loxonematidae	Θ (Cara)	— C (Penn)	(132,208,386)
Mathildidae	P (Guad)	— R	(8)
Meekospiridae	C (Vise)	— P (Guad)	(132,208)
Melanopsidae	K (Albi)	— R	(3)
Micromelaniidae	K (l)	— R	(136)
Modulidae	K (u)	— R	(75)
Naticidae	Tr (Carn)	— R	(112,323)
Oleacinidae	T (Ol)	— R	(323)
Omalaxidae	K (Albi)	— R	(8)
Omalogyridae	T (Eo)	— R	
Ovulidae	K (Ceno)	— R	(75,229)
Palaeozygopleuridae	D (Sieg)	— C (l)	(189,208)
Planaxidae	T (Pale)	— R	(1)
Pomatiasidae	K (u)	— R	(75)
Potamididae	P	— R	(213)
Procerithiidae	D (m)	— K (Maes)	(208,318)
Pseudomelaniiidae	S	— T (Mi)	(75,229)
Pseudozygopleuridae	D (Gedi)	— Tr (Ladi)	(8,189,208)
Purpurinidae	Tr (Nori)	— J (Kimm)	(71,176)
Rissoidae	J (l)	— R	(75,229)
Seraphsidae	T (Pale)	— R	
Spirostylidae	Tr	— J	(229)
Stiliferidae	T (Eo)	— R	(75)
Strombidae	K (Haut)	— R	(324)
Struthiolariidae	T (Dani)	— R	(75)
Subulitidae	Θ (Aren)	— P (Guad)	(132,208)
Thiaridae	J (u)	— R	(229)
Tonnidae	K (Maes)	— R	(347)
Trichotropididae	K (Turo)	— R	(322)
Triphoridae	K (Albi)	— R	(75)
Turritellidae	D (Give)	— R	(189,208)
Vanikoridae	K (Sant)	— R	(186,322)
Vermetidae	Tr (m)	— R	(176)
Vitrinellidae	J	— R	(75)
Xenophoridae	J (Tith)	— R	(243)
Zygopleuridae	Tr (Ladi)	— J (Kimm)	(208)
Or. Neogastropoda			
Buccinidae	K (Albi)	— R	(347)
Buccinulidae	T (Ol)	— R	(75)
Cancellariidae	K (Albi)	— R	(347)
Conidae	K (Maes)	— R	(347)
Fasciolariidae	K (Apti)	— R	(3,322,347)
Harpidae	K (Maes)	— R	(347)
Heligmotoniidae	T (Pale)		(1)
Magilidae	K (Camp)	— R	(75,347)
Marginellidae	K (Maes)	— R	(213,243,347)

Melongenidae	K (Albi)	— R	(347)
Mitridae	K (Ceno)	— R	(213)
Muricidae	K (Albi)	— R	(243,323,347)
Nassariidae	K (Turo)	— R	(347)
Neptuneidae	K (u)	— R	(75)
Olividae	K (Camp)	— R	(213,322,347)
Pyrenidae	T (Dani)	— R	(347)
Strepsiduridae	K (u)	— T (Ol)	(75)
Terebridae	K (Maes)	— R	(347)
Turridae	K (Ceno)	— R	(322,347)
Vasidae	K (Albi)	— R	(75,322,347)
Vaupalmeriidae	T (Pale)	— R	(1)
Volutidae	K (Ceno)	— R	(322,347)
Or. Entomotaeniata			
Ceritellidae	J (l)	— K (Ceno)	(206,213)
Itieriidae	J (Oxfo)	— K (Maes)	(229,320)
Nerineidae	J (l)	— K (Maes)	(3,206,213)
Nerinellidae	J (l)	— K (Seno)	(229)
Pyramidellidae	D (Give)	— R	(189)
Streptacidiidae	C (Vise)	— P (Guad)	(208)
Or. Cephalaspidea			
Actæonidae	C (Vise)	— R	(132,206,213)
Acteonellidae	J (l)	— K (Seno)	(229)
Atyidae	T (Pale)	— R	(75)
Bullidae	J (Plie)	— R	(213)
Cylindrobullidae	K (Camp)	— R	(319)
Diaphanidae	J (Bajo)	— R	(229)
Hydatinidae	J (l)	— R	(229)
Philinidae	K (Seno)	— R	(75,229)
Retusidae	J	— R	(75)
Ringiculidae	K (Apti)	— R	(136)
Scaphandridae	K (Camp)	— R	(75,319)
Or. Thecosomata			
Cavolinidae	K (u)	— R	(229)
Spiratellidae	T (Eo-l)	— R	(132)
Or. Basommatophora			
Acroreidae	K (u)	— T (Eo)	
Amphibolidae	T (Plio)	— R	(75)
Siphonariidae	J (Call)	— R	(75,229)
Trimusculidae	T (Ol)	— R	(229)
Or. Sacoglossa			
Arthessidae	T (Eo-l)	— R	(181)
Elysiacea	T (Eo)	— R	(213)
Juliidae	T (Eo-m)	— R	(75)

Or. Anaspidea			
Akeratidae	J (Bath)	— R	(229)
Aplysiidae	T (Mi)	— R	(229)

Or. Gymnosomata			
Clionidae	T (Eo-m)	— R	(213)

Or. Notaspidea			
Umbraculidae	T (Eo)	— R	(75)

Or. Nudibranchia			
*incertae sedis	C (Mosc)		(156)

## Cl. Scaphopoda

[Classification from Palmer (1974).]

Or. Dentalioida			
Dentaliidae	D (u)	— R	(208)
Laevidentaliidae	Θ (Llvi)	— R	(250)

Or. Siphonodentalioida			
Cadulidae	K (l)	— R	(208)
Siphonodentaliidae	P (Sakm)	— R	(385)

## Cl. Rostroconchia

[Classification and stratigraphic ranges based principally on Pojeta and Runnegar (1976).]

Or. Ribeirioida			
Ribeiriidae	Є (uTom)	— Θ (Ashg)	(198,205,291)
Technophoridae	Є (Dres)	— Θ (Ashg)	(249)

Or. Ischyriinoida			
Ischyriiniidae	Θ (Trem)	— Θ (Ashg)	

Or. Conocardioida			
Bransoniidae	Θ (Trem)	— P (Guad)	(314)
Conocardiidae	D (Sieg)	— P (Sakm)	(316,385)
Eopteriidae	Є (Trep)	— Θ (Cara)	(249,314)
Hippocardiidae	Θ (Lde)	— C (Mosc)	(137a)
?Myona	Є (uMid)	— Є (Dres)	(195,210,291)

## Cl. Bivalvia

[Classification and stratigraphic ranges principally from the *Treatise*, Pt. N, with emendations as listed. Exclusively nonmarine families and 5 extant marine families without known fossil representatives are not listed.]

Or. Fordilloida			
Fordillidae	Є (Atda)	— Є (Boto)	(150,247)

\*May be related to unlisted extant families.

?Or. Tuarangiida			
Tuarangiidae	Є (uMid)		(199b)
Or. Solemyoida			
Solemyidae	D (Sieg)	— R	(132)
Or. Nuculoida			
Ctenodontidae	Θ (Llde)	— J (Bath)	(71,136)
Isoarcidae	J (Bajo)	— K (Seno)	
Malletiidae	Θ (Llde)	— R	(136)
Nuculanidae	D (Give)	— R	(316)
Nuculidae	Θ (Trem)	— R	
Praenuculidae	Θ (Trem)	— D (l)	(136,248)
Thoraliidae	Θ (l)		(211)
Or. Mytiloida			
Colpomyidae	Θ (Aren)	— Θ (Ashg)	(248)
Mysidiellidae	Tr (l)	— Tr (Rhae)	
Mytilidae	D (Fame)	— R	(132)
Pinnidae	C (Vise)	— R	(132)
Or. Praecardioida			
Antipleuridae	Θ (m)	— D (Fras)	
Butovicellidae	S (u)		
Praecardiidae	Θ (Aren)	— Tr (Carn)	
Or. Arcoida			
Arcidae	J (Bajo)	— R	(129)
Cucullaeidae	J (Hett)	— R	(129,135)
Cyrtodontidae	Θ (Aren)	— D (m)	(132)
Glycymerididae	K (Apti)	— R	(135)
Limopsidae	Tr (Ladi)	— R	(129a)
Manzanellidae	P (Leon)	— R	
Noetiidae	K (Apti)	— R	
Parallelodontidae	Θ (Llde)	— R	(181)
Philobryidae	T (Eo-u)	— R	
Or. Pterioida			
Ambonychiidae	Θ (Llde)	— D (Fras)	(132,136)
Anomiidae	P (l)	— R	(216)
Aviculopectinidae	D (Fras)	— J (Tith)	(316)
Bakevelliidae	P (Guad)	— T (Eo-l)	
Buchiidae	Tr (Carn)	— K (Maes)	
Cassianellidae	Tr (Ladi)	— Tr (Nori)	(176)
Chondrodontidae	K (Albi)	— K (Camp)	
Dattidae	Tr (Rhae)		
Deltopectinidae	P (Leon)		
Dimyidae	J (Bath)	— R	
Entoliidae	C (Tour)	— K (Seno)	
Gryphaeidae	Tr (Carn)	— R	

Inoceramidae	P (l)	— K (Maes)	(135)
Isognomonidae	P (Sakm)	— R	(136)
Kochiidae	D (Sieg)	— D (u)	
Leiopectinidae	Θ (Ashg)	— D (l)	
Limidae	C (l)	— R	
Lithiotidae	J (Plie)	.	(129)
Lunulacardiidae	Θ (l)	— C (l)	
Malleidae	J (Bajo)	— R	(129)
Monopteriidae	C (Bash)	— C (Step)	
Monotidae	Tr (Carn)	— Tr (Nori)	
Myalinidae	C (Tour)	— J (l)	
Ostreidae	P (Guad)	— R	(136)
Oxytomidae	P (Leon)	— K (u)	
Pectinidae	Tr (Ladi)	— R	(129a)
Pergamidiidae	Tr (Carn)	— J (Sine)	
Plicatulidae	Tr (Ladi)	— R	
Posidoniidae	C (l)	— K (Coni)	
Pseudomonotidae	C (Vise)	— P (u)	
Pteriidae	P (Sakm)	— R	(136,216)
Pterineidae	Θ (Livi)	— P (Guad)	(136)
Pterinopectinidae	S (u)	— P (Leon)	
Pulvinitidae	J (Bath)	— R	(129)
Spondylidae	Tr (m)	— R	(176)
Terquemiidae	P (l)	— J (Kimm)	(216)
Or. Actinodontoida (= Modiomorphoidea)			
Allodesmatidae	Θ (Cara)	— S (u)	
Carydiidae	D (Emsi)	— D (u)	
Cycloconchidae	Θ (Aren)	— Θ (Ashg)	(248)
Lyrodesmatidae	Θ (Aren)	— S (l)	(248)
Modiomorphidae	Θ (Trem)	— P (Guad)	
Or. Unionoida (mostly freshwater)			
Pachycardiidae	Tr (l)	— J (Plie)	(135)
Or. Trigonioida			
Myophoriidae	D (Gedi)	— Tr (Rhae)	
Scaphellinidae	P (Leon)		
Trigoniidae	Tr (Anis)	— R	
Or. Veneroida			
Arcticidae	Tr (Rhae)	— R	
Astartidae	D (l)	— R	
Babinkidae	Θ (Aren)	— Θ (Llde)	(199,210)
Cardiidae	Tr (Nori)	— R	
Cardiliidae	T (Ol)	— R	
Cardiniidae	Θ (Llde)	— R	
Carditidae	D (Gedi)	— R	
Ceratomyopsidae	J (Toar)	— J (Kimm)	(129)

Chamidae	T (Dani)	— R	
Condylocardiidae	T (Eo-m)	— R	
Cooperellidae	T (Mi-m)	— R	
Corbiculidae	J (Toar)	— R	(71)
Crassatellidae	D (l)	— R	
Cultellidae	K (l)	— R	
Cyamiidae	T (Mi)	— R	
Dicerocardiidae	Tr (Carn)	— K (Maes)	
Donacidae	K (u)	— R	
Dreissenidae	T (Eo)	— R	
Erycinidae	T (Pale)	— R	
Euloxidae	T (Mi)		
Fimbriidae	C	— R	
Gaimardiidae	T (Mi)	— R	
Galeommatidae	T (Eo-u)	— R	
Glossidae	K (Maes)	— R	(361a)
Hippopodiidae	Tr (Nori)	— J (Tith)	(129a)
Icanotiidae	K (Haut)	— K (Maes)	
Kelliellidae	T (Eo-l)	— R	
Kelliidae	T (Pale)	— R	
Lahilliidae	K (u)	— T (Mi)	
Leptonidae	T (Pale)	— R	
Lucinidae	S	— R	
Lymnocardiidae	T (Mi-l)	— R	
Mactridae	K (Camp)	— R	(135)
Mactromyidae	D (Eife)	— R	(316)
Mecynodontidae	D (Eife)		
Mesodesmatidae	T (Eo)	— R	
Montacutidae	T (Eo-m)	— R	
Myophoricardiidae	Tr (l)	— Tr (Nori)	
Neoleptonidae	T (Plio)	— R	
Neomiodontidae	J (Hett)	— K (Ceno)	(135)
Permophoridae	C (l)	— K (Camp)	
Petricolidae	T (Eo)	— R	
Pisidiidae	K (l)	— R	(135)
Pollicidae	K (Ceno)	— K (u)	
Psammobiidae	K (u)	— R	
Ptychomyidae	K (l)	— K (u)	
Quenstedtiidae	J (Toar)	— J (Tith)	(129)
Rzehakiidae	T (Mi-m)	— T (Mi-u)	
Scrobiculariidae	T (Eo)	— R	
Semelidae	T (Eo)	— R	
Solecurtidae	T (Eo-l)	— R	
Solenidae	T (Eo-l)	— R	
Sowerbyidae	J (Toar)	— J (Tith)	(129)
Sportellidae	J	— R	
Tancrediidae	Tr (Carn)	— K (Maes)	(135)

Tellinidae	K (l)	— R	
Thyasiridae	Tr (m)	— R	
Trapeziidae	K (Maes)	— R	
Tridacnidae	T (Eo-m)	— R	
Turtoniidae	T (Mi)	— R	
Ungulinidae	K (u)	— R	
Unicardiopsidae	J (m)	— J (Oxfo)	
Veneridae	K (Albi)	— R	
Vesicomyidae	T (Ol-u)	— R	
 Or. Myoida			
Corbulidae	C (Tour)	— R	(132)
Erodonidae	T (Eo-u)	— R	
Gastrochaenidae	J (Bath)	— R	(129)
Hiatellidae	P (l)	— R	
Myidae	T (Pale)	— R	
Pholadidae	J (Bajo)	— R	(129)
Pleurodesmatidae	T (Ol-u)	— T (Mi-l)	
Raetomyidae	T (Eo)		
Spheniopsidae	T (Eo-m)	— R	
Teredinidae	K (Apti)	— R	(135)
 Or. Hippuritoida			
Caprinidae	K (Barr)	— K (Maes)	
Caprotinidae	K (Barr)	— K (Turo)	
Diceratidae	J (Oxfo)	— K (Vala)	
Hippuritidae	K (Turo)	— K (Maes)	
Megalodontidae	S (l)	— K (Apti)	
Monopleuridae	K (Vala)	— K (Maes)	
Radiolitidae	K (Barr)	— K (Maes)	
Requieniidae	J (Tith)	— K (Maes)	
 Or. Pholadomyoida			
Burmesiidae	Tr (Rhae)	— J (Hett)	(129,135)
Ceratomyidae	Tr (Rhae)	— J (Tith)	(129)
Chaenomyidae	C (u)	— T (Eo)	(290)
Clavagellidae	K (Turo)	— R	
Cleidothaeridae	T (Ol)	— R	(290)
Cuspidariidae	Tr (Ladi)	— R	(132)
Edmondiidae	D (Fras)	— P (Dzhu)	(216,316)
Grammysiidae	Θ (Cara)	— P (Guad)	(136)
Laternulidae	Tr (Carn)	— R	(129a)
Lyonsiidae	T (Eo)	— R	
Margaritariidae	T (Mi)		
Megadesmidae	C (u)	— P (Dzhu)	(216)
Myochamidae	T (Mi)	— R	
Myopholadidae	J (Toar)	— K (Albi)	(129)
Orthonotidae	Θ (Llde)	— D (Give)	(248)

Pandoridae	T (Eo)	— R	(290)
Periplomatidae	K (Albi)	— R	(135)
Pholadomyidae	C (I)	— R	
Pleuromyidae	Tr (Nori)	— K (I)	(129a, 290)
Poromyidae	K (u)	— R	
Thraciidae	J (Hett)	— R	(71,129)
Verticordiidae	K (Maes)	— R	(361a)

## Cl. Cephalopoda

[Classification and stratigraphic ranges for the Nautiloidea and Ammonoidea basically follow Teichert (1967) and *The Ammonoidea* (House and Senior, eds. 1981), with augmentation from the *Treatise*, Pts. K and L, and *Osnovy Paleontologii*. Classification of the Coleoidea is from Jeletsky (1965). 25 extant families of coleoids without known fossil representatives are not listed.]

## Or. Ellesmerocerida

Acaroceratidae	€ (Trep)	(45)	
Balkoceratidae	€ (Trep)	(45)	
Baltoceratidae	Θ (Trem)	— Θ (Cara)	(73)
Bassleroceratidae	Θ (Trem)	— Θ (Cara)	(45,73)
Bathmoceratidae	Θ (Aren)	— Θ (Llde)	
Cyrtocerinidae	Θ (Cara)	— Θ (Ashg)	(45,73)
Ellesmeroceratidae	€ (Trep)	— Θ (Llvi)	(45,73)
Huaiheceratidae	€ (Trep)	— Θ (Trem)	(45,73)
Plectronoceratidae	€ (Fran)	— Θ (Llde)	(73)
Protocycloceratidae	Θ (Trem)	— Θ (Llvi)	(45)
Shideleroceratidae	Θ (Ashg)		

## Or. Orthocerida

Brachycycloceratidae	C (Namu)	— C (Step)	
Clinoceratidae	Θ (Llvi)	— Θ (Ashg)	(73,99)
Dawsonoceratidae	S (I)	— S (u)	
Engorthoceratidae	D (Eife)		(353)
Folioceratidae	D (Eife)		(353)
Geisonoceratidae	Θ (Llvi)	— D (Give)	(353)
Lamellorthoceratidae	D (Sieg)	— D (Eife)	(353)
Mysterioceratidae	Θ (Llde)	— Θ (Cara)	(45,73,99)
Offleyoceratidae	S (m)	— D (Gedi)	(353)
Orthoceratidae	Θ (Trem)	— Tr (Nori)	(73)
Paraphragmitidae	S (m)		
Proteoceratidae	Θ (Llvi)	— S (m)	(99)
Pseudorthoceratidae	Θ (Llvi)	— P (I)	(73)
Sactorthoceratidae	Θ (Aren)	— Θ (Llde)	(45,73,99)
Sphooceratidae	S (m)		
Stereoplasmoceratidae	Θ (Aren)	— Θ (Cara)	(45,73)
Troedssonellidae	Θ (Trem)	— Θ (Llde)	(73,229)

Or. Ascocerida			
Ascoceratidae	Θ (Cara)	— S (u)	(73)
Choanoceratidae	S (u)		
Hebetoceratidae	Θ (Llde)	— Θ (Ashg)	(73)
Or. Intejocerida			
Bajkocolceratidae	Θ (Aren)	— Θ (Llvi)	
Intejoceratidae	Θ (Aren)	— Θ (Llvi)	(73)
Majoceratidae	Θ (Trem)	— Θ (Aren)	(73)
Padunoceratidae	Θ (Aren)	— Θ (Llvi)	
Or. Protactinocerida			
Protactinoceratidae	€ (Trep)	— Θ (Trem)	(44,73)
Or. Actinocerida			
Actinoceratidae	Θ (Aren)	— S (l)	
Armenoceratidae	Θ (Aren)	— S (m)	
Carbactinoceratidae	C (Vise)	— C (Step)	
Ellinoceratidae	Θ (Llvi)	— Θ (Llde)	
Georginidae	Θ (Llvi)		(73)
Gonioceratidae	Θ (Llvi)	— Θ (Ashg)	(73)
Huroniidae	Θ (Llvi)	— S (m)	(73)
Meitanoceratidae	Θ (Llvi)	— Θ (Llde)	(73)
Ormoceratidae	Θ (Aren)	— C (Vise)	
Polydesmiidae	Θ (Aren)	— Θ (Llde)	(73)
Wutinoceratidae	Θ (Aren)	— Θ (Llvi)	
Or. Yanhecerida			
Yanheceratidae	€ (Trep)		(44)
Or. Endocerida			
Allotrioceratidae	Θ (Aren)	— Θ (Llde)	(45,73)
Botryceratidae	Θ (Ashg)		(73)
Chihlioceratidae	Θ (Aren)	— Θ (Llvi)	(73,98,229)
Cyrtendoceratidae	Θ (Aren)	— Θ (Cara)	(98)
Emmonsoceratidae	Θ (Aren)	— Θ (Llde)	(45,229)
Endoceratidae	Θ (Trem)	— S (m)	(73)
Humeoceratidae	S (m)		
Manchuroceratidae	Θ (Aren)	— Θ (Cara)	(73)
Nartheoceratidae	Θ (Cara)	— S (l)	(73,229)
Piloceratidae	Θ (Trem)	— Θ (Llvi)	(73)
Proterocameroceratidae	Θ (Trem)	— Θ (Llde)	
Thylacoceratidae	Θ (Aren)	— Θ (Llvi)	(73,98,229)
Yorkoceratidae	Θ (Aren)		(73)
Or. Discosorida			
Brevicoceratidae	S (m)	— D (Fame)	(353)
Cytogomphoceratidae	Θ (Llvi)	— S (l)	(45,73)
Devonocheilidae	D (Emsi)	— D (Fame)	(353)
Discosoridae	S (m)	— D (Give)	

Entimoceratidae	D (Emsi)	— D (Fame)	(353)
Greenlandoceratidae	Θ (Ashg)	— S (l)	(73)
Lowoceratidae	Θ (Cara)	— S (m)	(73)
Mandaloceratidae	S (m)		
Mecynoceratidae	D (Emsi)	— D (Fame)	(353)
Mesoceratidae	S (m)		
Naedyceratidae	S (m)	— D (Fame)	(353)
Phragmoceratidae	S (m)	— D (Gedi)	(353)
Ruedemannoceratidae	Θ (Llvi)	— Θ (Cara)	(45,73)
Taxyceratidae	D (Gedi)	— D (Fame)	(353)
Westonoceratidae	Θ (Llde)	— S (l)	
Or. Oncocerida			
Acleistoceratidae	S (m)	— D (Fame)	(358)
Aktjubocheilidae	D (Fame)		(358)
Archiacoceratidae	D (Sieg)	— D (Give)	(358)
Bolloceratidae	D (Eife)	— D (Fras)	(358)
Cyrtoceratidae	S (u)	— D (Fras)	(358)
Diestoceratidae	Θ (Cara)	— Θ (Ashg)	(73,99)
Graciloceratidae	Θ (Llde)	— Θ (Ashg)	(45,73)
Hemiphragmoceratidae	S (m)	— S (u)	
Jovellaniidae	Θ (Llvi)	— D (Fame)	(73,358)
Karoceratidae	S (l)	— D (Fame)	(358)
Nothoceratidae	S (u)	— D (Fras)	(358)
Oncoceratidae	Θ (Llvi)	— S (u)	
Polyelasmoceratidae	S (m)	— D (Fame)	(358)
Poterioceratidae	D (Gedi)	— C (Namu)	(358)
Ptenoceratidae	D (Sieg)	— D (Eife)	(358)
Trimeroceratidae	S (m)		
Tripleuroceratidae	S (u)	— C (l)	(358)
Tripteroceratidae	Θ (Llvi)	— Θ (Ashg)	(99)
Valcouroceratidae	Θ (Llde)	— Θ (Ashg)	(73,99)
Or. Nautilida			
Aturiidae	T (Pale)	— T (Mi)	
Centroceratidae	D (Eife)	— P (l)	(132)
Clydonautilidae	Tr (Anis)	— Tr (Nori)	
Cymatoceratidae	J (m)	— T (Ol-u)	
Ephippioceratidae	C (Vise)	— P (l)	
Gonionautilidae	Tr (Nori)		
Grypoceratidae	C (l)	— Tr (Nori)	(132)
Hercoglossidae	J (u)	— T (Ol)	
Koninckioceratidae	C (l)	— P (l)	
Liroceratidae	C (Vise)	— Tr (Nori)	
Nautilidae	Tr (Ladi)	— R	
Paracenoceratidae	J (m)	— K (Albi)	
Permoceratidae	P (l)		
Pseudonautilidae	J (Tith)	— K (Vala)	(229)

Rhiphaeoceratidae	P (l)		
Rutoceratidae	D (Sieg)	— C (l)	(132)
Siberionautilidae	Tr (Carn)		
Solenochilidae	C (Tour)	— P (Leon)	
Syringonautilidae	Tr (Anis)	— Tr (Nori)	
Tainoceratidae	C (l)	— Tr (Nori)	(132)
Tetragonoceratidae	D (Eife)		(353)
Trigonoceratidae	C (l)	— P (l)	
 Or. Tarphycerida			
Apsidoceratidae	Θ (Cara)	— Θ (Ashg)	(99)
Barrandeoceratidae	Θ (Livi)	— D (Eife)	(99,353)
Estonioceratidae	Θ (Aren)	— Θ (Livi)	(73)
Lechritrochoceratidae	Θ (Livi)	— S (u)	(73)
Lituitidae	Θ (Aren)	— Θ (Ashg)	
Nephriticeratidae	D (Emsi)	— D (Fras)	(353)
Plectoceratidae	Θ (Livi)	— Θ (Ashg)	(99)
Tarphyceratidae	Θ (Aren)		(45,73)
Trocholitidae	Θ (Aren)	— S (u)	
Uranoceratidae	S (m)		
 Or. incertae sedis			
Neptunoceratidae	C (u)		
 Or. Anarcestida			
Agoniatiidae	D (Eife)	— D (Give)	(139,140,229)
Anarcestidae	D (Emsi)	— D (Fame)	(229)
Auguritidae	D (Emsi)		(139)
Bactritidae	Θ (Aren)	— P (Dzhu)	
Beloceratidae	D (Fras)		(229)
Devonopronoritidae	D (Fras)		(139)
Gephuroceratidae	D (Fras)	— D (Fame)	(140,229)
Maenioceratidae	D (Give)		(139)
Mimagoniatitidae	D (Emsi)	— D (Eife)	(139)
Mimoceratidae	D (Emsi)	— D (Eife)	(139)
Mimosphinctidae	D (Emsi)	— D (Eife)	(139)
Parabactritidae	P (Asse)	— P (Leon)	
Pharciceratidae	D (Fras)		(229)
Phenacoceratidae	D (Fras)	— D (Fame)	(139)
Pinacitidae	D (Eife)	— D (Give)	(139)
Triainoceratidae	D (Fras)		(139)
 Or. Clymeniida			
Biloclymeniidae	D (Fame)		(140)
Carinoclymeniidae	D (Fame)		(140)
Clymeniidae	D (Fame)		
Costaclymeniidae	D (Fame)		(140)
Cymaclymeniidae	D (Fame)		(140)
Cyrtoclymeniidae	D (Fame)		

Glatziellidae	D (Fame)	
Gonioclymeniidae	D (Fame)	
Hexaclymeniidae	D (Fame)	
Miroclymeniidae	D (Fame)	(140)
Parawocklumeridae	D (Fame)	(139)
Rectoclymeniidae	D (Fame)	
Sellaclymeniidae	D (Fame)	(140)
Wocklumeridae	D (Fame)	
 Or. Goniatitida		
Adrianitidae	C (Step)	— P (Guad) (111)
Agathiceratidae	C (Vise)	— P (Guad) (182)
Anthracoceratidae	C (Namu)	— C (Mosc) (111,182)
Berkhoceratidae	C (Vise)	— C (Namu) (182)
Bisatoceratidae	C (Namu)	— C (Step) (182)
Cheiloceratidae	D (Fame)	(111,139)
Christioceratidae	C (Bash)	— C (Mosc) (182)
Cravenoceratidae	C (Vise)	— C (Namu) (182)
Cyclobidae	P (Leon)	— P (Dzhu)
Delepinoceratidae	C (Vise)	— C (Namu) (182)
Dimeroceratidae	D (Fame)	(139)
Dimorphoceratidae	C (Vise)	— C (Namu) (111,182)
Eogonioloboceratidae	C (Vise)	— C (Namu) (182)
Gastrioceratidae	C (Namu)	— P (Asse) (111)
Girtyoceratidae	C (Vise)	— C (Namu) (182)
Glaphyritidae	C (Namu)	— P (Asse) (182)
Goniatitidae	C (Vise)	(182)
Gonioloboceratidae	C (Mosc)	— P (Asse) (182)
Homoceratidae	C (Namu)	— C (Bash) (111,182)
Hyattoceratidae	P (Leon)	— P (Guad) (182)
Intoceratidae	C (Tour)	— C (Vise) (182)
Karagandoceratidae	C (Tour)	(182)
Marathonitidae	C (Step)	— P (Guad) (182)
Maximitidae	C (Mosc)	— C (Step) (111,182)
Metalegoceratidae	P (Asse)	— P (Guad) (111,182)
Munsteroceratidae	C (Tour)	— C (Namu) (182)
Neodimorphoceratidae	C (Namu)	— C (Step) (182)
Neoglyphioceratidae	C (Vise)	— C (Namu) (111)
Neoicoceratidae	C (Mosc)	— P (Asse) (182)
Nomismoceratidae	C (Vise)	— C (Namu) (111)
Orulganitidae	C (Bash)	(182)
Paragastrioceratidae	P (Asse)	— P (Dzhu) (111)
Pericyclidae	C (Tour)	— C (Vise) (182)
Perrinitidae	P (Asse)	— P (Leon) (111,182,208)
Popanoceratidae	P (Asse)	— P (Guad) (111)
Praeglyphioceratidae	D (Fame)	(111)
Prionoceratidae	D (Fame)	— C (Namu) (139)

Prolobitidae	D (Give)	— D (Fame)	(139)
Pseudohaloritidae	C (Step)	— P (Dzhu)	(182)
Pseudoparalegoceratidae	C (Bash)	— C (Mosc)	(182)
Ramositidae	C (Namu)		(182)
Reticuloceratidae	C (Namu)	— C (Mosc)	(111,182)
Rhymnoceratidae	C (Vise)	— C (Namu)	(182)
Schistoceratidae	C (Bash)	— C (Step)	(111)
Shumarditidae	C (Step)		(182)
Sinotitidae	D (Fame)		(139)
Somoholitidae	C (Mosc)	— P (Leon)	(111,182)
Sporadoceratidae	D (Fame)		(111)
Thalassoceratidae	C (Mosc)	— P (Guad)	(182)
Tornoceratidae	D (Give)	— D (Fame)	(111)
Vidrioceratidae	C (Step)	— P (Dzhu)	(182)
Welleritidae	C (Bash)	— C (Mosc)	(111)
Wiedeyoceratidae	C (Bash)	— C (Step)	(182)
 Or. Ceratitida			
Acrochordiceratidae	Tr (Olen)	— Tr (Anis)	(355)
Aploceratidae	Tr (Anis)		(355)
Araxoceratidae	P (Guad)	— P (Dzhu)	(208)
Arcestidae	Tr (Anis)	— Tr (Rhae)	
Arpaditidae	Tr (Ladi)	— Tr (Nori)	
Aspenitidae	Tr (Olen)		(355)
Badiotitidae	Tr (Carn)		(355)
Balatonitidae	Tr (Anis)		(355)
Beneckeiiidae	Tr (Olen)	— Tr (Anis)	(355)
Carnitidae	Tr (Ladi)	— Tr (Carn)	
Ceratitidae	Tr (Anis)	— Tr (Ladi)	
Choristoceratidae	Tr (Nori)		(355)
Cladiscitidae	Tr (Anis)	— Tr (Nori)	
Clionitidae	Tr (Ladi)	— Tr (Nori)	(355)
Clydonitidae	Tr (Carn)	— Tr (Nori)	
Cochloceratidae	Tr (Nori)		
Columbitidae	Tr (Olen)		(355)
Cycloceltitidae	Tr (Nori)		(355)
Cyrtopleuritidae	Tr (Nori)		(355)
Danubitidae	Tr (Anis)	— Tr (Ladi)	(355)
Didymitidae	Tr (Nori)		
Dinaritidae	Tr (Olen)	— Tr (Anis)	(355)
Distichitidae	Tr (Nori)		(355)
Dzhulfitidae	P (Dzhu)		(111)
Episculitidae	Tr (Nori)		(355)
Gymnitidae	Tr (Anis)	— Tr (Nori)	(355)
Haloritidae	Tr (Carn)	— Tr (Nori)	
Hedenstroemiidae	Tr (Olen)		
Heraclitidae	Tr (Nori)		
Hungaritidae	Tr (Ladi)	— Tr (Carn)	(355)

Isculitidae	Tr (Anis)	(355)
Japonitidae	Tr (Olen)	— Tr (Anis) (355)
Joannitidae	Tr (Ladi)	— Tr (Carn) (355)
Keyserlingitidae	Tr (Olen)	(355)
Lecanitidae	Tr (Carn)	(355)
Lobitidae	Tr (Ladi)	— Tr (Carn)
Longobarditidae	Tr (Anis)	— Tr (Ladi) (355)
Meekoceratidae	Tr (Indu)	— Tr (Olen) (355)
Megaphyllitidae	Tr (Anis)	— Tr (Rhae)
Melagathiceratidae	Tr (Olen)	(355)
Metasibiritidae	Tr (Nori)	
Nannitidae	Tr (Carn)	(355)
Nathorstitidae	Tr (Ladi)	
Noridiscitidae	Tr (Nori)	(355)
Noritidae	Tr (Olen)	— Tr (Carn) (355)
Olenikitidae	Tr (Olen)	(355)
Ophiceratidae	Tr (Indu)	— Tr (Olen) (355)
Otoceratidae	P (Dzhu)	— Tr (Indu)
Paraceltitidae	P (Leon)	— P (Guad) (208)
Paranannitidae	Tr (Olen)	
Parapopanoceratidae	Tr (Anis)	(355)
Pinacoceratidae	Tr (Ladi)	— Tr (Nori) (355)
Prionitidae	Tr (Olen)	
Procarnitidae	Tr (Olen)	(355)
Proteusitidae	Tr (Anis)	(355)
Ptychitidae	Tr (Anis)	— Tr (Ladi)
Rhabdoceratidae	Tr (Nori)	(355)
Rimkinitidae	Tr (Ladi)	(355)
Sageceratidae	Tr (Olen)	— Tr (Carn) (355)
Sagenitidae	Tr (Ladi)	— Tr (Nori) (355)
Sandlingitidae	Tr (Carn)	(355)
Sibiritidae	Tr (Olen)	(355)
Sphingitidae	Tr (Ladi)	— Tr (Carn) (355)
Stephanitidae	Tr (Olen)	
Sturiidae	Tr (Olen)	— Tr (Ladi) (355)
Thanamitidae	Tr (Ladi)	(355)
Thetiditidae	Tr (Nori)	(355)
Thisbitidae	Tr (Carn)	— Tr (Nori)
Tibetitidae	Tr (Nori)	(355)
Trachyceratidae	Tr (Ladi)	— Tr (Carn) (355)
Tropiceltitidae	Tr (Carn)	— Tr (Nori) (355)
Tropitidae	Tr (Carn)	— Tr (Nori)
Ussuriidae	Tr (Olen)	
Xenodiscidae	P (Guad)	— Tr (Olen) (208,355)
Or. Prolecanita		
Daraelitidae	C (Vise)	— P (Guad)
Episageceratidae	P (Guad)	— Tr (Indu) (208,355)

Medlicottiidae	C (Mosc)	— P (Dzhu)	(111,182)
Prodromitidae	C (Tour)		
Prolecanitidae	C (Tour)	— C (Namu)	(111,182)
Pronoritidae	C (Vise)	— P (Guad)	(111)
Shikhanitidae	P (Asse)		(208)
Sundaitidae	P (Sakm)	— P (Guad)	(182)
 Or. Phyllocerida			
Acanthoceratidae	K (Albi)	— K (Coni)	(382)
Astiericeratidae	K (Albi)		(382)
Brancoiceratidae	K (Albi)	— K (Ceno)	(382)
Coilopoceratidae	K (Coni)	— K (Maes)	(382)
Collignoniceratidae	K (Turo)	— K (Maes)	(382)
Deshayesitidae	K (Apti)	— K (Albi)	(382)
Desmoceratidae	K (Haut)	— K (Camp)	(382)
Discophyllitidae	Tr (Carn)	— Tr (Nori)	
Douvilleiceratidae	K (Barr)	— K (Albi)	
Engonoceratidae	K (Albi)	— K (Turo)	
Eodesmoceratidae	K (Vala)	— K (Barr)	(382)
Flickiidae	K (Albi)	— K (Turo)	(382)
Forbesiceratidae	K (Albi)	— K (Ceno)	(382)
Hoplitidae	K (Albi)	— K (Ceno)	
Juraphyllitidae	J (Hett)	— J (Plie)	
Kossmaticeratidae	K (Albi)	— K (Maes)	
Leymeriellidae	K (Albi)		(382)
Lyelliceratidae	K (Albi)	— K (Ceno)	(382)
Muniericeratidae	K (Turo)	— K (Maes)	(382)
Pachydiscidae	K (Albi)	— K (Maes)	
Parahoplitidae	K (Apti)	— K (Albi)	(382)
Phylloceratidae	J (Hett)	— K (Maes)	
Placenticeratidae	K (Albi)	— K (Camp)	(382)
Pulchelliidae	K (Haut)	— K (Apti)	
Schloenbachiiidae	K (Albi)	— K (Turo)	
Silesitidae	K (Barr)	— K (Albi)	
Sphenodiscidae	K (Sant)	— K (Maes)	(382)
Tissotiidae	K (Coni)	— K (Sant)	(165,382)
Trochleiceratidae	K (Apti)	— K (Albi)	(382)
Ussuritidae	Tr (Olen)	— Tr (Rhae)	
Vascoceratidae	K (Ceno)	— K (Coni)	(382)
 Or. Lytokerida			
Ancyloceratidae	K (Vala)	— K (Albi)	(382)
Anisoceratidae	K (Albi)	— K (Turo)	
Baculitidae	K (Albi)	— K (Maes)	
Bochianitidae	J (Tith)	— K (Apti)	
Hamitidae	K (Albi)	— K (Turo)	(382)
Hemihoplitidae	K (Barr)		(382)
Heteroceratidae	K (Barr)	— K (Apti)	

Labeceratidae	K (Albi)	— K (Ceno)	(382)
Lytoceratidae	J (Sine)	— K (Ceno)	
Macroscaphitidae	K (Haut)	— K (Albi)	(382)
Protetragonitidae	J (Tith)	— K (Albi)	
Ptychoceratidae	K (Haut)	— K (Apti)	(382)
Scaphitidae	K (Albi)	— K (Maes)	
Tetragonitidae	K (Barr)	— K (Maes)	
Turrilitidae	K (Albi)	— K (Maes)	(382)

## Or. Ammonitida

Amaltheidae	J (Plie)		
Arietitidae	J (Hett)	— J (Sine)	(81)
Aspidoceratidae	J (Call)	— J (Tith)	
Ataxioceratidae	J (Kimm)	— J (Tith)	(81)
Aulacostephanidae	J (Oxfo)	— J (Tith)	(81)
Binneyitidae	K (Haut)	— K (Coni)	(81)
Cardioceratidae	J (Bajo)	— J (Kimm)	(81)
Coeloceratidae	J (Sine)	— J (Plie)	(81)
Craspeditidae	K (Berr)	— K (Barr)	(81)
Cymbitidae	J (Hett)	— J (Plie)	(81)
Dactylioceratidae	J (Plie)	— J (Toar)	
Dorsoplanitidae	J (Kimm)	— J (Tith)	(81)
Echioceratidae	J (Sine)		(81)
Graphoceratidae	J (Bajo)		
Haploceratidae	J (Kimm)	— K (Barr)	(81)
Hildoceratidae	J (Plie)	— J (Bajo)	(81)
Himalayitidae	J (Tith)	— K (Berr)	(81)
Holcodiscidae	K (Haut)	— K (Barr)	(81)
Kosmoceratidae	J (Bath)	— J (Call)	(81)
Liparoceratidae	J (Plie)		
Lissoceratidae	J (Bajo)	— J (Oxfo)	(81)
Macrocephalitidae	J (Bath)	— J (Call)	
Morphoceratidae	J (Bajo)	— J (Bath)	
Neocomitidae	K (Berr)	— K (Haut)	(81)
Olcostephanidae	J (Tith)	— K (Haut)	(81)
Oosterellidae	K (Vala)	— K (Haut)	
Oppeliidae	J (Bajo)	— K (Haut)	(81)
Otoitidae	J (Bajo)		
Oxynoticeratidae	J (Sine)	— J (Plie)	
Pachyceratidae	J (Call)	— J (Oxfo)	
Perisphinctidae	J (Bajo)	— J (Tith)	
Phricodoceratidae	J (Sine)	— J (Plie)	(81)
Phymatoceratidae	J (Toar)	— J (Bajo)	(81)
Polymorphitidae	J (Plie)		
Psiloceratidae	J (Hett)		(81)
Reineckeidae	J (Call)		(81)
Schlottheimiidae	J (Hett)	— J (Sine)	(81)

Simoceratidae	J (Tith)	(81)
Sonniniidae	J (Bajo)	(81)
Sphaeroceratidae	J (Bajo)	— J (Call) (81)
Spiroceratidae	J (Bajo)	— J (Call) (81)
Stephanoceratidae	J (Bajo)	— J (Bath)
Strigoceratidae	J (Bajo)	— J (Call) (81)
Tulitidae	J (Bath)	— J (Call)
Virgatitidae	J (Kimm)	— J (Tith) (81)
Xipheroceratidae	J (Sine)	— J (Plie) (81)
 Or. Belemnitida		
Bayanoteuthidae	T (Eo-u)	(334)
Belemnitellidae	K (Ceno)	— K (Maes) (334)
Belemnitidae	J (Hett)	— J (Call) (333)
Belemnoteuthididae	J (Call)	— J (Kimm) (333)
Belemnopseidae	J (Toar)	— K (Barr) (333,334)
Chondroteuthidae	J (Toar)	(333)
Cylindroteuthididae	J (Bajo)	— K (Apti) (333,334)
Dicoelitidae	J (Bajo)	— J (Tith) (333)
Dimitobelidae	K (Barr)	— K (Maes) (334)
Diplobelidae	J (Tith)	— K (Ceno) (333,334)
Duvaliidae	J (Toar)	— K (Apti) (333,334)
Hastitidae	J (Plie)	— J (Tith) (333)
Oxyteuthidae	K (Vala)	— K (Apti) (334)
 Or. Octopida		
Argonautidae	T (Mi)	— R (132)
*Palaeoctopodidae	K (Coni)	(132,243)
 Or. Sepiida		
Belemnoseidae	T (Dani)	— T (Ol-l) (243)
Belopteridae	T (Eo)	(243)
Belocephiliidae	T (Eo-m)	(243)
Groenlandibelidae	K (Ceno)	— K (Maes) (243)
Sepiidae	T (Eo-u)	— R (243)
Spirulidae	T (Mi)	— R (243)
Spirulirostridae	T (Eo)	— T (Mi-m) (243)
Vasseuriidae	T (Eo)	(243)
 Or. Teuthida		
Geopeltididae	J (Toar)	— J (Tith) (243)
Gonatidae	T (Plio)	— R (50)
Kelaenidae	J (Kimm)	— J (Tith) (243)
Leptoteuthididae	K (Ceno)	— K (Seno) (243)
Lioteuthididae	J (Toar)	(243)
Loliginidae	T (Eo-m)	— R (50)
Loligosepiidae	Tr (u)	— J (Tith) (243)

\*May be related to unlisted extant families.

Necroteuthididae	T (Ol-l)		(243)
Ommastrephidae	T (Plio)	— R	(50)
Onychoteuthidae	T (Plio)	— R	(50)
Palaeoliginidae	J (Kimm)	— K (Sant)	(243)
Pleisioteuthidae	J (Toar)	— K (Maes)	(243)
Teudopseidae	J (Toar)	.	(243)
Trachyteuthididae	J (Kimm)	— K (Maes)	(243)
Or. Phragmoteuthida			
Phragmoteuthidae	P (Guad)	— J (Sine)	(132,181)
Or. Aulacocerida			
Aulacoceratidae	D (Fras)	— Tr (Nori)	
Chitinoteuthidae	J (Hett)	— J (Plie)	
Xiphoteuthididae	Tr (Anis)	— J (Oxfo)	

## ?MOLLUSCA

## Cl. Cricoconarida (= Tentaculitoidea)

[Sources of classification and stratigraphic ranges as indicated.]

## Or. Tentaculitida

Gotlandellitidae	S (l)	— S (m)	(184)
Rossiitiidae	S (u)	— D (Fras)	(184)
Tentaculitidae	Θ (Trem)	— D (Fras)	(132,184)
Volynitidae	S (m)	— D (Gedi)	(184)

## Or. Homoctenida

Homoctenidae	D (Sieg)	— D (Fras)	(30)
--------------	----------	------------	------

## Or. Dacryococonarida

Nowakiidae	S (u)	— D (Fras)	(132)
Striatostyliolinidae	D (Sieg)	— D (Give)	(30)
Styliolinidae	S (m)	— D (Fame)	(132)

## Cl. Hyolitha (= Calyptoptomatida)

[Classification primarily follows Rozanov et al. (1969) with additional families and stratigraphic ranges as indicated. *Wyattia* (Taylor, 1966) is included in the Globorilidae.]

## Or. Orthothecida

Allathecidiae	Є (ITom)	— Θ	(70,289)
Ceratothecidae	S (u)	— D (Gedi)	(132)
Circothecidae	V (N-Da)	— Θ (Cara)	(198,289,358)
Isitithecidae	Є (uTom)	— Є (Boto)	(340)
Orthothecidae	Є (ITom)	— D (m)	(132,198,289,358)
Pauxillitidae	Θ (Trem)	— Θ (Llde)	(132)
Tetrathecidae	Є (ITom)	— Θ (l)	(287,340,358)

## Or. Hyolithida

Amydaicoridiae	Є (Atda)	— Є (mMid)	(358)
----------------	----------	------------	-------

Golicornidae	€ (Atda)	— € (Boto)	(358)
Hyolithidae	€ (uTom)	— P (Guad)	(132,195,358)
Pterigothecidae	Θ (l)	— D	(358)
Sulcavitiidae	€ (Atda)	— S (l)	(205,208,358)
Trapevitidae	€ (uTom)	— € (Atda)	(69,289,358)
Or. Globorilida			
Globorilidae	V (N-Da)	— € (IMid)	(163,172,253,348)
Or. Camerothecida			
Camerothecidae	€ (Boto)		(197,208,220)
Diplothecidae	€ (Boto)		(197,208,220)
SIPUNCULIDA (1 extant family)	D (m)	— R	(54)
ECHIURIDA (5 extant families)			
* <i>Coprinoscolex</i>	C (Mosc)		(158)
? <i>Protechiurus</i>	V (u)		(120)

## ANNELIDA

## Cl. Polychaeta

[Classification from Clark (1969) with significant additions of extinct families from Kielan-Jaworska (1966); stratigraphic ranges in part from the *Treatise*, Pt. W. Approximately 44 extant families without known fossil representatives are not listed.]

Or. Amphinomorpha			
Amphinomidae	C (Mosc)	— R	(302,354)
Or. Eunicemorpha			
Archaeopriionidae	Θ (Cara)		(203)
*Atraktopriionidae	Θ (m)	— P (Guad)	(166,341)
Dorvilleidae	J (Call)	— R	(342)
Kallopriionidae	Θ (m)	— P (Guad)	(166)
Kielanopriionidae	D (Fras)	— P (Gaud)	(341,343)
Lumbrinereidae	C (Namu)	— R	(304)
Mochtyellidae	Θ (m)	— P (Gaud)	(166,341)
Paulinitidae	Θ (Trem)	— C	(167)
Polychaeturidae	Θ (m)	— D (Fras)	(167,349)
Ramphopriionidae	Θ (m)		(166)
Rhytipriionidae	Θ (m)	— S	(166)
*Skalenopriionidae	Θ (Ashg)	— P	(166)
Symmetropriionidae	Θ (Ashg)	— S	(166)
Tetrapriionidae	Θ (m)	— S (m)	(166)
Xaniopriionidae	Θ (m)	— D (Fras)	(166,341)

\*May be related to unlisted extant families.

Or. Phyllocoemorpha			
Aphroditidae	D (u)	— R	(303,354)
Fossundecimidae	C (Mosc)	—	(303,354)
Hesionidae	C (Mosc)	— R	(303,354)
Glyceridae	Θ	— R	(208)
Goniadidae	C (Vise)	— R	(303)
Nephthyidae	C (Vise)	— R	(303,354)
Nereidae	Θ (m)	— R	(208)
Palmyridae	C (Mosc)	— R	(354)
Phyllodocidae	S	— R	(208)
Sigalionidae	Θ (m)	— R	(208)
Sprigginidae	V (u)	—	(117,119)
Or. Spiromorpha			
Cirratulidae	T (Mi)	— R	(208)
?Keiloritidae	Θ	— S	(208)
Sabellariidae	C	— R	(208)
Spionidae	€ (Boto)	— R	(119)
Or. Drilomorpha			
Arenicolidae	Tr (Rhae)	— R	(208)
Opheliidae	€ (Boto)	— R	(119)
Or. Terebellomorpha			
Pectinariidae	C (Namu)	— R	(89)
Terebellidae	€ (mMid)	— R	(119)
Or. Flabelligerimorpha			
Flabelligeridae	C (Mosc)	— R	(354)
Or. Serpulimorpha			
Sabellidae	C (1)	— R	(229)
Serpulidae	Θ	— R	(208)
Incertae sedis			
<i>Eopolychaetus</i>	Θ (Ashg)	—	(181,208)
<i>Pontobdellepsis</i>	Θ (Ashg)	—	(181,208)
<i>Ramesses</i>	C (Namu)	—	(304)
<i>Soris</i>	C (Namu)	—	(304)
Burgessochaetidae	€ (mMid)	—	(60)
Canadiidae	€ (mMid)	— € (uMid)	(60)
Insilicoryphidae	€ (mMid)	—	(60)
Palaeoscolecidae	€ (Boto)	— S (u)	(60,119,208)
Peronochaetidae	€ (mMid)	—	(60)
Stephanoscolecidae	€ (mMid)	—	(60)
Cl. Myzostomaria (6 extant families)			
	D (m)	— R	(62)
Cl. Clitellata			

[Classification from Clark (1969). The great majority of families are nonmarine.]

Subcl. Oligochaeta (3 extant marine families)  
 \*incertae sedis C (Bash) (178)

Subcl. Hirudinea (2 extant marine families)  
 \*incertae sedis J (Bajo) (177)

?ANNELIDA [Groups included largely follow Glaessner (1976).]  
 Or. Cribriocyathida

[Classification and stratigraphic ranges mostly from the *Treatise*, Pt. E (revised). *Cloudina* is included in the Vologdinophyllidae.]

Acherocyathidae	€ (Atda)	(146)
Akademiophyllidae	€ (Atda)	(146)
Capillicyathidae	€ (Boto)	
Conoidocyathidae	€ (Boto)	— € (IMid)
Cribriocyathidae	€ (Boto)	— € (IMid)
Leibaellidae	€ (Atda)	
Pyxidocyathidae	€ (Boto)	— € (IMid)
Vologdinophyllidae	V (u)	— € (Atda) (115,146)

Or. Coleolida  
 Coleolidae V (N-Da) — C (Tour) (132,329)

?Or. Volborthellida

[Considered a distinct phylum, "Agnata," by Yochelson (1977) but a part of Spiromorpha (Polychaeta) by Glaessner (1976). Classification is from the *Treatise*, Pt. W.]

Salterellidae	€ (Boto)	(388,391)
Volborthellidae	€ (Atda)	— € (m) (131,132,163,388)
Vologdinellidae	€ (uMid)	(351)

?Or. Machaeridia  
 Lepidocoleidae Θ (m) — D (l) (251)  
 Plumulitidae D (l) (149)  
 Turrilepodidae Θ (m) — (Namu) (251)

Or. Mitrosagophora

[Classification from Bengtson (1970, 1977a).]

Lapworthellidae	€ (ITom)	— € (m) (132,198,289,340)
Tannuolinidae	€ (Boto)	(395)
Tommotiidae	€ (ITom)	— € (Atda) (69,198,289)

Or. Thambetolepida (=Coeloscleritophora)  
 Chancelloriidae € (ITom) — € (u) (18,69,74)  
 Siphogonuchitidae € (uTom) (253)  
 Wiwaxiidae (=Sachitidae) € (ITom) — € (mMid) (15,18,61,150,202)

\*May be related to unlisted extant families.

## Incertae sedis

Anabaritidae	V (N-Da)	— € (uTom)	(69,329,358)
Ancientidae	Θ (m)	— Θ (Ashg)	(284,299a)
Angustiochreidae	V (N-Da)	— € (ITom)	(358)
Cornulitidae	Θ (Cara)	— C (Tour)	(132)
Platysolenitidae	€ (ITom)	— € (Atda)	(163,289)

## INCERTAE SEDIS

[Miscellaneous taxa of familial or probable familial status but of unknown taxonomic affinities.]

Amiskwiidae	€ (mMid)	(55)	
Konyriidae	Θ (Llv)	— Θ (Llde)	(23a)
Opabinidae	€ (mMid)	(61,373)	
Banffia	€ (mMid)	(61)	
Cambrotubulus	V (N-Da)	— (uTom)	(69,329,358)
Deuteronectanebos	C (Namu)	(304)	
Dinomischus	€ (mMid)	(56)	
Etacystis	C (Mosc)	(219)	
Hallucigenia	€ (mMid)	(57)	
Lenargyrion (=Utaphospha)	€ (Atda)	— Θ (Aren)	(16,260)
Nectocaris	€ (mMid)	(61)	
Oesia	€ (mMid)	(61)	
Portalia	€ (mMid)	(61)	
Redoubtia	€ (mMid)	(61)	
Rhomboconiculum	€ (Atda)	— € (Boto)	(32,33)
Sunnaginia	€ (ITom)	— € (Atda)	(289)
Tullimonstrum	C (Mosc)	(266)	
Tummulduria	€ (ITom)	(358)	
Worthenella	€ (mMid)	(61)	

## ARTHROPODA

## Chelicerata

## Cl. Merostomata

[Classification and stratigraphic ranges from the *Treatise*, Pt. P, with emendations as indicated. Only marine taxa are listed.]

## ?Or. Aglaspida

Aglaspididae	€ (Fran)	— Θ (m)	(20,254)
?Beckwithiidae	€ (uMid)	— € (Dres)	(132,254,337)

## Or. Chasmataspida

Chasmataspidae	Θ (Llv)	(132)
?Diploaspidae	D (Emsi)	(132)
?Heteroaspidae	D (Emsi)	(132)

Or. Xiphosurida			
<i>Drabovaspis</i>	Θ (m)	—	(22)
Archaeolimulidae	Θ (m)	—	(22)
Belinuridae	D (Gedi)	— C (Step)	(97)
Bunodidae	S (u)	—	(85)
Elleriidae	D (u)	—	(97)
Eolimulidae	€ (Atda)	—	(20,196)
Kiaeriidae	D (Gedi)	—	(97)
Limulidae	Tr (l)	— R	(97)
Mesolimulidae	Tr (l)	— K (Ceno)	(97)
Paleolimulidae	D (Fame)	— P (Leon)	(97)
Pseudoniscidae	S (m)	— D (Sieg)	(85,132)
Weinberginidae	D (Sieg)	—	(132)

Incertae sedis			
?Gdowia	€ (uTom)	—	(329)

## Or. Eurypterida

[Classification from Størmer (1974); marine taxa only.]

Carcinosomatidae	Θ (Llvi)	— D (Emsi)	(279)
Dolichopteridae	Θ (Llvi)	— D (Sieg)	
Drepanopteridae	Θ (Caro)	— D (Fame)	(279,338)
Eurypteridae	Θ (Llde)	— D (Emsi)	(132,208)
Hibbertopteridae	D (Eife)	— C (Step)	(279,338)
Hughmilleriidae	Θ (Llde)	— D (Fame)	(132,208)
Kokomopteridae	S (u)	—	(338)
Megalograptidae	Θ (Caro)	— S (m)	(132,144)
Mixopteridae	S (m)	— S (u)	(338)
Pageidae	D (Gedi)	— D (Sieg)	(279,338)
Parastylonuridae	Θ (Ashg)	— D (Fame)	
Pterygotidae	Θ (Llvi)	— D (Fame)	(132,208,279)
Rhenopteridae	D (Gedi)	— D (Fras)	(279)
Slimoniidae	S (l)	— D (Gedi)	(338)
Stylonuridae	Θ (Llde)	— D (Fame)	

## Cl. Arachnida

[Classification from the *Treatise*, Pt. P; marine taxa only.]

Or. Scorpionida			
Dolichophonidae	S (m)	— S (u)	(132,208,244)
Palaeophonidae	S (u)	—	(208,244)
Palaeoscorpiidae	D (Sieg)	—	(132,208,244)
?Waeringscorpii	D (Emsi)	—	(208,244)

## Cl. Pycnogonida

Or. Palaeopantopoda			
Palaeopantopodidae	D (Sieg)	—	(132,208)

Or. Palaeoisopoda			
Palaeoisopodidae	D (Sieg)		(23,132,208)
Or. Pantopoda			
* <i>Palaeotheca</i>	D (Sieg)		(23)
Trilobitomorpha			
Cl. Trilobita			
[Classification primarily from the <i>Treatise</i> , Pt. 0, except for the Agnostida which is from Öpik (1963). Stratigraphic ranges are from numerous sources, as indicated.]			
?Or. Agnostida (=Miomera)			
Agnostidae	Є (Boto)	— Θ (Ashg)	(132,208)
Clavagnostidae	Є (uMid)	— Є (Dres)	(132,148,193)
Condylopygidae	Є (Boto)	— Є (uMid)	(110,132,208)
Diplagnostidae	Є (mMid)	— Θ (Trem)	(132,148,193)
Discagnostidae	Є (Dres)		(226)
Eodiscidae	Є (Atda)	— Є (uMid)	(110,132,193)
Pagetiidae	Є (Atda)	— Є (mMid)	(132,148,287)
Phalacromidae	Є (uMid)	— Є (Dres)	(171,172)
Sphaeragnostidae	Θ (Ashg)		(226)
Trinodidae	Є (Dres)	— Θ (Ashg)	(226)
Or. Redlichiida			
Abadiellidae	Є (Atda)	— Є (lMid)	(132,141,895)
Bathynotidae	Є (Boto)	— Є (lMid)	(148,193,208)
Chengkouiidae	Є (Boto)		(393)
Daguinaspididae	Є (Atda)		(132,141)
Despujolsiidae	Є (Atda)		(141)
Dolerolenidae	Є (Atda)	— Є (Boto)	(141,195,261)
?Ellipsocephalidae	Є (Atda)	— Є (mMid)	(110,141,261)
Emuellidae	Є (lMid)		(173,245)
Gigantopygidae	Є (Boto)		(141,173)
Hicksiidae	Є (Boto)		(141)
Kueichowiidae	Є (Boto)		(173,393)
Longduiidae	Є (Boto)		(393)
Mayiellidae	Є (Boto)		(42,173)
Neoredlichiidae	Є (Atda)	— Є (Boto)	(141,173,262)
Olenellidae	Є (Atda)	— Є (mMid)	(110,132,262)
Paradoxididae	Є (Atda)	— Є (uMid)	(110,173,262)
Protolenidae	Є (lTom)	— Є (mMid)	(92,141,173,262)
Redlichiidae	Є (Atda)	— Є (mMid)	(132,141,173,262)
Saukiandidae	Є (Boto)		(141,262)
Yinitidae	Є (Atda)	— Є (Boto)	(195,393)
Yunnanocephalidae	Є (Atda)		(170,195)

\*May be related to unlisted extant families.

Or. Corynexochida			
Balangiidae	€ (Boto)	— € (IMid)	(173,195)
Corynexochidae	€ (Boto)	— € (Fran)	(227,261,262,314)
Dinesidae	€ (Boto)	— € (uMid)	(148,263)
Dolichometopidae	€ (Boto)	— € (Dres)	(193,261,263)
Dorypygidae	€ (Atda)	— € (Trep)	(110,261,379)
Ogygopsidae	€ (Boto)	— € (mMid)	(110,193)
Oryctocephalidae	€ (Atda)	— € (uMid)	(132,173,195,393)
Zacanthoididae	€ (Boto)	— € (uMid)	(110,193,261)
Or. Ptychopariida			
Agraulidae	€ (IMid)	— € (Dres)	(110,148,296)
Alokistocaridae	€ (Boto)	— € (Fran)	(110,193,381)
Alsataspidae	Θ (Trem)	— Θ (Llvi)	(132,379)
Andrarinidae	€ (mMid)	— € (Fran)	(110,148,193)
Annamitiidae	€ (uMid)		(276)
Anomocaridae	€ (IMid)	— Θ (Trem)	(148,263,379)
Apatokephalopsidae	Θ (Trem)		(394)
Asaphidae	€ (uMid)	— Θ (Ashg)	(132,172,173)
Asaphiscidae	€ (IMid)	— Θ (Trem)	(173,263,379)
Aulacodigmatidae	€ (uMid)	— € (Dres)	(226)
Auritamidae	€ (uMid)	— € (Dres)	(226)
Avoninidae	€ (uMid)	— € (Fran)	(208)
Bolaspidae	€ (mMid)	— € (uMid)	(110,193)
Burlingiidae	€ (mMid)	— € (Fran)	(87,132,172,173)
Catillicephalidae	€ (uMid)	— € (Trep)	(148,349)
Ceratopygidae	€ (uMid)	— Θ (Trem)	(132,173,208)
Cheilocephalidae	€ (Dres)	— € (Fran)	(132,208)
Conocoryphidae	€ (Atda)	— Θ (Trem)	(110,132,173,193)
Coosellidae	€ (uMid)	— € (Dres)	(110,191,208)
Crepicephalidae	€ (IMid)	— € (Dres)	(148,173,263)
Cyclopygidae	Θ (Trem)	— Θ (Ashg)	(132)
Damesellidae	€ (uMid)	— € (Dres)	(110,173,227)
Decoroproctidae	S (I)	— D (Sieg)	(2)
Diceratocephalidae	€ (Dres)		(170,173,381)
Dikelocephalidae	€ (Dres)	— € (Trep)	(110,132,208)
Dikelocephalinidae	Θ (Trem)	— Θ (Llvi)	(132)
Dionididae	Θ (Llvi)	— Θ (Ashg)	(132)
Dokimocephalidae	€ (Dres)	— Θ (Trem)	(110,132,379)
Elviniidae	€ (Dres)	— Θ (Trem)	(132,173,208,315)
Emmrichellidae	€ (Boto)	— Θ (Trem)	(262,378)
Endymioniidae	Θ (Llvi)	— Θ (Cara)	(132)
Entomaspidae	€ (Trep)	— Θ (Trem)	(208)
Erixaniidae	€ (Dres)		(226)
Eurekiidae	€ (Fran)	— Θ (Trem)	(110,132,208,379)
Hapalopleuridae	Θ (Trem)	— Θ (Cara)	(132,378)
Harpidae	Θ (Trem)	— D (Fras)	(132)

Harpididae	€ (Fran)	— Θ (Aren)	(87,132,208)
Holotrachelidae	Θ (Ashg)	—	(132)
Housiidae	€ (Dres)	— € (Fran)	(110,132)
Hungaiidae	€ (Trep)	— Θ (Ashg)	(208,378)
Hypermecaspidae	Θ (Aren)	— Θ (m)	(208)
Idahoiidae	€ (Dres)	— € (Trep)	(110,132,208,227)
Illaenidae	Θ (Trem)	— S (u)	(132,379)
Illaenuridae	€ (Fran)	— € (Trep)	(110,132,208)
Ityophoridae	Θ (Cara)	— Θ (Ashg)	(132)
Kaolishaniidae	€ (Dres)	— € (Trep)	(193,195,314)
Kingstoniidae	€ (uMid)	— Θ (Trem)	(170,183,379)
Komaspididae	€ (uMid)	— Θ (Cara)	(132,172,312,378)
Lecanopygidiae	€ (Fran)	— Θ (Aren)	(110,132,208)
Leiostegiidae	€ (uMid)	— Θ (Trem)	(172,173,378)
Liostracinidae	€ (uMid)	— € (Trep)	(172,173,208)
Loganellidae	€ (Fran)	— € (Trep)	(87,208)
Lonchocephalidae	€ (uMid)	— € (Trep)	(191,208,227)
Marjumiidae	€ (mMid)	— Θ (Trem)	(132,172,173)
Menomoniidae	€ (uMid)	— € (Fran)	(87,110,191,208)
Missisquoiidae	€ (Fran)	— Θ (Trem)	(313,378,394)
Nepeidae	€ (uMid)	— € (Dres)	(132,227)
Nileidae	€ (Trep)	— Θ (Ashg)	(5,379)
Norwoodiidae	€ (uMid)	— Θ (Trem)	(191,227,378)
Olenidae	€ (Dres)	— Θ (Cara)	(132,193,375)
Orometopidae	Θ (Trem)	— Θ (Aren)	(132)
Pagodiidae	€ (uMid)	— € (Trep)	(132,170,173,379)
Papyriaspidae	€ (lMid)	— € (Trep)	(170,173,349,381)
Parabolinoididae	€ (Dres)	— € (Fran)	(110,208)
Phillipsiidae	D (Fame)	— P (Guad)	(132)
Phillipsinellidae	Θ (Ashg)	—	(132,378)
Placosematidae	€ (uMid)	— € (Dres)	(226)
Plectrigeridae	€ (uMid)	— € (Dres)	(226)
Plethopeltidae	€ (uMid)	— Θ (Trem)	(110,191,378)
Polycyrtaspidae	€ (Dres)	—	(226)
Pterocephalidae	€ (Dres)	— € (Trep)	(132,208,313)
Ptychaspidae	€ (Fran)	— Θ (Trem)	(110,132,195)
Ptychopariidae	€ (Boto)	— Θ (Trem)	(110,148,262,378)
Raphiophoridae	Θ (Aren)	— S (u)	(132)
Raymondinidae	€ (uMid)	— € (Trep)	(110,208)
Remopleurididae	€ (uMid)	— Θ (Ashg)	(132,148)
Rhyssometopidae	€ (uMid)	— € (Dres)	(226)
Saukiidae	€ (Dres)	— € (Trep)	(132,195,208,227)
Shirakiellidae	€ (Fran)	—	(172,173,195)
Shumardiidae	€ (Fran)	— Θ (Ashg)	(132,313,350)
Solenopleuridae	€ (Boto)	— Θ (Trem)	(110,262,263,379)
Styginiidae	Θ (Aren)	— Θ (Ashg)	(132)
Taihungshaniidae	Θ (Trem)	— Θ Llvi	(132)

Telephinidae	Θ (Llvi)	— Θ (Ashg)	(132)
Thysanopeltidae	Θ (Cara)	— D (Fras)	(132)
Tricrepicephalidae	€ (uMid)	— € (Dres)	(110,191,227)
Trinucleidae	Θ (Aren)	— Θ (Ashg)	(132)
Tsinaniidae	€ (Fran)	— € (Trep)	(172,173,381)
Or. Proetida			
Aulacopleuridae	Θ (Aren)	— C (Bash)	(103,132)
Bathyuridae	Θ (Trem)	— Θ (Cara)	(103,132)
Brachymetopidae	S (m)	— C (Step)	(103,132,138)
Celmidae	Θ (Llvi)		(103,132)
Dimeropygidae	Θ (Aren)	— Θ (Ashg)	(103,132)
Glaphuridae	Θ (Aren)	— Θ (Ashg)	(103)
Proetidae	Θ (Aren)	— P (Leon)	(103,132)
Or. Phacopida			
Calmoniidae	S (m)	— D (Fras)	(2,208)
?Calymenidae	Θ (Trem)	— D (Eife)	(2,132,208)
Cheiruridae	€ (u)	— D (Give)	(2,132)
Dalmanitidae	Θ (Aren)	— D (Fras)	(132)
Enocrinuridae	Θ (Trem)	— D (Gedi)	(2,132,379)
?Homalonotidae	Θ (Trem)	— D (Fras)	(2,132)
Monorakidae	Θ (Cara)	— Θ (Ashg)	(132)
Phacopidae	S (l)	— D (Fame)	(132)
Pliomeridae	Θ (Trem)	— Θ (Ashg)	(132)
Pseudotrinitidae	D (Emsi)	— D (Eife)	(2)
Pterygometopidae	Θ (Aren)	— S (l)	(132)
Synphoriidae	S (m)	— D (Eife)	(2,138)
Or. Lichida			
Lichakephalidae	Θ (Trem)		(132)
Lichidae	Θ (Trem)	— D (Fras)	(132,208)
Or. Odontopleurida			
Eoacidaspidae	€ (uMid)	— € (Fran)	(148,283)
Eodontopleuridae	€ (Boto)		(393)
Odontopleuridae	Θ (Aren)	— D (Fras)	(132)
Or. Nektaspidida			
Naraoiidae	€ (mMid)		(374)
Incertae sedis			
Amgaspididae	€ (lMid)	— € (uMid)	(148,173,263)
Edelsteinaspidae	€ (lMid)	— € (mMid)	(148,173,263,288)
Granulariidae	€ (Boto)	— € (uMid)	(148,261)
Hanburiidae	€ (mMid)		(148)
Isocolidae	Θ (Aren)	— Θ (Ashg)	(5,378)
Jakutidae	€ (Boto)	— € (lMid)	(148,173,263)
Myindidae	Θ (Trem)		(379)
Namanoiidae	€ (lMid)	— € (uMid)	(148,173)

Sarkiidae	$\Theta$ (m)	(208)	
Cl. Merostomoidea			
[Classification after Bergström (1971, 1980).]			
Or. Emeraldellida			
Emerallidae	$\epsilon$ (mMid)	(41,61)	
Or. Limulavida			
Sidneyiidae	$\epsilon$ (mMid)		
Or. Cheloniellida			
Cheloniellidae	D (Sieg)	(132,208)	
Or. uncertain			
<i>Helmetia</i>	$\epsilon$ (mMid)	(377)	
Paleomeridae	$\epsilon$ (Atda)	(21)	
Strabopidae	$\epsilon$ (Fran) — $\Theta$ (Cara)	(132,254)	
Cl. Marrelomorpha			
[Classification after Bergström (1980).]			
Or. Mimetasterida			
<i>Mimetaster</i>	$\epsilon$ (mMid) — D (Sieg)	(132,377)	
Or. Acercostraca			
Vachoniidae	D (Sieg)	(132)	
Or. Cyclina (=Cycloidea)			
Cyclidae	C (Tour)	— Tr (Nori)	(132)
Hemitrochiscidae	P (u)	— Tr (u)	
Mesoprosopidae	Tr (u)		
Crustacea			
Cl. Incertae sedis			
?Or. Leanchoiliida			
Leanchoiliidae	$\epsilon$ (mMid)	(377)	
Waptiida			
Waptiidae	$\epsilon$ (mMid)	(377)	
Cl. Cephalocarida			
*?Dala	$\epsilon$ (Dres)	(215)	
Cl. Branchiopoda			
[Classification from the <i>Treatise</i> , Pt. R. Only marine families are listed; three extant marine families without known fossil representatives are not listed.]			

\*May be related to unlisted extant families.

Incertae sedis		
?Parvancorinidae	V (u)	(208)

Or. Diplostraca (?Conchostraca)		
incertae sedis	D (Gedi)	— D (Emsi) (132)

Or. Anostraca		
Gilsonicarididae	D (Sieg)	(132)

## Cl. Ostracoda

[Classification and stratigraphic ranges from the *Treatise*, Pt. Q, and *The Fossil Record*, with emendations as indicated. Six extant families without known fossil representatives are not listed; exclusively nonmarine families also are not listed.]

## Or. Archaeocopida (= Bradoriida)

Alutidae	€ (Atda)	(170,195,228)
Beyrichonidae	€ (Atda)	— Θ (Trem) (132,228,295,297)
Bradoriidae	€ (uTom)	— € (Fran) (70,74,228,314)
Comptalutidae	€ (IMid)	— € (uMid) (228,314)
Indianidae	€ (Boto)	— € (mMid) (228,295,314)
Monasteriidae	€ (mMid)	(160)
Oepikalutidae	€ (mMid)	— € (uMid) (160)
Svealutidae	€ (mMid)	— € (Dres) (160,228,296,314)

## Or. Leperditicopida

Isochilinidae	Θ (Trem)	— D (Give)
Leperiditiidae	Θ (Aren)	— D (Give)
incertae sedis	€ (l)	(177)

## Or. Palaeocopida

Aechminellidae	S (m)	— P (Sakm) (144)
Aechminidae	Θ (Cara)	— D (Give) (208)
Amphissitidae	D (Emsi)	— P (Guad) (125)
Aparchitidae	Θ (Aren)	— C (Step)
Arcyzonidae	D (Eife)	— D (Fras) (125, 224)
Bassleratiidae	Θ (Aren)	— Θ (Cara)
Beyrichiidae	Θ (m)	— C (Tour) (26)
Beyrichiopsidae	D (Sieg)	— P (Guad) (125)
Bolliidae	Θ (Aren)	— D (Give) (125,298)
Cardiniferellidae	C (Namu)	
Chilobolbinidae	Θ (m)	— S (m)
?Conchoprimitiidae	Θ (l)	— Θ (Ashg) (299)
Craspedobolbinidae	S (l)	— D (Fame) (159)
?Cryptophyllidae	Θ (Cara)	— J (u) (299)
Ctenoloculinidae	S (m)	— C(l) (27)
Drepanellidae	Θ (Cara)	— D (Fame)
?Eridoconchidae	Θ (Cara)	— D (Fame) (159,299)
Eurychilinidae	Θ (Aren)	— S (l) (298)
Geisinidae	D (Eife)	— P (Sakm) (125)

Glyptopleuridae	D (Give)	— P (Sakm)	
Hollinellidae	S (u)	— Tr (Indu)	(27)
Hollinidae	D (m)	— C (Step)	(27)
Jaanussoniidae	Θ (m)	— D	(299)
Judahellidae	Tr (Indu)	— Tr (Ladi)	(325)
Kellettinidae	C (Tour)	— P (Guad)	
Kirkbyellidae	S (m)	— C (Mosc)	
Kirkbyidae	D (Eife)	— Tr (Indu)	(224,325)
Kloendenellidae	S (l)	— C (Step)	
Knightinidae	D (u)	— Tr (Indu)	(325)
Leperditellidae	Θ (Aren)	— P (Leon)	
Lichwiiniidae	D (Give)	— P (Guad)	
Miltonellidae	P (Guad)		
Nodellidae	D (Emsi)	— D (Fame)	(125)
Oepikellidae	Θ (Cara)	— Θ (Ashg)	
Paraparachitidae	D (Gedi)	— P (Guad)	(125)
Piretellidae	Θ (Aren)	— Θ (Ashg)	(298)
Placideidae	D (Give)	— P (Guad)	
Pribylitidae	S (u)	— D (Fame)	(125)
Primitiopsidae	Θ (Llde)	— D (Give)	
Punciidae	Θ (Aren)	— R	
Quadrijugatoridae	Θ (Aren)	— Θ (Cara)	
Richinidae	Θ (Cara)	— D (Fras)	(125)
Sansabellidae	C (Vise)	— C (Step)	
?Schmidtellidae	Θ (Cara)	— D (m)	(298,299)
Scrobiculidae	D (Emsi)	— P (Guad)	(125)
Sigmoopsidae	Θ (Aren)	— Θ (Ashg)	
Tetradellidae	€ (lMid)	— S (l)	(177)
Tvaerenellidae	Θ (Aren)	— Θ (Ashg)	
Youngiellidae	D (Give)	— C (Step)	
Zygbolbidae	Θ (m)	— S (m)	
 Or. Myodocopida			
Bolbozooidae	S (l)	— D (Fame)	
Cyprellidae	C (Vise)		
Cypridinellidae	D	— C (Vise)	
Cyprosinidae	D (Gedi)	— D (Fame)	
Entomoconchidae	D (Gedi)	— C (Step)	
Entomozoidae	Θ (Llvi)	— P (Asse)	
Polycopidae	C (Vise)	— R	
Rhombinidae	C (Tour)	— C (Vise)	
Thaumatocyprididae	J (Bajo)	— R	
 Or. Podocopida			
Acronotellidae	Θ (Cara)	— S (u)	
Bairdiidae	Θ (Ashg)	— R	
Bairdiocyprididae	S (l)	— P	
Barychilinidae	Θ (Cara)	— D (Fras)	(159)

•

Beecherellidae	S (m)	— D (Fame)	(125,144)
Berounellidae	S (u)	— D (Fame)	
Brachycytheridae	J (Kimm)	— R	
Bufinidae	D (Eife)	— C (Penn)	(125)
Bythocytheridae	Θ (Ashg)	— R	
Cavellinidae	S (u)	— Tr (Indu)	(325)
Cyprididae	J (Bath)	— R	
Cytherellidae	S (m)	— R	(144)
Cytherettidae	K (Coni)	— R	
Cytheridae	K (Neoc)	— R	(4)
Cytherideidae	C (Mosc)	— R	
Cytherissinellidae	Tr (Indu)	— Tr (Olen)	
Cytheruridae	J (Bajo)	— R	
Egorovitinidae	C (Vise)		(127)
Healdiidae	D (Gedi)	— K (Apti)	(125)
Hemicytheridae	T (Eo-l)	— R	
Krausellidae	Θ (Llde)	— D (Fras)	(159)
Leguminocythereididae	T (Pale)	— R	
Leptocytheridae	T (Eo-u)	— R	
Loxoconchidae	J (Bajo)	— R	
Macrocypridiidae	J (Oxo)	— R	
Pachydomellidae	S (m)	— D (Fame)	(224)
Paracyrididae	J (Hett)	— R	
Paradoxostomatidae	K (Albi)	— R	(4)
Pectocytheridae	K (Maes)	— R	
Permianidae	P (Dzhu)		
Pontocypridiidae	K (u)	— R	(208)
Progonocytheridae	J (Sine)	— K (Maes)	
Protocytheridae	J (Sine)	— R	
Quasillitidae	D (Gedi)	— P (Guad)	(125)
Ropolonellidae	D (Sieg)	— D (Fras)	(125)
Schizocytheridae	K (Ceno)	— R	
Schulerideidae	J (Toar)	— R	
Sinusuellidae	P (Dzhu)		
Thlipsuridae	S (l)	— D (Fame)	(125,224)
Tomelliidae	P (Dzhu)		
Trachyleberididae	J (Bajo)	— R	
Xestoleberididae	K (Sant)	— R	
Incertae sedis			
Buregiidae		D (Give)	
?Cl. Euthycarcinoidea			
[May not be crustaceans.]			
Or. Euthycarinida			
Euthycarinidae	C (Mosc)	— Tr (Olen)	(132,301)
Sottyxeridae	C (Mosc)	— C (Step)	(301)

## Cl. Copepoda

[Approximately 30 extant marine families; freshwater families not considered.]

Calanoida -			
(?Ascomyzontidae)	K (Apti)	- R	(72)
incertae sedis	J (Call)		(132,278)

## Cl. Cirripedia

[Classification from the *Treatise*, Pt. R; stratigraphic ranges from *The Fossil Record*. Approximately 17 extant families without fossil representatives (including all in the Rhizocephala and Apoda) are not listed.]

## Or. Thoracica

<i>Priscanermarinus</i>	Є (mMid)		(51a)
Balanidae	T (Eo-l)	- R	
Brachylepadidae	J (Tith)	- T (Mi-u)	
Chthamalidae	K (Camp)	- R	
Cyprilepadidae	S (u)		
Lepadidae	T (Eo-m)	- R	
Oxynaspidae	T (Eo-m)	- R	
Poecilasmatidae	T (Eo-u)	- R	
Praelepadidae	C (Mosc)		
Scalpellidae	C (u)	- R	
Stramentidae	K (Albi)	- K (Seno)	
Verrucidae	K (Ceno)	- R	

## Or. Acrothoracica

Trypetesidae	C (Step)	- R	
--------------	----------	-----	--

## Or. Ascothoracica

incertae sedis	K (Turo)	- R	(181)
----------------	----------	-----	-------

## Cl. Malacostraca

[Classification and stratigraphic ranges based on the *Treatise*, Pt. R, with emendations as indicated. Only marine families are listed, and approximately 90 extant families without known fossil representatives are not listed.]

## Or. Leptostraca

Nebaliidae	P (Guad)	- R	(132)
------------	----------	-----	-------

## ?Or. Canadaspidida

Canadaspididae	Є (mMid)		(36,39)
Perspicarididae	Є (mMid)	- Є (uMid)	(37,276a)

## Or. Hymenostraca

Hymenocarididae	Є (Boto)	- Θ (Aren)	(39,264)
-----------------	----------	------------	----------

## Or. Hoplostraca

Sairocarididae	C (Vise)	- C (Mosc)	(301,303)
----------------	----------	------------	-----------

## Or. Archaeostraca

Aristozoidae	S (u)	- D (Eife)	(279)
--------------	-------	------------	-------

•

Austriocarididae	D (Emsi)	— Tr (Carn)	(279)
Ceratiocarididae	Θ (Trem)	— C (Mosc)	(46,303)
Echinocarididae	D (Sieg)	— C (Tour)	(46,279)
Ohiocarididae	D (Fame)		(279)
Pephricarididae	D (Fame)		(279)
Rhinocarididae	D (Sieg)	— C (Mosc)	(279)
Or. Incertae sedis			
<i>Anomalocaris</i>	Є (Boto)	— Є (mMid)	(38)
<i>Carnarvonia</i>	Є (mMid)		
<i>Protocaris</i>	Є (l)	— Є (mMid)	(35)
Discinocarididae	Θ (m)	— S (u)	
Peltocarididae	Θ (Ashg)	— Tr (Carn)	
Tuzoïidae	Є (Atda)	— Є (uMid)	(61,121,276a)
Or. Palaeostomatopoda			
Perimecturidae	D (u)	— C (Namu)	(300)
Or. Stomatopoda			
Gonodactylidae	K (l)	— R	(300)
Lysiosquillidae	T (Eo)	— R	
Sculdidae	J (Tith)	— K (u)	
Squilliidae	K (l)	— R	
Tyrannophontidae	C (Namu)	— C (Mosc)	(300,303,307)
Or. Aeschronectida			
<i>Aenigmatocarididae</i>	C (Vise)	— C (Step)	(307)
<i>Aratidecthidae</i>	C (Vise)	— C (Mosc)	
<i>Kallidecthidae</i>	C (Bash)	— C (Step)	(300,307,308)
Or. Eocaridacea			
<i>Anthracophausiidae</i>	C (Vise)	— C (Mosc)	(300,303)
<i>Eocarididae</i>	D (Give)	— D (Fame)	(300)
<i>Essoidiidae</i>	C (Mosc)		(303)
Or. Belotelsonidea			
Belotelsonidae	C (Vise)	— C (Mosc)	(303,307)
Or. Waterstonellidea			
Waterstonellidae	C (Vise)		
Or. Decapoda			
Albuneidae	T (Ol-l)	— R	
Astacidae	J (u)	— R	
Atelecyclidae	T (Eo-l)	— R	
Atyidae	T (Ol-u)	— R	
Axiidae	J (Toar)	— R	
Calappidae	K (Haut)	— R	(383)
Callianassidae	J (u)	— R	
Cancridae	T (Eo-m)	— R	
Cancrinidae	J (Tith)		

MILWAUKEE PUBLIC MUSEUM

---

# Contributions

in  
BIOLOGY  
and  
GEOLOGY

---

Number 51

November 29, 1982

A Compendium of Fossil Marine Families

J. John Sepkoski, Jr.

Carcineretidae	K (Apti)	— K (Maes)	(383)
Carpiliidae	K (Ceno)	— R	(383)
Coenobitidae	T (Mi)	— R	
Coleiidae	Tr (u)	— K (l)	
Cymonomidae	K (Haut)	— R	(383)
Dakoticancridae	K (Seno)		
Dorippidae	K (Albi)	— R	
Dromiidae	K (Albi)	— R	(383)
Dynomenidae	J (Tith)	— R	(383)
Eocarcinimidae	J (Plie)		
Erymidae	P (Guad)	— K (Camp)	(9,305)
Eryonidae	J (l)	— K (l)	
Galatheidae	J (Bajo)	— R	
Geryonidae	T (Eo-l)	— R	
Glypheidae	P (Sakm)	— R	(101a,300)
Goneplacidae	T (Pale)	— R	
Grapsidae	T (Eo-m)	— R	
Homolidae	J (Tith)	— R	
Laomediidae	T (Mi)	— R	
Latreilliidae	K (Ceno)	— R	(383)
Leucosiidae	T (Eo-m)	— R	
Majidae	T (Eo-l)	— R	
Mecochiridae	Tr (l)	— K (Camp)	(9)
Nephropidae	J (m)	— R	
Ocypodidae	T (Eo-m)	— R	
Oplophoridae	J (Call)	— R	
Paguridae	J (Plie)	— R	
Palaemonidae	T (Eo-l)	— R	
Palaeopalaemonidae	D (Fame)		(93,300,306)
Palaeopemphicidae	P (Guad)		(132)
Palicidae	K (Albi)	— R	
Palinuridae	J (l)	— R	
Parastacidae	T (Eo)	— R	
Parthenopidae	T (Eo-m)	— R	
Pemphicidae	Tr (m)		
Penaeidae	Tr (u)	— R	(132)
Pinnotheridae	T (Eo-l)	— R	(52)
Platychelidae	Tr (u)		
Polychelidae	J (m)	— R	
Porcellanidae	K (Ceno)	— R	
Portunidae	T (Eo-l)	— R	
Potamidae	T (Mi-u)	— R	
Prosopidae	J (Bajo)	— R	(383)
Raninidae	K (Albi)	— R	
Retroplumidae	T (Eo-m)	— R	
Scyllaridae	K (Albi)	— R	
Tetrachelidae	Tr (u)		

Thalassinidae	Q (Plei)	— R	
Tymolidae	T (Eo-m)	— R	
Udorellidae	J (Tith)		
Uncinidae	J (Toar)		
Xanthidae	K (Apti)	— R	(383)
Or. Mysidacea (marine only)			
Jerometichenoriidae	P (Sakm)		
Mysidae	T (Mi-l)	— R	
Notocarididae	P (Guad)		(300)
Pygocephalidae	C (Vise)	— P (l)	(305)
Tealliocarididae	C (Tour)	— C (Vise)	(303)
Or. Cumacea (extant)			
<i>Nannastacus</i>	P (Guad)		(132)
<i>Palaeocuma</i>	Tr (m)		
Or. Tanaidacea (extant)			
<i>Charassocarcinus</i>	J (Bajo)		
<i>Ophthalmapseudes</i>	P (Guad)		(132)
<i>Palaeotanais</i>	J (Bajo)		(208)
Anthracocarididae	C (Tour)		(303)
Cryptocarididae	C (Mosc)		
Or. Isopoda			
Amphisopidae	P (Guad)	— R	
Archaeoniscidae	J (u)		
Armadillidiidae	T (Mi-u)	— R	
Cirolanidae	T (Mi-u)	— R	
Epicaridea	J (Oxfo)	— R	(62)
Idoteidae	T (Ol-l)	— R	
Oniscidae	T (Eo-u)	— R	
Palaeophreatoicidae	C (Mosc)	— P (Guad)	(303,305)
Porcellionidae	T (Eo-u)	— R	
Sphaeromidae	K (u)	— R	
Trichoniscidae	T (Eo-u)	— R	
Urdidae	J (Bath)	— K (Apti)	(243)
Or. Amphipoda			
Gammaridae	T (Eo-u)	— R	

## Uniramia

## Cl. Tardigrada (3 extant marine families)

? <i>Aysheaia</i>	Є (mMid)	(375,376)
? <i>Xenusion</i>	Є (uTom)	(145)

## Arthropod incertae sedis

## Cl. Incertae sedis

## Or. Bostrichopodida

Bostrichopodidae	C (Tour)	
Incertae sedis		
<i>Actaeus</i>	€ (mMid)	(377)
<i>Alalcomenaeus</i>	€ (mMid)	(377)
<i>Bohemillidae</i>	Θ — Θ (Ashg)	(208)
<i>Burgessia</i>	€ (mMid)	(377)
<i>Corcoraniidae</i>	Θ (Trem)	(151)
<i>Habelia</i>	€ (mMid)	(377)
<i>Kablikia</i>	Θ (Llde)	(132,208)
<i>Marriocarida</i>	€ (mMid) — Θ (Ashg)	(208)
<i>Molaria</i>	€ (mMid)	(377)
<i>Mollisoniidae</i>	€ (mMid)	(377)
<i>Odaraidae</i>	€ (mMid)	(377)
<i>Pahvantia</i>	€ (uMid)	(276a)
<i>Plenocaris</i>	€ (mMid)	(377)
? <i>Pomerania</i>	€ (uTom)	(162,186)
<i>Praecambrididae</i>	V (u)	(123,208)
<i>Protocarididae</i>	€ (mMid) — € (uMid)	(40,276a,377)
<i>Serracaris</i>	€ (Boto)	(38)
<i>Skania</i>	€ (mMid)	(377)
<i>Tegopelte</i>	€ (mMid)	(377)
<i>Vendomiidae</i>	V (u)	(162,231)
<i>Yohoiidae</i>	€ (mMid) — € (u)	(132,208)

## PHORONIDA (1 extant family)

? <i>Helicosyrinx</i>	Θ	(62)
incertae sedis	K (Maes) — R	(208)

## BRYOZOA

[Classification based on the *Treatise*, Pt. G, and stratigraphic ranges on *The Fossil Record*, with emendations as indicated; references to the *Treatise* (208) are to Part A. Approximately 37 extant families without fossil records are not listed.]

## Cl. Stenolaemata

Or. Cyclostomata		
Cavidae	J (Bath) — K (Maes)	
Cerymboporidae	K (Ceno) — R	
Clausidae	K (Vala) — K (Maes)	
Corynotrypidae	Θ (Aren) — P (u)	(84a)
Crisiidae	K (Ceno) — R	
Crownoporidae	Θ (Livi) — Θ (Cara)	(84a)
Cytididae	J (Kimm) — R	
Diaperoeciidae	J (Bath) — R	
Diastoporidae	Θ (Cara) — R	

Eleidae	J (Bath)	— T (Dani)
Entalophoridae	J (Plie)	— R
Frondiporidae	J (Bath)	— R
Hederellidae	S (m)	— C (Step) (84a)
Heteroporidae	Tr (Nori)	— R
Horneridae	K (Apti)	— R (359)
Leiosoeciidae	J (Bath)	— T (Eo-l)
Lichenoporidae	K (Vala)	— R
Lobosoeciidae	K (Ceno)	
Multisparsidae	J (Bajo)	— K (l)
Oncousoeciidae	J (Plie)	— R
Petaloporidae	K (Haut)	— R
Phaceloporidae	S (l)	— D
Plagioeciidae	J (Bath)	— R
Reptariidae	S (m)	— D (m) (84a)
Semiceidae	K (Ceno)	— K (Maes) (359)
Sulcocavidae	K (Coni)	— K (Maes) (359)
Terviidae	T (Eo-m)	— R
Theonoidae	J (Bajo)	— R
Tretocycloeciidae	K (Ceno)	— R
Tubuliporidae	D (Eife)	— R
 Or. Cystoporata		
Actinotrypidae	C (Vise)	— P (Guad) (208,285)
Ceramoporidae	Θ (Aren)	— D (Gedi) (229)
Constellariidae	Θ (Llde)	— S (m)
Dianulitidae	Θ (Aren)	— Θ (Ashg) (200)
Etherellidae	P (Leon)	— P (Guad) (208,285a)
Fistuliporidae	Θ (Llvi)	— P (Dzhu) (209,285)
Goniocladiidae	D (u)	— P (Dzhu) (209,229,285)
Hexagonellidae	D (m)	— P (Dzhu) (209,285)
Rhinoporidae	Θ (Llvi)	— S (m)
Sulcoreteporidae	D (Gedi)	— P (Guad) (285)
 Or. Trepostomata		
Aisenvergiidae	C (Tour)	
Amplexoporidae	Θ (Cara)	— S (u)
Anisotrypidae	D (u)	— P (Dzhu) (285a)
Araxoporidae	C (Tour)	— Tr (Indu) (285)
Atactotoechidae	Θ (Aren)	— D (Fras) (23b)
Batostomellidae	Θ (Aren)	— S (u)
Cycloporidae	C (Vise)	(285a)
Dittoporidae	Θ (Aren)	— Θ (Ashg)
Dyscritellidae	D (u)	— P (Dzhu) (285a)
Eridotrypellidae	D (Give)	— P (Guad) (209,229)
Halloporidae	Θ (Llde)	— D (Emsi)
Heterotrypidae	Θ (Cara)	— D (Fras)
Leioclemidae	C (Tour)	— P (Asse) (285a)

Mesotrypidae	Θ (Llde)	— Θ (Ashg)	(229)
Monticuliporidae	Θ (Llvi)	— S (u)	
Stenoporidae	D (Gedi)	— P (Dzhu)	(285a)
Trematoporidae	Θ (Aren)	— D (Eife)	
Ulrichotrypellidae	D (u)	— P (Guad)	(285)
 Or. Cryptostomata			
Acanthocladidae	Θ (Llvi)	— P (Guad)	(285a)
Arhrostylidae	Θ (Cara)	— P (Asse)	(285a)
Bactroporidae	D (Give)		
Escharoporidae	Θ (Llde)	— C (Vise)	
Fenestellidae	Θ (Cara)	— Tr (Indu)	
Fenestraliidae	C (Tour)	— P (Guad)	(209,285)
Girtyoporidae	C (Vise)	— Tr (Indu)	(285a)
Hyphasmoporidae	D (u)	— P (Dzhu)	(209,285)
Intraporidae	D (m)	— C (Vise)	
Nikiforovellidae	D (m)	— P (Guad)	(285)
Phragmopheridae	C (Step)		(285a)
Phylloporinidae	Θ (Llvi)	— P (Leon)	(285a)
Polyporidae	Θ (Cara)	— Tr (Indu)	(209,285)
Ptilodictyidae	Θ (Llde)	— D (Sieg)	
Rhabdomesidae	S (u)	— P (Guad)	(285a)
Rhinidiptyidae	Θ (Cara)	— S	(229)
Rhomboporidae	C (Vise)	— P (u)	
Septatoporidae	C (Tour)	— P (Dzhu)	(209,285)
Steptoporidae	D (u)	— P (Guad)	(285a)
Stictoporellidae	Θ (Aren)	— C (Namu)	
Timanodictyidae	P (Asse)	— P (Guad)	(208,229,285a)
Virgatellidae	Θ (m)		
?Worthenoporidae	C (Tour)	— C (Vise)	

## Cl. Gymnolaemata

Or. Ctenostomata			
Arachnidiidae	K (Maes)	— R	(361)
Ascodictyidae	S (l)	— P	
Cookobryozoonidae	T (Mi-m)	— T (Plio)	(246)
Immergentiidae	T (Mi-l)	— R	(246)
Orbignyoporidae	K (Maes)	— T (Plio)	(246)
Penetrantiidae	Tr (Rhae)	— R	(246)
Ropalonariidae	Θ (Cara)	— P (Sakm)	(84a)
Spathiporidae	J (Call)	— R	(246)
Terebriporidae	K (Maes)	— R	(246)
Vesiculariidae	K (Maes)	— R	
Vinellidae	Θ (Aren)	— K (Camp)	
Voigtellidae	K (Camp)	— K (Maes)	(246)

## Or. Cheiostomata

Adeonidae	T (Eo-l)	— R
-----------	----------	-----

Aeteidae	T (Eo-u)	— R
Alysidiidae	T (Mi-m)	— R
Andrioporidae	K (Ceno)	— T (Dani)
Arachnopusiidae	T (Eo-m)	— R
Aspidostomatidae	K (Ceno)	— R (84)
Bifaxariidae	T (Eo-u)	— R (43)
Bitectiporidae	T (Ol-u)	— R
Calloporidae	K (Albi)	— R
Calpensiidae	T (Eo-m)	— R
Calpidoporidae	K (Ceno)	— K (Sant)
Catenicellidae	T (Eo-u)	— R (115a)
Cellariidae	K (Sant)	— R
Celleporidae	K (Camp)	— R
Chaperiellidae	T (Ol-l)	— R
Cheiloporinidae	K (Sant)	— R
Conescharellinidae	T (Eo-u)	— R (115a)
Coscinopleuridae	K (Sant)	— T (Eo-l)
Crepidacanthidae	T (Eo-m)	— R
Cribrilinidae	K (Sant)	— R
Ctenoporidae	K (Ceno)	
Cyclicoporidae	K (Maes)	— R
Disheloporidae	K (Coni)	— K (Sant)
Dysnoetoporidae	K (Camp)	— K (Maes)
Electridae	J (Tith)	— R
Eurystomellidae	Q (Plei)	— R
Exochellidae	K (Sant)	— R
Farciminariidae	K (Maes)	— R
Fusicellariidae	K (Turo)	— T (Mi)
Gigantoporidae	K (Maes)	— R
Hiantoporidae	K (Sant)	— R
Hincksinidae	K (Camp)	— R
Hippoporinidae	K (Camp)	— R
Hippothoidae	K (Ceno)	— R
Lekythoporidae	T (Eo-u)	— R
Lunulitidae	K (Turo)	— R
Mamilloporidae	K (Maes)	— R
Membranicellariidae	K (Maes)	— R
Membraniporidae	K (Ceno)	— R
Microporellidae	T (Mi-u)	— R
Microporidae	K (Sant)	— R
Mucronellidae	K (Camp)	— R
Myagroporidae	K (Coni)	— K (Sant)
Myriozoidae	T (Eo-u)	— R (115a)
Nephroporidae	K (Maes)	
Onychocellidae	K (Ceno)	— R
Orbitaliporidae	K (Camp)	— R
Otoporidae	K (Ceno)	— K (Camp)

Parmulariidae	K (Camp)	— R
Pasytheidae	T (Eo-l)	— R
Pelmatoporidae	K (Turo)	— R
Petraliidae	T (Eo-u)	— R
Phylactellidae	T (Pale)	— R
Phylactelliporidae	T (Eo-m)	— R
Platyglenidae	K (Maes)	— T (Dani)
Porinidae	K (Turo)	— R
Prostomariidae	T (Mi)	
Reteporidae	T (Eo-l)	— R
Rhacheoporidae	K (Coni)	— K (Maes)
Schizoporellidae	K (Maes)	— R
Scrupariidae	T (Eo-l)	— R
Scrupocellaridae	T (Eo-u)	— R
Setosellidae	K (Maes)	— R
Steginoporellidae	T (Eo-m)	(43)
Stomachetosellidae	T (Eo-m)	— R
Taractoporidae	K (Coni)	— K (Sant)
Thalamoporellidae	K (Camp)	— R
Thoracoporidae	K (Turo)	— K (Sant)
Tubucellariidae	T (Eo-m)	— R
Umbonulidae	T (Eo-m)	— R
Wawaliidae	J (u)	— R
		(84)

## BRACHIOPODA

## Cl. Inarticulata

[Classification from the *Treatise*, Pt. H; stratigraphic ranges augmented with *The Fossil Record* and a number of other sources, as listed.]

Or. Lingulida		
Andobolidae	Θ (Ashg)	
Craniopsidae	Θ (Cara)	— C (Tour)
Elkaniidae	Є (Fran)	— Θ (Aren) (295)
Lingulasmatidae	Θ (Cara)	— Θ (Ashg)
Lingulidae	S (m)	— R
Obolidae	Є (uTom)	— Θ (Ashg) (196,384)
Paterulidae	Θ (Aren)	— S (m)
Trimerellidae	Θ (Cara)	— S (u)
Or. Acrotretida		
Acrothelidae	Є (uTom)	— Θ (Trem) (70,252)
Acrotretidae	Є (lTom)	— S (u) (358)
Botsfordiidae	Є (Atda)	— Є (mMid) (163,314,350)
Craniidae	Θ (Llv)	— R
Curticiidae	Є (uMid)	— Є (Fran) (132,192)
Discinidae	Θ (Llde)	— R
Eoconulidae	Θ (Cara)	

Ptychopeltidae	$\Theta$ (m)		
Siphonotretidae	$\epsilon$ (uMid)	— $\Theta$ (Cara)	(192,314)
Trematidae	$\Theta$ (Cara)	— D (Gedi)	
Or. Obolellida			
Obolellidae	$\epsilon$ (Atda)	— $\epsilon$ (uMid)	(163,172,286)
Or. Paterinida			
Paterinidae	$\epsilon$ (uTom)	— $\Theta$ (Cara)	(286,287)
Or. Kutorginida			
Kutorginidae	$\epsilon$ (uTom)	— $\epsilon$ (mMid)	(132,286,351)
Yorkiidae	$\epsilon$ (Atda)	— $\epsilon$ (Boto)	(64,232,316)

## Cl. Articulata

[Classification from the *Treatise*, Pt. H, with additions from Cooper and Grant (1972-1977); stratigraphic ranges augmented with *The Fossil Record* and a number of other sources, as listed. References to the *Treatise* (208) are to Part A.]

## Or. Orthida

Angusticardiniidae	$\Theta$ (Aren)	— $\Theta$ (Ashg)	
Billingsellidae	$\epsilon$ (Boto)	— $\Theta$ (Aren)	(171,172)
Clitambonitidae	$\Theta$ (l)	— $\Theta$ (Ashg)	
Cremnorthidae	$\Theta$ (m)	— $\Theta$ (Cara)	
Dalmanellidae	$\Theta$ (Aren)	— C (l)	
Dicoelosiidae	$\Theta$ (Ashg)	— D (Fras)	(31)
Dolerorthidae	$\Theta$ (l)	— D (Emsi)	(31)
Eichwaldiidae	$\Theta$ (Cara)	— S (u)	
Enteletidae	$\Theta$ (Llde)	— P (Dzhu)	(128)
Eoothidae	$\epsilon$ (Boto)	— $\Theta$ (Aren)	(194,275)
Finkelnburgiidae	$\epsilon$ (Fran)	— $\Theta$ (Aren)	(196,297,380)
Gonambonitidae	$\Theta$ (Aren)	— $\Theta$ (m)	
Harknessellidae	$\Theta$ (Llde)	— $\Theta$ (Ashg)	(380)
Hesperonomiidae	$\Theta$ (Aren)		(282,380)
Heterorthidae	$\Theta$ (Llvi)	— S (u)	(31,380)
Hypsomyoniidae	D (m)	— D (u)	
Isogrammidae	C (l)	— P (Guad)	
Kayserellidae	D (Eife)	— D (Fras)	(31)
Kullervoidea	$\Theta$ (Llde)	— $\Theta$ (Ashg)	(380)
Leioriidae	$\epsilon$ (Boto)		(64)
Linoporellidae	$\Theta$ (Cara)	— S (m)	
Mystrophoridae	D (Eife)		(31)
Nisusiidae	$\epsilon$ (Atda)	— $\epsilon$ (uMid)	(171,275,286)
Orthidae	$\epsilon$ (m)	— P (Dzhu)	(368)
Orthidiellidae	$\Theta$ (Aren)	— $\Theta$ (Ashg)	(282)
Paurorthidae	$\Theta$ (Aren)	— $\Theta$ (m)	
Plaesiomiyidae	$\Theta$ (l)	— $\Theta$ (Ashg)	
Plectorthidae	$\Theta$ (Aren)	— S (m)	(31,380)

Polytoechiidae	Θ (Trem)	— Θ (Llvi)	(282)
Protorthidae	Є (Boto)	— Є (m)	(64)
Rhipidomellidae	S (l)	— P (Dzhu)	(368)
Saukrodictyidae	Θ (Ashg)	— S (l)	(31)
Skenidiidae	Θ (Aren)	— D (Fras)	(380)
Triplesiidae	Θ (Aren)	— S (m)	(282)
Tropidoleptidae	D (Gedi)	— D (Fras)	(31)
Tuvaellidae	S (l)		
 Or. Strophomenida			
Agelesiidae	P (Leon)		(66)
Anoplidiidae	S (u)	— D (Emsi)	(31)
Anoptambonitidae	Θ (Llde)	— Θ (Ashg)	(239a)
Aulostegidae	C (Step)	— P (Dzhu)	(208)
Bactryniidae	Tr (Rhae)		
Bimuriidae	Θ (Llde)	— Θ (Cara)	
Buxtoniidae	D (Fame)	— P (Guad)	(31,208)
Cadomellidae	J (Plie)	— J (Toar)	
Chonetellidae	P (Asse)	— P (Dzhu)	(128)
Chonetidae	Θ (Ashg)	— P (Dzhu)	
Chonostrophiidae	S (u)	— C (Tour)	
Christianiidae	Θ (Llde)	— Θ (Ashg)	
Cooperinidae	C (Penn)	— P (Guad)	(66)
Cyclacanthariidae	P (Sakm)	— P (Guad)	(66)
Davidsoniidae	D (Eife)	— D (Give)	(31,67)
Daviesiellidae	D (u)	— C (Vise)	
Dictyoclostidae	C (Tour)	— P (Dzhu)	(208)
Echinoconchidae	C (Tour)	— P (Dzhu)	(128)
Eocramatiidae	Θ (Llvi)		(380)
Eodevonariidae	D (Sieg)	— D (Eife)	(31)
Foliomenidae	Θ (Ashg)		
Gigantoprotidae	D (u)	— C (Namu)	
Hercosiidae	P (Sakm)	— P (Leon)	(66)
Institinidae	C (Vise)		
Leioproductidae	D (Give)	— P (Leon)	(31,208)
Leptaenidae	Θ (Cara)	— C (Vise)	(380)
Leptellinidae	Θ (Aren)	— S (u)	
Leptestiidae	Θ (Aren)	— Θ (Ashg)	
Linoproductidae	C (Tour)	— P (Dzhu)	(128)
Lyttoniidae	C (Step)	— P (Dzhu)	(128)
Marginiferidae	C (Vise)	— P (Dzhu)	(128)
Meekellidae	Θ (Cara)	— P (Dzhu)	(128)
Oepikinidae	Θ (Llvi)	— Θ (Ashg)	
Orthotetellidae	P (Sakm)	— P (Dzhu)	(66,128)
Orthotetidae	D (Sieg)	— P (Guad)	(31,208)
Overtoniidae	D (Fame)	— P (Leon)	(31,208)
Plectambonitidae	Θ (Aren)	— Θ (m)	

Poikilosakidae	C (Step)	— P (Guad)	(208)
Productellidae	D (Emsi)	— C (Namu)	
Productidae	C (Vise)	— C (Mosc)	
Richthofeniidae	P (Sakm)	— P (Dzhu)	(208)
Scacchinellidae	C (Step)	— P (Leon)	(208)
Scapharinidae	P (Guad)		(66)
Schuchertellidae	S (l)	— P (Dzhu)	(31,128)
Sinuatellidae	C (Vise)	— C (Namu)	
Sowerbyellidae	Θ (Llde)	— D (Eife)	
Spiridiophoridae	C (Step)	— P (Leon)	
Strophalosiidae	D (Emsi)	— P (Guad)	(208)
Stropheodontidae	Θ (Ashg)	— D (Fras)	(31)
Strophomenidae	Θ (Llvi)	— D (Sieg)	(31)
Taffiidae	Θ (Trem)	— Θ (Llvi)	(194)
Teguliferinidae	C (Penn)	— P (Guad)	(208)
Thecospiridae	Tr	— Tr (Rhae)	
Tschernyschewiidae	P (Asse)	— P (Dzhu)	(208,368)
 Or. Pentamerida			
Alimbellidae	Θ (l)		
Brevicameridae	Θ (l)		
Camerellidae	Θ (Aren)	— S (l)	(31,282)
Clarkellidae	Θ (Trem)	— Θ (Ashg)	(194)
Enantiosphenidae	D (Give)		(31)
Eostrophiidae	€ (mMid)		(232,314)
Gypidulidae	S (l)	— D (Fras)	(31)
Huenellidae	€ (Dres)	— Θ (l)	(172,350)
Lycophoriidae	Θ (l)	— Θ (m)	
Parallelelasmatidae	Θ (m)	— Θ (Cara)	(31)
Parastrophinidae	Θ (Cara)	— D (Sieg)	(31,380)
Pentameridae	S (l)	— D (Fras)	
Porambonitidae	Θ (Aren)	— S (l)	(282)
Stricklandiidae	S (l)	— S (m)	
Syntrophiidae	Θ (Aren)	— Θ (Ashg)	(31,380)
Syntrophopsidae	Θ (Trem)	— Θ (Llvi)	(194,282)
Tetralobulidae	Θ (Trem)	— Θ (Aren)	(194,282)
Virginidae	Θ (Ashg)	— S (l)	
 Or. Rhynchonellida			
Allorhynchidae	C (Tour)	— P (Guad)	(66)
Amphipellidae	P (Leon)		(66)
Ancistrorhynchidae	Θ (Llvi)	— Θ (Llde)	
Atriboniidae	D (Sieg)	— P (Dzhu)	
Austrirhynchiidae	Tr (Carn)	— Tr (Rhae)	
Basiliolidae	K (Coni)	— R	
Camarotoechiidae	Θ (Llde)	— P (Guad)	(208,282)
Camerophorinidae	D (Emsi)	— D (Give)	(208)
Cardiarinidae	C (Namu)	— C (Step)	

Cryptoporidae	T (Eo-l)	— R	
Dimerellidae	Tr (Anis)	— K (Haut)	
Eatoniidae	S (l)	— D (Emsi)	
Erymnariidae	T (Eo-l)	— T (Eo-u)	
Frieliidae	T (Plio)	— R	
Hemithyrididae	T (Eo-l)	— R	
Oligorhynchiidae	Θ (Llde)	— S (l)	(31,282)
Petasmatheridae	P (Leon)	— P (Guad)	(66)
Pontisiidae	P (Asse)	— P (Guad)	(66)
Pugnacidae	D (Gedi)	— P (Dzhu)	(368)
Rhynchonellidae	Tr (Olen)	— K (Ceno)	(240)
Rhynchoporidae	C (Tour)	— P (Guad)	
Rhynchotetradidae	C (Tour)	— P (Sakm)	(208)
Rhynchotrematidae	Θ (Llde)	— D (Give)	
Septirhynchiidae	J (Call)	— J (Kimm)	
Stenoscismatidae	D (Eife)	— P (Dzhu)	(128,208)
Tetracameridae	C (Tour)	— C (Namu)	
Trigonirhynchiidae	Θ (Llde)	— P (Sakm)	
Uncinulidae	S (m)	— P (Dzhu)	(208)
Wellerellidae	C (Tour)	— K (Maes)	
Yunnanellidae	D (Give)	— D (Fame)	(31)
 Or. Spiriferida			
Ambocoeliidae	S (u)	— Tr (Indu)	(128)
Anomaloriidae	P (Guad)		(66)
Athyrididae	S (m)	— Tr (Carn)	
Athyrisinidae	D (Emsi)	— Tr (Nori)	
Brachythryrididae	D (Fame)	— P (Leon)	(31,208)
Costspiriferidae	D (Sieg)	— D (Fras)	
Crenspiriferidae	C (Penn)	— P (Guad)	(66)
Cyrtiidae	S (l)	— D (Eife)	
Cyrtinidae	S (m)	— Tr (Carn)	
Cyrtospiriferidae	D (Eife)	— C (Mosc)	(31)
Delthyrididae	S (l)	— C (Tour)	
Elythidae	D (Emsi)	— P (Dzhu)	(31,208)
Fimbrispiriferidae	D (Emsi)	— D (Give)	
Koninckinidae	Tr (Rhae)	— J (Sine)	
Martiniidae	S (u)	— P (Dzhu)	(128)
Meristellidae	Θ (Cara)	— C (Tour)	
Microspiriferidae	D (Emsi)	— C (Vise)	
Nucleospiridae	S (m)	— P (Leon)	
Paraspiriferinidae	P (Asse)	— P (Guad)	(66)
Reticulariidae	S (m)	— P (Dzhu)	(31,368)
Reticulariinidae	C (Miss)	— P (Guad)	(66)
Retziidae	D (Gedi)	— Tr (Carn)	(31)
Rhynchospirinidae	S (m)	— D (Eife)	(208)
Saranostegidae	P (Leon)	— P (Guad)	(66)

Spinocyrtiidae	D (Emsi)	— D (Fame)	
Spiriferidae	C (Tour)	— Tr (Indu)	(368)
Spiriferinidae	D (Fame)	— J (Bajo)	(31)
Suessiidae	J (l)		
Syringothyrididae	D (Fame)	— P (Guad)	
Xenosariidae	P (Guad)	.	(66)
Xestotrematidae	P (Guad)		(66)
 Or. Atrypida			
Anoplothecidae	S (m)	— D (Eife)	
Atrypidae	Θ (Ashg)	— D (Fras)	(67)
Dayiidae	Θ (Cara)	— S (u)	(31)
Kayseriidae	D (Eife)		(31)
Leptocoeliidae	S (l)	— D (m)	
Lissatrypidae	Θ (Cara)	— D (Give)	(31,67)
Palaferellidae	S (u)	— D (Give)	(67)
Zygospiridae	Θ (Cara)	— S (m)	(67)
 Or. Terebratulida			
Cancellothyrididae	J (Plie)	— R	
Centronelliidae	D (Gedi)	— P (Dzhu)	(31)
Cheniothyrididae	J (m)		
Cranaenidae	D (Emsi)	— P (Sakm)	(31)
Cryptonellidae	D (Emsi)	— P (Guad)	(31,208)
Dallinidae	Tr (Rhae)	— R	
Dictyothyrididae	J (Bath)	— J (Oxfo)	
Dielasmatidae	C (Tour)	— Tr (Rhae)	
Dyscoliidae	K (Ceno)	— R	
Eudesiidae	J (Bath)		
Hercothyrididae	T (Eo)	— R	(65)
Heterelasminidae	D (u)	— P (Guad)	(208)
Kraussinidae	T (Mi-m)	— R	
Labaiidae	C (Mosc)	— Tr (Indu)	(208)
Laqueidae	T (Mi)	— R	
Megathyrididae	K (Ceno)	— R	
Mutationellidae	S (u)	— P (Guad)	(31,208)
Notothyrididae	C (Namu)	— P (Dzhu)	(208)
Orthotomidae	J (Plie)		
Platidiidae	T (Eo-l)	— R	
Pygopidae	J (Bajo)	— K (Neoc)	
Rhenorensselaeidae	D (Gedi)	— D (Emsi)	(31)
Rhipidothyrididae	D (Gedi)	— C (Vise)	(31)
Stringocephalidae	D (Eife)	— D (Give)	(31)
Tegulithyrididae	J (Call)		
Terebratellidae	K (Albi)	— R	
Terebratulidae	Tr (Ladi)	— R	
Zeilleriidae	Tr (Olen)	— K (Albi)	(240)

Or. Thecideida		
Thecideidae	J (l)	— R
Thecidellinidae	Tr (Rhae)	— R

CHAETOGNATHA (One extant family)		
C (Mosc)	— R	(55,302)

## ECHINODERMATA

## Cl. Edrioblastoidea

Or. Pentacystida		
Astrocystitidae	Θ (Cara)	(208)

## Cl. Crinoidea

[Classification and stratigraphic ranges from the *Treatise*, Pt. T, and Moore and Strimple (1973), with emendations as noted. Approximately 13 extant families without known fossil representatives are not listed.]

?Or. Echmatocrinida		
Echmatocrinidae	Є (mMid)	(331)

Or. Diplobathrida		
Anthemocrinidae	S (m)	— S (u)
Anthracocrinidae	Θ (Lvvi)	
Archaeocrinidae	Θ (Lvvi)	— Θ (Ashg) (164)
Cleioocrinidae	Θ (Ldde)	(132)
Dimerocrinidae	Θ (Ashg)	— D (Fras)
Elpidocrinidae	S (m)	— S (u) (107)
Emperocrinidae	S (m)	— S (u) (107)
Gazacrinidae	S (m)	
Hirneocrinidae	S (l)	— S (u) (106)
Lampterocrinidae	S (m)	
Nyctocrinidae	S (m)	
Opsiocrinidae	D (l)	— D (Give) (107)
Orthocrinidae	D (Emsi)	— D (m)
Prokopicrinidae	S (m)	— S (u) (106)
Reteocrinidae	Θ (Ldde)	— Θ (Ashg)
Rhodocrinidae	Θ (Ldde)	— C (Vise)
Spyridocrinidae	D (Gedi)	(132)

Or. Monobathrida		
Abacocrinidae	S (u)	
Acocrinidae	C (l)	— C (Step)
Actinocrinidae	C (Tour)	— P (Guad)
Amphorocrinidae	C (Tour)	— C (Namu)
Batocrinidae	C (Tour)	— C (Vise)
Carpocrinidae	S (m)	— D (m)
Clonocrinidae	S (u)	— D (Sieg)

Coelocrinidae	D (m)	— C (l)	
Dichocrinidae	C (l)	— P (Guad)	
Dolatocrinidae	D (Emsi)	— D (Give)	
Eucalyptocrinitidae	S (m)	— D (m)	
Eutelocrinidae	P (Guad)		(106)
Glyptocrinidae	Θ (Cara)	— S (u)	
Hapalocrinidae	S (m)	— D (Give)	
Hexacrinitidae	S (u)	— D (Fame)	
Marsupiocrinidae	S (m)	— D (l)	
Melocrinidae	Θ (Ashg)	— D (Fame)	
Paragaricocrinidae	C (Namu)	— P (Guad)	
Parahexocrinidae	D (l)		
Paramelocrinidae	S (u)		
Patelliocrinidae	Θ (Ashg)	— D (Emsi)	
Periechocrinidae	S (m)	— C (Vise)	
Platycrinidae	D (l)	— P (u)	
Polypeltidae	S (u)	— D (Emsi)	
Scyphocrinidae	S (u)	— D (Gedi)	
Stelidiocrinidae	S (u)		
Tanaocrinidae	Θ (Aren)	— Θ (Ashg)	(238)
Tunisiacrinidae	P (Guad)		
Xenocrinidae	Θ (Ashg)		
 Or. Taxocrinida			
Synerocrinidae	C (l)	— C (Step)	
Taxocrinidae	Θ (Cara)	— C (Namu)	
 Or. Sagenocrinida			
Calycocrinidae	D (Eife)	— P (Guad)	
Dactylocrinidae	S (u)	— P (Guad)	
?Edriocrinidae	D (Gedi)	— D (m)	
Euryocrinidae	D (Give)	— P (Guad)	
Gaulocrinidae	C (l)		
Homalocrinidae	S (u)		
Icthyocrinidae	S (l)	— C (l)	
Lecanocrinidae	S (u)	— D (Eife)	
Mespiocrinidae	C (l)	— P (Guad)	
Nipterocrinidae	S (m)	— C (Tour)	(199a)
Palaeoholopodidae	P (Guad)		
Prophyllocrinidae	P (Guad)		
Sagenocrinidae	S (u)	— P (l)	
 Or. Disparida			
Acolocrinidae	S (m)		(34)
Allageocrinidae	D (u)	— P (Guad)	
Anamesocrinidae	D (m)		
Anomalocrinidae	Θ (Cara)	— Θ (Ashg)	
Apodasmocrinidae	Θ (Cara)		(108,367)

Belemnocrinidae	C (l)		
Calceocrinidae	Θ (Llde)	— P (Leon)	
Catillocrinidae	Θ (Llvi)	— P (Guad)	(165a)
Cincinnaticrinidae	Θ (Cara)	— Θ (Ashg)	(108,367)
Eustenocrinidae	Θ (Aren)	— Θ (Cara)	
Haplocrinitidae	D (m)		
Heterocrinidae	Θ (Llvi)	— S (m)	(367)
Holynocrinidae	D (m)		
Homocrinidae	Θ (Cara)	— S (m)	
Iocrinidae	Θ (Llvi)	— Θ (Ashg)	
Myelodactylidae	S (l)	— D (Gedi)	
Paradoxocrinidae	P (Guad)		
Perissocrinidae	D (m)		
Perittocrinidae	Θ (Llvi)		
Pisocrinidae	S (m)	— D (u)	
Pygmaeocrinidae	S (u)	— D (m)	
Synbathocrinidae	S (m)	— P (Guad)	
Tunguskocrinidae	Θ (m)		
Zophocrinidae	S (u)	— D (m)	
Or. Hyboocrinida			
Baerocrinidae	Θ (Llvi)	— Θ (m)	
Cornucrinidae	Θ (Llvi)	— S (l)	(330)
Hyboocrinidae	Θ (Aren)	— Θ (Cara)	(183a)
Hybocystitidae	Θ (m)		
Or. Cladida			
Adinocrinidae	C (l)		
Aethocrinidae	Θ (Trem)	— Θ (Aren)	
Agassizocrinidae	C (Namu)	— P (l)	
Ampelocrinidae	C (l)	— P (Guad)	
Anobasicrinidae	C (Namu)	— P (u)	
Aphelecrinidae	C (Tour)	— C (Namu)	
Apographocrinidae	C (Bash)	— P (Guad)	
Arkaocrinidae	C (Bash)		
Barycrinidae	D (Emsi)	— C (Vise)	
Blothrocrinidae	C (Tour)	— P (Sakm)	
Botryocrinidae	Θ (m)	— D (u)	
Bursocrinidae	C (l)		
Cadocrinidae	P (Guad)		
Calceolispongiidae	P (Sakm)	— P (Guad)	
Carabocrinidae	Θ (Llde)	— Θ (Ashg)	
Catacrinidae	C (Bash)	— P (Guad)	
Cércidocrinidae	C (l)		
Clathrocrinidae	C (Step)		
Codiocrinidae	S (m)	— P (Guad)	
Corythocrinidae	C (l)		
Cricocrinidae	C (Mosc)		

Cromyocrinidae	C (Vise)	— P (Sakm)
Crotalocrinidae	S (m)	— D (Give)
Cupressocrinitidae	D (l)	— D (u)
Cupulocrinidae	Θ (Llde)	— Θ (Ashg)
Cyathocrinidae	Θ (Llvi)	— P (Guad)
Cymbiocrinidae	C (Namu)	— P (u)
Decadocrinidae	D (Give)	— C (Step)
Dendrocrinidae	Θ (Cara)	— D (u)
Diphiocrinidae	C (Bash)	— P (Sakm)
Enocrinidae	Tr (m)	
Erisocrinidae	C (Namu)	— P (Sakm)
Eupachycrinidae	C (Namu)	
Euspiocrinidae	Θ (Cara)	— C (Tour)
Exocrinidae	C (Bash)	— P (Sakm)
Galateocrinidae	C (Mosc)	— C (Step)
Gasterocrinidae	D (l)	— D (Give)
Graphioocrinidae	C (Tour)	— P (Guad)
Hydreionocrinidae	C (Vise)	
Indocrinidae	P (Leon)	— P (Guad)
Laudonocrinidae	C (Miss)	— P (l)
Lecythocrinidae	D (Give)	— C (l)
Lophocrinidae	C (Vise)	
Manicrinidae	S (m)	— S (u)
Mastigocrinidae	Θ (m)	— C (l)
Merocrinidae	Θ (m)	— Θ (Ashg)
Metabolocrinidae	Θ (m)	— D (l)
Molloocrinidae	C (Bash)	— P (Guad)
Ontariocrinidae	Θ (m)	
Ottawocrinidae	Θ (Cara)	
Pachyocrinidae	C (l)	— P (l)
?Paractocrinidae	Θ (l)	
Paradelocrinidae	C (Bash)	— P (Guad)
Pelecocrinidae	C (l)	— P (Guad)
Petalocrinidae	S (m)	
Phanocrinidae	C (Vise)	— C (Namu)
Pirasocrinidae	C (Bash)	— P (Guad)
Porocrinidae	Θ (Cara)	— Θ (Ashg)
Poteriocrinidae	D (Sieg)	— C (Namu)
Protothylacocrinidae	D (m)	
Protencrinidae	C (Mosc)	— P (Leon)
Rhenocrinidae	D (l)	— C (Mosc)
Scotiacrinidae	C (Vise)	
Seytalocrinidae	D (Give)	— P (Guad)
Sellardsicrinidae	C (Mosc)	
Spaniocrinidae	C (l)	— P (Guad)
Sphaerocrinidae	Θ (Llvi)	— D (Give)
Stachyocrinidae	P (Guad)	

Staphylocrinidae	C (Vise)	— P (u)
Stellarocrinidae	C (Namu)	— P (Sakm)
Streblocrinidae	D (m)	— P (Guad)
Sundocrinidae	P (l)	— P (Guad)
Sycocrinidae	C (Tour)	— P (Guad)
Texocrinidae	C (Bash)	— P (Sakm)
Thenarocrinidae	S (m)	
Timorechinidae	P (Guad)	
Trimerocrinidae	P (l)	— P (Guad)
Uocrinidae	C (Vise)	— P (Guad)
Zeocrinidae	C (Tour)	— P (Guad)
 Or. Millericrinida		
Apiocrinitidae	J (l)	— K (l)
Cyclocrinidae	J (l)	— K (l)
Dadocrinidae	Tr (Anis)	
Hyocrinidae	T (Dani)	— R
Millericrinidae	J (l)	— J (Tith)
 Or. Cyrtocrinida		
Eudesicrinidae	J (Plie)	— J (Toar)
Eugeniacrinidae	J (Bath)	— K (Camp)
Hemibrachiocrinidae	K (Vala)	— K (Barr)
Hemicrinidae	J (Tith)	— K (Albi)
Holopodidae	J (Tith)	— R
Phyllocrinidae	J (Bajo)	— K (Neoc)
Plicatocrinidae	J (Hett)	— J (Tith)
Sclerocrinidae	J (Oxfo)	— K (Haut)
 Or. Bourgueticrinida		
Bathycrinidae	K (Sant)	— R
Bourgueticrinidae	K (Turo)	— T (Eo)
 Or. Isocrinida		
Holocrinidae	Tr (Anis)	
Isocrinidae	Tr (Indu)	— R
Pentacrinitidae	J (Hett)	— J (Oxfo)
 Or. Comatulida		
Antedonidae	K (Albi)	— R
Asterometridae	T (Plio)	— R
Atelecrinidae	K (Camp)	— R
Comasteridae	T (Eo)	— R
Conometridae	K (Ceno)	— T (Mi)
Decameridae	K (Vala)	— K (Albi)
Himerometridae	T (Eo)	— R
Notocrinidae	J (Bath)	— R
Paracomatulidae	J (Toar)	— K (Haut)
Pterocomidae	J (Tith)	— T (Dani)

Solanocrinitidae	J (Plie)	— K (Coni)
Thalassometridae	T (Mi-l)	— R
Thiolliericrinidae	J (Bath)	— K (Haut)
Or. Uintocrinida		
Marsupitidae	K (Sant)	
Uintocrinidae	K (Sant)	
Or. Roveocrinida		
Roveocrinidae	Tr (Ladi)	— K (Camp)
Saccocomidae	J (Kimm)	— K (Camp)

## Cl. Edrioasteroidea

[Classification and stratigraphic ranges from Bell (1976) with emendations as indicated.]

Or. Isorophida			
Agelacrinitidae	D (Gedi)	— C (Step)	
Carneyellidae	Θ (Llde)	— Θ (Ashg)	
Cyathocystidae	Θ (Llvi)	— Θ (Ashg)	
Isorophidae	Θ (Cara)	— S (m)	
Lebetodiscidae	Θ (Llde)	— S (u)	
Pyrgocystidae	Θ (Llvi)	— D (Eife)	
Or. Edrioasterida			
Edrioasteridae	Θ (Cara)		
Totiglobidae	Є (mMid)	— Є (u)	(11)
Or. Stromatocystitida			
Campstostromatidae	Є (Boto)		(208,331)
Stromatocystitidae	Є (Atda)	— Є (mMid)	(11)
Walcottidiscus	Є (mMid)		
Incertae sedis			
?Tribrachidium	V (u)		(122,208)

## Cl. Helicoplacoidea

Helicoplacidae      Ε (Atda) — Ε (Boto) (208,331)

## Cl. Blastoidea

[Classification based on the *Treatise*, Pt. S and Breimer and Macurda (1972); stratigraphic ranges from *The Fossil Record*.]

Incertae sedis	Θ (Cara)	(39a)
Or. Fissiculata		
Codasteridae	C (Vise)	— P (Guad) (33a)
Neoschismatidae	C (Tour)	— P (Guad) (33a)
Nymphaeoblastidae	D (Emsi)	— P (Sakm) (33a)
Orophocrinidae	D (Emsi)	— P (Guad) (33a)
Phaenoschismatidae	S (m)	— C (Namu) (33a)

## Or. Spiraculata

Diploblastidae	C (Vise)	— C (Namu)
Granatocrinidae	C (Tour)	— C (Vise)
Nucleocrinidae	D (Emsi)	— D (Give)
Orbitremitidae	C (Vise)	— P (Guad)
Penetremitidae	D (Eife)	— P (Guad)
Schizoblastidae	D (Give)	— P (Guad)
Troosticrinidae	S (m)	— C (Vise)

## Cl. Parablastoidea

Blastocystidae	Θ (Aren)	— Θ (Llvi)	(132,330,331)
Meristoschismatidae	Θ (Llvi)	— Θ (Llde)	(238,330)

## "Cystoidea"

[Classification based on the *Treatise*, Pt. S; stratigraphic ranges from *The Fossil Record* with emendations as indicated.]

## Cl. Rhombifera

Callocystitidae	Θ (Ashg)	— D (Fras)
Caryocrinidae	Θ (Cara)	— D (Gedi)
Caryocystitidae	Θ (Aren)	— Θ (Ashg)
Cheirocrinidae	Θ (Trem)	— Θ (Ashg) (237)
Cystoblastidae	Θ (Llde)	— Θ (Cara) (237)
Echinoencrinitidae	Θ (Aren)	— Θ (Ashg)
Echinospaeritidae	Θ (Aren)	— Θ (Ashg)
Glyptocystitidae	Θ (Llde)	— Θ (Cara) (238)
Hemicosmitidae	Θ (Aren)	— Θ (Ashg)
Heterocystitidae	S (m)	
Macrocytellidae	Θ (Trem)	— Θ (Llde) (237)
Pleurocystitidae	Θ (Aren)	— D (Eife)
Polycosmitidae	Θ (Llvi)	
Rhombiferidae	Θ (Cara)	— Θ (Ashg)
Stichocystidae	Θ (Llvi)	— Θ (Llde) (238)
Thomacystidae	Θ (Ashg)	(238)

## Cl. Diploporeta

Aristocystitidae	Θ (Trem)	— S (m)
Asteroblastidae	Θ (Aren)	— Θ (Llvi)
Dactylocystitidae	Θ (Cara)	— S (m)
Eumorphocystitidae	Θ (Llde)	(238)
Glyptosphaeritidae	Θ (Trem)	— Θ (Cara)
Gomphocystitidae	Θ (Llvi)	— S (m)
Mesocystidae	Θ (Aren)	— Θ (Llvi)
Protocrinitidae	Θ (Llvi)	— Θ (Cara)
Sphaeronitidae	Θ (Trem)	— D (Eife)

## Cl. Eocrinioidea

[Classification and stratigraphic ranges from Sprinkle (1973), with emendations as indicated.]

Or. Imbricata (= Lepidocystoidea)			
Lepidocystidae	Є (Boto)		
Or. "number 1"			
Eocrinidae	Є (Atda)	— Є (uMid)	(83,331)
Lichenoididae	Є (uMid)		
Rhopalocystidae	Θ (Trem)		
Or. "number 2"			
<i>Cambrocrinus</i>	Є (Dres)		
<i>Eocystites</i>	Є (Atda)	— Є (mMid)	(171)
<i>Eustypocystis</i>	Є (uMid)		
<i>Nolichuckia</i>	Є (uMid)		
<i>Pareocrinus</i>	Є (uMid)		
Ascocystitidae	Θ (Llvi)		(238)
Macrocystellidae	Θ (Trem)	— Θ (Llde)	(132)
Or. Coronata			
Stephanocrinidae	Θ (Llde)	— S (u)	
Incertae sedis			
<i>Cymbionites</i>	Є (lMid)		(314,316b)
<i>Peridionites</i>	Є (lMid)		(314,316b)
Cryptocrinidae	Θ (Llvi)	— S (m)	(238)
Lingulocystidae	Θ (Trem)	— Θ (Llvi)	
?Palaeocystitidae	Θ (Llde)		(39b,238)
Rhipidocystidae	Θ (Aren)	— Θ (Cara)	
Trachelocrinidae	Є (mMid)	— Є (Fran)	

## Cl. Paracrinidea

[Classification from Parsely and Mintz (1975); sources of stratigraphic ranges as indicated.]

Or. Comarocystitida			
Amygdalocystitidae	Θ (Llvi)	— Θ (Cara)	(238)
Camarocystitidae	Θ (Llvi)	— S (l)	(109,238)
Or. Platycystitida			
Malocystitidae	Θ (Llvi)	— Θ (Cara)	(238)
Platycystitidae	Θ (Llvi)	— Θ (Llde)	(108)
Incertae sedis			
Springerocystitidae	Θ (l)	— Θ (Llde)	(330)
Cl. Ctenocystoidea	Є (mMid)		(277)
Cl. Homostelea			
Or. Cincta			
Gyrocystidae	Є (mMid)		(208,331)
Trochocystitidae	Є (mMid)	— Є (uMid)	(208,331)
Cl. Homoiostelea			

[Classification and stratigraphic ranges based on the *Treatise*, Pt. S.]

Or. Soluta			
Dendrocystitidae	Є (Boto)	— D (Gedi)	(331)
Girvanicystidae	Θ (Ashg)		
Iowacystidae	Θ (Lde)	— Θ (Ashg)	(175,238)
Minervaecystidae	Є (Trep)	— Θ (Trem)	(331)
Rutroclypeidae	D (Gedi)		
Syringocrinidae	Θ (Cara)		(238)

Cl. Stylophora

[Classification and stratigraphic ranges mostly from Derstler (1979).]

Or. Cornuta			
Amygdalothecidae	Θ (Aren)	— Θ (Livi)	
Ceratocystidae	Є (mMid)	— Є (Fran)	(331)
Cothurnocystidae	Є (Trep)	— Θ (Ashg)	(208)
Phyllocystidae	Є (uMid)	— Θ (Livi)	
incertae sedis	Є (Boto)		(331)
Or. Mitrata			
Anomalocystitidae	Θ (Cara)	— D (Give)	
Kirkocystidae	Θ (Aren)	— Θ (Ashg)	
Lagynocystidae	Θ (Cara)		
Paranacystidae	Θ (Cara)	— D (Sieg)	

Cl. Echinoidea

[Classification and stratigraphic ranges based on the *Treatise*, Pt. U, and *The Fossil Record*, with emendations as noted. Five extant families without known fossil representatives are not listed.]

Or. Bothriocidaroida			
Bothriocidaridae	Θ (Lde)	— Θ (Ashg)	(174)
Or. Echinocystitoida			
Echinocystitidae	S	— P (Asse)	(181)
Lepidesthidae	D	— P (Asse)	(181)
Lepidocentridae	Θ (Ashg)	— C (Miss)	
Or. Palaechinoidea (= Melonechinoida)			
Cravenechinidae	S (m)	— P (Asse)	
Palaechinidae	C (Tour)	— C (Namu)	(255)
Or. Cidaroida			
Archaeocidaridae	D (Eife)	— P (Guad)	
Cidaridae	Tr (Carn)	— R	(316a)
Miocidaridae	C (l)	— J (Oxfo)	(316a)
Psychocidaridae	J (u)	— R	
Or. Echinothurioida			
Echinothuriidae	K (Sant)	— R	(132)

Pelanechinidae	J (Oxfo)	(132)
Or. Diadematoidea		
Aspidodiadematidae	J (Bath)	— R
Diadematidae	Tr (Nori)	— R
Or. Pedinoida		
Pedinidae	Tr (Nori)	— R
Or. Pygasteroida		
Pygasteridae	J (Sine)	— K (Ceno) (167)
Or. Salenioida		
Acrosaleniidae	J (Hett)	— K (Maes)
Saleniidae	J (Kimm)	— R
Or. Hemicidaroida		
Hemicidaridae	J (Hett)	— K (Maes)
Pseudodiadematidae	Tr (Nori)	— K (Ceno)
Or. Phymosomatoidea		
Phymosomatidae	J (Hett)	— R
Stomechinidae	J (Plie)	— R
Or. Arbacioida		
Arbaciidae	J (Bajo)	— R (167)
Or. Temnopleuroidea		
Glyphocyphidae	J (Plie)	— T (Eo-m)
Temnopleuridae	K (Ceno)	— R
Toxopneustidae	K (Camp)	— R
Or. Echinoida		
Echinidae	T (Eo-m)	— R (396)
Echinometridae	T (Pale)	— R
Parasaleniidae	T (Eo-u)	— R
Strongylocentrotidae	T (Ol-u)	— R
incertae sedis	K (Neoc)	(167)
Or. Plesiocidaroida		
Tiarechinidae	Tr (Carn)	
Or. Orthopsida		
Orthopsidae	J (Hett)	— K (Maes) (167)
Or. Holocryptopoda		
Anorthopygidae	K (Albi)	— K (Ceno)
Conoclypidae	T (Eo-m)	— T (Mi)
Conulidae	J (Call)	— T (Eo)
Discoididae	K (Apti)	— K (Maes)
Echinoneidae	K (u)	— R
Galeritidae	K (Sant)	— K (Maes)
Holocryptidae	J (Plie)	— K (Maes)

Oligopygidae	K (Seno)	— T (Ol)
Or. Clypeasteroida		
Abertellidae	T (Mi)	
Arachnoididae	T (Ol-u)	— R
Astriclypeidae	T (Ol-l)	— R
Clypeasteridae	T (Eo-u)	— R
Dendrasteridae	T (Plio)	— R
Echinarachniidae	T (Ol)	— R
Eoscutellidae	T (Eo)	
Fibulariidae	T (Pale)	— R
Laganidae	T (Eo-u)	— R
Mellitidae	T (Mi-l)	— R
Monophorasteridae	T (Eo-m)	(396)
Neolaganidae	T (Eo-u)	— T (Ol-u)
Protoscutellidae	T (Eo-m)	— T (Eo-u)
Rotulidae	T (Mi)	— R
Scutasteridae	T (Mi-l)	
Scutellidae	T (Ol)	— T (Mi)
Or. Cassiduloida		
Apatopygidae	T (Plio)	— R
Archaciidae	K (Neoc)	— K (Ceno)
Cassidulidae	K (Albi)	— R
Clypeidae	J (Bajo)	— K (Ceno)
Clypeolampadidae	K (Ceno)	— K (Maes)
Echinolampadidae	K (Ceno)	— R
Faujasiiidae	K (Ceno)	— T (Eo-u)
Galeropygidae	J (Toar)	— J (Kimm)
Nucleolitidae	J (Bajo)	— K (Maes)
Pliolampadidae	K (Camp)	(255) — R
Or. Holasteroida		
Collyritidae	J (Sine)	— K (Vala)
Disasteridae	J (Plie)	— K (Vala)
Holasteridae	K (Vala)	— R
Somaliasteridae	K (Seno)	— T (Pale)
Stenonasteridae	K (Coni)	— K (Camp)
Urechinidae	T (Mi-l)	— R
Or. Spatangoida		
Asterostomatidae	T (Pale)	— R
Brissidae	K (Ceno)	— R
Hemasteridae	K (Apti)	— R
Loveniidae	T (Eo-m)	— R
Micrasteridae	K (Ceno)	— T (Eo-m)
Palaeostomatidae	K (Ceno)	— R
Pericosmidae	T (Eo-l)	— R
Schizasteridae	K (Ceno)	— R

Spatangidae	T (Eo-m)	— R
Toxasteridae	K (Berr)	— R
Or. Neolampadoida		
Neolampadidae	T (Eo-u)	— R
Incertae sedis		
Heterocidaridae	J (Plie)	— J (Oxfo)

## Cl. Stelleroidea

[Classification from the *Treatise*, Pt. U, and stratigraphic ranges from *The Fossil Record*, with emendations as indicated. 29 extant families without known fossil representatives are not listed.]

Or. Goniactinida		
Archegonasteridae	Θ (Aren)	
Archophiactinidae	Θ (Ashg)	— D (u)
Chinianasteridae	Θ (Trem)	— Θ (Aren)
Helianthasteridae	D (Sieg)	— D (u)
Villebrunasteriidae	Θ (Trem)	
Or. Platyasterida		
Palasteriscidae	Θ (Cara)	— D (Eife)
Or. Paxillosida		
Astropectinidae	Tr (Ladi)	— R
Benthopectinidae	K (Albi)	— R
Lepidasteridae	S (m)	— D (u)
Luidiidae	T (Mi)	— R
Palasterinidae	S (l)	— D (l)
Petrasteridae	Θ (Aren)	— S
Or. Valvatida		
Eoactinidae	S (l)	
Goniasteridae	J (Plie)	— R
Hudsonasteridae	Θ (Livi)	— S (u) (181)
Mesopalaeasteridae	Θ (Ashg)	— D (Fras)
Monasteridae	C (u)	— P (l)
Neopalaeasteridae	C (Tour)	
Odontasteridae	J (Bath)	— R
Ophidiasteridae	K (Ceno)	— R
Palaeasteridae	S (m)	
Promopalaeasteridae	Θ (Cara)	— S (174)
Sphaerasteridae	J (Bath)	— R
Stauranderasteridae	J (Bath)	— T (Dani)
Xenasteridae	D (Eife)	
Or. Spinulosida		
Asterinidae	J (Bajo)	— R
Lepyriactinidae	S (l)	
Schuchertiidae	Θ (Cara)	— S (m)

Solasteridae	J (Plie)	— R
Taeniactinidae	Θ (Cara)	— C (Tour)
Tropidasteridae	J (Plie)	
Valvasteridae	J (Hett)	— R
Or. Forcipulatida		
Asteriidae	J (Plie)	— R
Brisingidae	T (Ol-l)	— R
Calliasterellidae	C (Tour)	— K (Maes)
Cnemidactinidae	Θ (Llvi)	— Θ (Ashg) (181)
Compsasteridae	D (l)	— J (Bath)
Urasterellidae	Θ (Llvi)	— C (u) (181)
Or. Stenurida		
Bdellacommidae	S (u)	— D (Eife)
Eophiuridae	Θ (Aren)	
Palaeuridae	Θ (Aren)	— D (Eife)
Phragmactinidae	Θ (Ashg)	(238)
Pradesuridae	Θ (Aren)	— D (Eife)
Rhopalocomidae	S (u)	— D (Fras)
Stenasteridae	Θ (Cara)	— Θ (Ashg)
Or. Oegophiurida		
Enocrinasteridae	Θ (Ashg)	— C (Tour)
Furcasteridae	Θ (Ashg)	— C (Tour)
Klasmuridae	D (u)	
Lapworthuridae	Θ (Aren)	— D (l)
Protasteridae	Θ (Cara)	— C (Tour)
Or. Phrymophiurida		
Eospondylidae	D (Sieg)	
Gorgonocephalidae	T (Ol)	— R
Onychasteridae	C (l)	
Or. Ophiurida		
Amphiuridae	K (Ceno)	— R
Ophiacanthidae	J (Plie)	— R
Ophiocomidae	K (Ceno)	— R
Ophiodermatidae	J (Hett)	— R
Ophiuridae	C (Tour)	— R
Ophiurinidae	S (l)	— C (Tour) (181)

## Cl. Holothuroidea

[Classification and stratigraphic ranges based on the *Treatise*, Pt. U, with augmentation from Kukalová-Peck (1973) and Pawson (1980). Names in quotes indicate sclerite taxa. Approximately 23 families of extant holothurians have been erected.]

Or Dendrochirotida “Calclamnidae”	Θ (Trem)	— R
--------------------------------------	----------	-----

"Priscopedaatidae"	C (Tour)	— R	(128a)
"Schlumbergeritidae"	J (u)		
Or. Aspidochirotida			
Holothuriidae	J (Tith)	— R	
"Stichopitidae"	C (Vise)	— R	(132)
Or. Elasipodida (extant)			
? <i>Eldonia</i>	Є (mMid)		(132)
"Paleochiridotidae"	C (Tour)	— C (u)	(128a)
"Protocaudinidae"	Θ (m)	— J (u)	
Or. Apodida			
"Achistridae"	C (Tour)	— K (Barr)	(128a,132)
Chiridotidae	K	— R	
Myriotrichidae	D (Give)	— R	(132)
Synaptidae	J (Oxfo)	— R	(132)
Or. Molpadiida			
Caudinidae	J (u)	— R	
Molpadiidae	J (u)	— R	
Or. Arthrochirotida			
Palaeocucumariidae	D (Gedi)		
Incertae sedis			
"Calclyridae"	C (u)	— P (Guad)	(132)
"Etheridgellidae"	C (Step)	— J (Oxfo)	(132)
Cl. Cyclocystoidea			
Cyclocystoididae	Є (lMid)	— D (Eife)	
Cl. Ophiocistioidea			
[Classification and stratigraphic ranges based on the <i>Treatise</i> , Pt. U, and <i>The Fossil Record</i> .]			
Eucladiidae	S (u)		
Rhenosquamidae	D (Eife)		
Rotasacciidae	D (Give)		(134)
Sollasiniidae	S (m)	— S (u)	
Volchoviidae	Θ (Aren)	— Θ (Ashg)	

## POGONOPHORA

?Or. Hyolithelminthes			
Hyolithellidae	V (N-Da)	— Є (mMid)	(12,208,327)
Torellellidae	V (N-Da)	— P (Sakm)	(326,329,371)
?Or. Sabelliditida			
Saarinidae	V (N-Da)	— Є (uTom)	(326,327)
Sabelliditidae	V (N-Da)	— Є (lMid)	(68,289,326,327)

Or. Thecanephria or Athecanephria (5 extant families)  
 \*incertae sedis                  Θ                  — T (Plio)                  (326)

### ?PHYLUM

#### Cl. Conodonta

[Assemblage classification based on the *Treatise*, Pt. W (revised), with emended stratigraphic ranges based on Fahraeus (1976) and other sources, as listed.]

#### Or. Paraconodontida

Amphigeisinidae	Є (Atda)	—	Є (mMid)	(33)
Furnishinidae	V (N-Da)	—	Θ (Trem)	(15,16,183,329)
Westergaardodinidae	Є (mMid)	—	Θ (m)	(15,16,188)

#### Or. Conodontophorida

Acanthodontidae	Θ (Trem)	—	Θ (Aren)	
Anchignathodontidae	C (Tour)	—	Tr (l)	
Bactrognathidae	D (Fame)	—	C (Tour)	(132)
Balognathidae	Θ (Aren)	—	S (l)	(339)
Belodellidae	Θ (l)	—	D (Fras)	(132)
Cavusgnathidae	D (u)	—	P (l)	
Chirognathidae	Θ (Llvi)	—	Θ (Cara)	
Clavohamulidae	Є (Trep)	—	Θ (Aren)	(204)
Cordylodontidae	Є (Trep)	—	Θ (Trem)	
Cryptotaxidae	D (Eife)	—	D (Fame)	
Cyrtodontidae	Θ (Aren)	—	Θ (Ashg)	
Distomodontidae	S (l)	—	S (m)	
Drepanoistodontidae	Θ (Trem)	—	Θ (Ashg)	
Elictognathidae	D (Fame)	—	C (Tour)	
Elloniidae	C (Bash)	—	Tr (Rhae)	
Fryxellodontidae	Θ (Trem)			
Gondolellidae	C (Mosc)	—	P (Leon)	
Hibbardellidae	Θ (Cara)	—	C (Step)	
Icriodontidae	Θ (Cara)	—	D (Fame)	(132)
Idiognathodontidae	D (Fame)	—	P (Leon)	(86,132)
Juanognathidae	Θ (l)	—	Θ (m)	
Kockenellidae	S (l)	—	S (u)	
Mestognathidae	C (Vise)	—	C (Namu)	
Multioistodontidae	Θ (Aren)	—	Θ (Ashg)	
Oepikodontidae	Θ (Aren)			
Oistodontidae	Θ (Trem)	—	Θ (Cara)	(204)
Oneotodontidae	Є (Trep)	—	Θ (Trem)	
Panderodontidae	Θ (Aren)	—	D (Give)	(91)
Paracordylodontidae	Θ (Aren)			(132)

\*May be related to unlisted extant families.

Periodontidae	Θ (Aren)	— Θ (Ashg)	
Phragmodontidae	Θ (Cara)	— Θ (Ashg)	
Polygnathidae	Θ (Ashg)	— C (Bash)	(63)
Polyplacognathidae	Θ (l)	— Θ (Cara)	
Prionodontidae	Θ (Aren)	— Θ (Livi)	
Proconodontidae	€ (Dres)	— Θ (Trem)	(183)
Protopanderodontidae	Θ (Trem)		
Pterospathodontidae	S (l)	— S (u)	
Pygodontidae	Θ (Cara)		
Rhipidognathidae	Θ (l)	— S (m)	(63)
Scolopodontidae	Θ (Trem)	— Θ (Cara)	
Strachanognathidae	Θ (Cara)	— Θ (Ashg)	
Teridontidae	€ (Fran)	— Θ (Trem)	
Ulrichodinidae	Θ (Aren)		
Xaniognathidae	P (l)	— Tr (u)	
Incertae sedis			
?Lochrieidae	C (Namu)		(201,309)
?Odontogriphidae	€ (mMid)		(53)

## HEMICORDATA

## Cl. Graptolithina

[Classification and stratigraphic ranges based on the *Treatise*, Pt. V (revised), with emendations as indicated.]

Or. Dendroidea			
Acanthograptidae	€ (Trep)	— D (Sieg)	(51,132)
Anisograptidae	Θ (Trem)	— Θ (Cara)	
Dendrograptidae	€ (uMid)	— C (Namu)	(208,265)
Ptilograptidae	Θ (Aren)	— S (u)	(51,194)
Or. Tuboidea			
Idiotubidae	€ (u)	— S (m)	
Tubidendridae	Θ (Trem)	— S (m)	
Or. Cameroidea			
Bithecocamaridae	Θ (Trem)		
Cysticamaridae	Θ (Trem)	— Θ (Aren)	(132)
Or. Stolonoidea			
Stolonodendridae	Θ (Trem)		(51,265)
Or. Crustoidea			
Hormograptidae	Θ (Llde)	— Θ (Cara)	(51)
Wimanicrustidae	Θ (Aren)		
Or. Dithecoidea			
Bulmanidendridae	€ (uMid)		(221)
Chaunograptidae	€ (uMid)	— D (Gedi)	(132,229,265)

Dithecodendridae	Є (uMid)	—	Є (u)	(229,311)
Siberiograptidae	Є (u)	—	S (l)	(229)
Or. Archaeodendrida				
Archaeodendridae	Є (uMid)			(221,265)
Or. Graptoloidea				
Abrograptidae	Θ (Aren)	—	Θ (Cara)	
Corynoididae	Θ (Cara)			
Cryptograptidae	Θ (Aren)	—	Θ (Ashg)	(51)
Cyrtograptidae	S (l)	—	S (u)	(51)
Dicaulograptidae	Θ (l)			
Dichograptidae	Θ (Trem)	—	Θ (Cara)	(51)
Dicranograptidae	Θ (Llde)	—	Θ (Ashg)	(51)
Dimorphograptidae	S (l)			
Diplograptidae	Θ (Lvvi)	—	S (l)	(51)
Glossograptidae	Θ (Aren)	—	Θ (Ashg)	(51)
Lasiograptidae	Θ (Trem)	—	Θ (Ashg)	(51,194)
Monograptidae	S (l)	—	D (Emsi)	
Nemagraptidae	Θ (Cara)			(51)
Perirograptidae	Θ (Ashg)			
Retiolitidae	Θ (Llde)	—	S (u)	(51)
Sinograptidae	Θ (Aren)	—	Θ (Lvvi)	
Incertae sedis				
Acanthastida	Θ (Trem)			
Graptoblasti	Θ (Trem)	—	Θ (Llde)	
Graptovermida	Θ (Trem)			
Cl. Pterobranchia				
Or. Rhabdopleurida				
Rhabdopleuridae	Θ (Trem)	—	R	(181,208)
?incertae sedis	Є (mMid)			(61)
Or. Cephalodiscida				
Cephalodiscidae	Θ (Trem)	—	R	(208)
Eocephalodiscidae	Θ (Trem)			(208)
Cl. Enteropneusta (4 extant families)				
?incertae sedis	Є (mMid)			(61)

## HORDATA

Cl. Ascidiacea (Urochordata) (13 extant families)			
*incertae sedis	Є (Trep)		(214)
Cl. Acrania (Cephalochordata) (1 extant family)			
*?Pikaia	Є (mMid)		(58,61)

May be related to unlisted extant families.

## Agnatha

[Classification from Nelson (1976) with stratigraphic ranges from *Osnovy Paleontologii* and *The Fossil Record*. One extant marine family without known fossil representatives is not listed; exclusively freshwater fossil families also are not listed.]

## Cl. Cephalaspidomorphi

Or. Cephalaspidiformes (=Osteostraci)	(mostly freshwater)	
Tremataspidae	Θ (Cara)	— S (u)
Sclerodontidae	S (u)	

## Or. Petromyzoniformes (= Cyclostomata)

Petromyzontidae	C (Mosc)	— R	(6)
-----------------	----------	-----	-----

## Cl. Pteraspidomorphi

## Or. Pteraspidomorphes (= Heterostraci)

?Anatolepis	€ (Fran)	— Θ (Llv)	(259)
Amphiaspidae	D (Sieg)	— D (Eife)	(130)
?Arandaspididae	Θ (Llv)		(274)
Astraspidae	Θ (Cara)	— Θ (Ashg)	
Cardipeltidae	D (Sieg)		
Cyathaspidae	S (l)	— D (Gedi)	(77,130)
Drepanaspidae	S (u)	— D (Eife)	
Eriptychiidae	Θ (Cara)	— S (u)	
Obliaspidae	D (Sieg)		
Polybranchiaspidae	D (Sieg)		
Pteraspidae	D (Gedi)	— D (Give)	

## Or. Thelodontiformes (= Coelolepida)

Apalolepididae	D (Gedi)	(356)
Loganiidae	S (m)	— S (u)
Thelodontidae	Θ (Aren)	— D (Eife)
Turiniidae	D (Gedi)	— D (Fras)

## Incertae sedis

<i>Palaeospondylus</i>	D (m)	(212)
------------------------	-------	-------

## Cl. Placodermi

[Classification and stratigraphic ranges from Denison (1978). Only families with marine representatives are listed.]

Or. Stensioellida		
Stensioellidae	D (Sieg)	

Or. Pseudopetalichthyida		
Paraplesiobatidae	D (Sieg)	

Or. Rhenanida		
Asterosteidae	D (Sieg)	— D (Fras)

Or. Ptyctodontida			
Ptyctodontidae	D (Eife)	— D (Fame)	
Or. Acanthothoraci			
Palaearcanthaspidae	D (Gedi)	— D (Sieg)	
Or. Petalichthyida			
Macropetalichthyidae	D (Sieg)	— D (Fras)	
Or. Arthrodira			
Actinolepidae	D (Sieg)		
Brachydeiridae	D (Fras)		
Buchanosteidae	D (Emsi)		
Bungartiidae	D (Fame)		
Coccosteidae	D (Eife)	— D (Fame)	
Dinichthyidae	D (Give)	— D (Fame)	
Gemuendenaspidae	D (Sieg)		
Heterosteidae	D (Give)		
Holonematidae	D (Sieg)	— D (Fame)	
Homosteidae	D (Sieg)	— D (Give)	
Leiosteidae	D (Fras)		
Leptosteidae	D (Fras)		
Mylostomatidae	D (Fras)	— D (Fame)	
Pholidosteidae	D (Fras)		
Rachiosteidae	D (Give)	— D (Fras)	
Selenosteidae	D (Fras)	— D (Fame)	(229)
Titanichthyidae	D (Fame)		
Trematosteidae	D (Fras)		
Williamsaspidae	D (Emsi)		
Or. Antiarcha			
Bothriolepidae	D (Fras)		
incertae sedis	D (Give)	— D (Fras)	
?Incertae sedis			
Cratoselachidae	C (Miss)		(280)

## I. Chondrichthyes

[Classification from Nelson (1976) and Zangerl (1981), with sources of stratigraphic ranges as indicated. Two extant marine families without fossil representatives and three fossil freshwater families are not listed.]

Or. Iniopterygia			
Iniopterygidae	C (Mosc)		(392)
Sibyrhynchidae	C (Mosc)		(392)
incertae sedis	C (Namu)		(392)
Or. Bradyodontida			
<i>Jimpohlia</i>	C (Mosc)		(392)
Acanthorhinidae	J (Plie)		(229)

Callorhynchidae	K (u)	— R	(280)
Chimaeropsidae	J (l)	— J (Tith)	(132)
Chimaeridae	C (Mosc)	— R	(392)
Cochliodontidae	D (Fame)	— P (Guad)	(132)
Copodontidae	D (Fras)	— C (Mosc)	(132)
Deltoptychiidae	C (l)	.	(392)
Menaspidae	C (l)	— P (Guad)	(212,392)
Myriacanthidae	J (Hett)	— J (Tith)	(132,229)
Rhinochimaeridae	J (u)	— R	(280)
Squaloraiidae	J (Sine)	.	(132)
Or. Euseelachii			
Ctenacanthidae	D (Give)	— C (Vise)	(132)
"Hybodontidae"	D (Fras)	— T (Pale)	(280,392)
Phoebodontidae	D (m)	— C (Mosc)	(392)
Pseudalatiidae	Tr (Rhae)	.	(257)
Ptychodontidae	K (Barr)	— T (Dani)	(132)
Tamiobatidae	D (m)	— C (u)	(392)
Tristychiidae	C (Vise)	— C (Namu)	(132)
Or. Desmiodontida			
<i>Desmodus</i>	C (l)	.	(392)
<i>Heteropetalus</i>	C (Namu)	.	(392)
Or. Cladoselachida			
Cladoselachidae	D (Fras)	.	(392)
Or. Coronodontida			
Coronodontidae	D (Fras)	.	(392)
Or. Symmoriida			
Stethacanthidae	D (u)	— C (u)	(392)
Symmoriidae	C (l)	— C (Mosc)	(392)
Or. Eugeneodontida			
<i>Campodus</i>	C (Namu)	.	(392)
Agassizodontidae	C (Namu)	— P (Guad)	(392)
Caseodontidae	C (Mosc)	— P (Guad)	(392)
Edestidae	C (Namu)	— K (u)	(280,392)
Eugeneodontidae	C (Mosc)	.	(392)
Or. Orodontida			
Orodontidae	D (u)	— C (Mosc)	(392)
Or. Petalodontida			
Petalodontidae	C (Tour)	— P (Guad)	(132,392)
Or. Heterodontiformes			
Heterodontidae	J (Tith)	— R	(132)
Palaeospinacidae	J (Sine)	— T (Pale)	(132)
Or. Hexanchiformes			

Chlamydoselachidae	T (Mi)	— R	(280)
Hexanchidae	J (Sine)	— R	(132)
Or. Lamniformes			
Carcharhinidae	K (Maes)	— R	(132)
Lamnidae	K (Apti)	— R	(132)
Odontaspidae	J (u)	— R	(280)
Orectolobidae	J (Tith)	— R	(132)
Orthacodontidae	J (Sine)	— T (Eo)	(132,280)
Scyliorhinidae	J (Tith)	— R	(132)
Sphyrnidae	T (Eo-m)	— R	(132)
Or. Squaliformes			
Pristiophoridae	K (Maes)	— R	(132)
Squalidae	J (Kimm)	— R	(229)
Squatinidae	C (Namu)	— R	(392)
Or. Rajiformes			
Dasyatidae	K (Turo)	— R	(132)
Myliobatidae	J (u)	— R	(280)
Pristidae	K (Turo)	— R	(132)
Rajidae	K (Ceno)	— R	(132)
Rhinobatidae	J (l)	— R	(280)
Torpedinidae	T (Pale)	— R	(280)
Incertae sedis			
Chondrenchelyidae	C (Vise)	— C (Namu)	(132,392)
Helodontidae	D (u)	— P (Guad)	(132,280)
Polysentoridae	C (Mosc)		(392)

## Cl. Acanthodii

[Classification and stratigraphic ranges from Denison (1979). Nonmarine families are not listed.]

Or. Climatiformes			
Climatiidae	S (m)	— D (Eife)	
Gyracanthidae	D (Sieg)	— C (Mosc)	
Or. Ischnacanthiformes			
Ischnacanthidae	S (u)	— C (Vise)	
Or. Acanthodiformes			
Acanthodidae	D (Fras)	— C (l)	

## Cl. Osteichthyes

[Classification from Nelson (1976); stratigraphic ranges based on Romer (1966), with emendations as indicated. Approximately 160 extant marine families without known fossil representatives are not listed; exclusively non-marine families also are not listed.]

Or. Dipteriformes (=Dipnoi) (only a few marine forms)

Dipnorhynchidae	D (Sieg)	— D (Fame)	(77,113,132)
Dipteridae	D (Eife)	— D (Fame)	(132)
Or. Coelacanthiformes (=Crossopterygii)			
Coelacanthidae	C (Vise)	— P (u)	(132,353a)
Diplocercidae	D (Fras)	— D (Fame)	(132,353a)
Euporosteidae	D (Give)	— D (Fras)	(132,353a)
Eusthenopteridae	D (Give)		(132,353a)
Holoptychidae	D (Give)	— D (Fame)	(132,353a)
*Laugiidae	Tr (Indu)	— J (Tith)	(242)
Osteolepididae	D (Eife)	— D (u)	(132,353a)
Porolepididae	D (Sieg)	— (Give)	(132,353a)
Or. Palaeonisciformes (only a few marine forms)			
Centrolepididae	J (Sine)		(132)
Palaeoniscidae	D (m)	— J (u)	(229)
Stegotrachelidae	D (Give)	— C (Tour)	(132)
Or. Perleidiformes (only a few marine forms)			
Cleithrolepididae	Tr (Anis)	— Tr (Carn)	(132)
Perleididae	Tr (Anis)	— Tr (Carn)	(132)
Or. Acipenseriformes (mostly freshwater)			
Acipenseridae	K (Camp)	— R	(132)
Or. Semionotiformes			
Hadrodontidae	K (u)		(354a)
Lepisosteidae	K (Camp)	— R	(132)
Semionotidae	P (Dzhu)	— K (Ceno)	(132)
Or. Pycnodontiformes			
Pycnodontidae	Tr (u)	— T (Eo-m)	
Or. Amiiformes (= Protospondyli)			
Catervariolidae	J (Tith)		(132)
Caturidae	Tr (Ladi)	— K (Ceno)	(132)
Macrosemiidae	Tr (Ladi)	— K (Ceno)	(132)
Pachycormidae	Tr (Nori)	— K (Ceno)	(132,229)
?Palaeolabridae	K (u)	— T (Pale)	(235)
Paracentrophoridae	Tr (l)		(218)
Parasemionotidae	Tr (Indu)	— Tr (u)	(132)
Promecosominidae	Tr (m)		(218)
Or. Aspidorhynchiformes			
Aspidorhynchidae	J (Bath)	— K (Maes)	(132)
Or. Pholidophoriformes			
Galkiniidae	J (Kimm)	— T (Eo-u)	(132,229,235)
Ichthyokentemidae	J (Tith)	— K (Berr)	(132)
Ligulellidae	J (Tith)		(132)

\*May be related to unlisted extant families.

Majokiidae	J (Tith)		(132)
Pleuropholidae	J (Tith)	— K (Berr)	(132)
Pholidophoridae	Tr (Ladi)	— K (Berr)	(132)
Or. Leptolepiformes			
Leptolepidae	Tr (Nori)	— K (Ceno)	(132)
Or. Ichthyodectiformes			
Allothrissopidae	J (Kimm)		(236)
Ichthyodectidae	J (Kimm)	— K (Camp)	(236)
Sauroidontidae	K (Ceno)	— K (Maes)	(236)
Or. Osteoglossiformes			
Tselfatiidae	K (Maes)		(132)
Or. Clupeiformes			
Chirocentridae	J (Tith)	— R	(132)
Clupeidae	J (Tith)	— R	(132)
Engraulidae	T (Eo-l)	— R	(229,335)
Or. Elopiformes			
Albulidae	K (Albi)	— R	(154)
?Crossognathidae	K (Neoc)	— K (Turo)	(236)
Elopidae	J (Kimm)	— R	(102,132)
Megalopidae	J (Kimm)	— R	(102)
Osmeroididae	K (Albi)	— K (Coni)	(102)
Pachyrhizodontidae	K (u)		
Thryptodontidae	K (u)		
Or. Anguilliformes			
Anguillavidae	K (u)		
Anguillidae	T (Eo-l)	— R	(335)
Congridae	K (u)	— R	
Heterenchelyidae	T (Mi)	— R	
Moringuidae	T (Eo-u)	— R	(335)
Muraenesocidae	T (Eo-l)	— R	(335)
Muraenidae	T (Eo-l)	— R	(335)
Nemichthyidae	T (Eo)	— R	
Nettastomatidae	T (Eo-u)	— R	(335)
Ophichthidae	T (Eo-m)	— R	(335)
Urenchelyidae	K (Ceno)	— K (Maes)	
Xenocongridae	T (Eo)	— R	
Or. Notacanthiformes			
Halosauridae	K (Ceno)	— R	(132)
Notacanthidae	K (Maes)	— R	(132)
Or. Salmoniformes			
Alepocephalidae	T (Eo-m)	— R	(335)
Argentinidae	T (Eo-l)	— R	(335)
Bathylagidae	T (Eo-m)	— R	

Chauliodontidae	T (Mi-u)	— R	(229)
Gonostomatidae	K (Ceno)	— R	(132)
Osmeridae	T (Eo-l)	— R	(335)
Protostomiatiidae	K (u)		
Salmonidae	K (Camp)	— R	(132)
Sternopychidae	T (Ol-l)	— R	(229)
Stomiatiidae	K (Ceno)	— R	(132)
Or. Gonorhynchiformes			
Chanidae	K (Albi)	— R	(132)
Gonorhynchidae	K (Ceno)	— R	(132)
Or. Siluriformes (only a few marine forms)			
Aridae	T (Eo-u)	— R	(335)
Bagridae	T (Eo-l)	— R	
Plotosidae	T (Ol)	— R	
Or. Myctophiformes			
Myctophidae	K (Ceno)	— R	(126)
Sardinioidae	K (Ceno)	— K (Camp)	(126)
Or. Aulopiformes			
Aulopidae	T (Eo-l)	— R	(281,335)
Cheirothricidae	K (u)		(281)
Chlorophthalmidae	T (Eo-l)	— R	(281,335)
Cimolichthyidae	K (Ceno)	— K (Camp)	(126,281)
Dercetidae	K (Ceno)	— T (Pale)	(126,281)
Enchodontidae	K (Albi)	— T (Pale)	(132,281)
Halecidae	K (Ceno)	— K (Turo)	(126,281)
Paralepididae	T (Eo)	— R	(281)
Prionepididae	K (Ceno)	— K (Turo)	(126,281)
Synodontidae	T (Eo-l)	— R	(281,335)
Tomagnatidae	K (u)		(281)
Or. Polymixiiformes			
Polymixiidae	K (Ceno)	— R	(132)
Sphenocephalidae	K (u)		(132)
Or. Gadiformes			
Bregmacerotidae	T (Eo-u)	— R	(335)
Carapidae	T (Eo-m)	— R	(50)
Gadidae	K (Camp)	— R	(132,229)
Macrouridae	T (Pale)	— R	(132)
Merlucciidae	T (Eo-l)	— R	(335)
Moridae	T (Eo-l)	— R	(335)
Ophidiidae	T (Eo-l)	— R	(335)
Or. Batrachoidiformes			
Batrachoididae	T (Mi-u)	— R	(132)
Or. Lophiiformes			

Antennariidae	T (Eo-m)	— R	(132)
Chaunacidae	T (Eo-u)	— R	(335)
Lophiidae	T (Eo-m)	— R	(132)
Or. Atheriniformes			
Atherinidae	T (Eo-l)	— R	(132,335)
Belonidae	T (Eo)	— R	
Cyprinodontidae	T (Ol-l)	— R	(132)
Exocoetidae	K (Maes)	— R	(132)
Forficidae	T (Mi)		
Scomberesocidae	T (Eo)	— R	
Or. Lampridiformes			
Lamprididae	T (Mi-u)	— R	(132)
Lophotidae	T (Eo-u)	— R	(335)
Veliferidae	T (Eo-l)	— R	(132)
Or. Ctenothrissiformes			
Aulolepididae	K (Ceno)		(132,229)
Ctenothrissidae	K (Ceno)	— K (Turo)	(132)
Or. Beryciformes			
Aipichthyidae	K (u)		
Berycidae	K (u)	— R	
Dinopterygidae	K (u)		
Diretmidae	T (Mi)	— R	
Holocentridae	K (u)	— R	
Melamphaeidae	T (Mi-m)	— R	(50)
Monocentridae	T (Pale)	— R	
Pharmacichthyidae	K (u)		
Pycnosteroididae	K (u)		
Trachichthyidae	K (Ceno)	— R	
incertae sedis	J (Tith)		(132)
Or. Zeiformes			
Caproidae	T (Eo-u)	— R	(335)
Zeidae	T (Pale)	— R	
Or. Syngnathiformes			
Aulostomidae	T (Eo-l)	— R	(132)
Centriscidae	T (Eo)	— R	(229)
Fistulariidae	T (Eo)	— R	(229)
Macrorhamphosidae	T (Ol)	— R	
Ramphosoidae	T (Eo-l)	— T (Eo-m)	
Solenostomidae	T (Eo)	— R	
Syngnathidae	T (Eo-m)	— R	
incertae sedis	K (Ceno)		
Or. Gasterosteiformes			
Aulorhynchidae	T (Eo-m)	— R	(132)
Gasterosteidae	T (Mi-u)	— R	(132)

Or. Scorpaeeniformes			
Agonidae	T (Eo-l)	— R	(335)
Anoplopomatidae	T (Mi-u)	— R	
Cottidae	T (Eo-l)	— R	
Cyclopteridae	Q (Plei)	— R	
Hexagrammidae	T (Mi-u)	— R	
Platycephalidae	T (Eo-u)	— R	(335)
Scorpaenidae	T (Eo-l)	— R	(132,335)
Synancejidae	T (Eo-l)	— R	(132,335)
Triglidae	T (Eo-l)	— R	(132,335)
Zaniolepididae	T (Eo-l)	— R	(335)
Or. Dactylopteriformes			
Dactylopteridae	T (Eo-u)	— R	(335)
Or. Perciformes			
Acanthuridae	T (Eo-m)	— R	(132)
Ammodytidae	T (Eo)	— R	
Apogonidae	T (Eo-l)	— R	(335)
Blenniidae	T (Eo-m)	— R	(132)
Blochiidae	K (Ceno)	— T (Eo-m)	(132)
Branchiostegidae	T (Eo-u)	— R	(335)
Callionymidae	T (Eo-l)	— R	(132)
Carangidae	T (Eo-l)	— R	(335)
Centropomidae	T (Eo-l)	— R	(335)
Cepolidae	T (Eo)	— R	
Chaetodontidae	K (u)	— R	(229)
Champsodontidae	T (Ol)	— R	
Channidae	K (Neoc)	— R	(229)
Chiasmodontidae	T (Ol)	— R	(229)
Clinidae	T (Eo)	— R	
Echeneidae	T (Ol)	— R	
Embiotocidae	T (Eo)	— R	
Enoplosidae	T (Eo)	— R	
Ephippidae	T (Eo)	— R	
Gempylidae	K (u)	— R	
Gerridae	T (Eo-m)	— R	(335)
Gobiidae	T (Eo-m)	— R	(132)
Istiophoridae	K (u)	— R	
Kuhliidae	T (Eo-m)	— R	(335)
Labridae	T (Pale)	— R	
Lactariidae	T (Mi)	— R	
Leiognathidae	T (Ol)	— R	
Leptoscopidae	T (Mi)	— R	
Lobotidae	T (Ol)	— R	
Lutjanidae	T (Eo-l)	— R	(335)
Menidae	T (Dani)	— R	(132)
Mugilidae	T (Ol-l)	— R	(132)

Mullidae	T (Mi)	— R	
Nemipteridae	T (Eo-l)	— R	(335)
Nameidae	T (Eo-l)	— R	(132)
Nototheniidae	T (Mi)	— R	
Oplegnathidae	T (Plio)	— R	
Owstoniidae	T (Plio)	— R	
Palaeorhynchidae	T (Eo)	— R	
Phyllodontidae	K (Camp)	— T (Eo-u)	(132,154)
Plesiopidae	T (Eo-u)	— R	(335)
Polynemidae	T (Mi-u)	— R	(132)
Pomacentridae	T (Eo)	— R	
Pomadasytidae	T (Eo-m)	— R	(335)
Pomatomidae	T (Eo-l)	— R	(335)
Princanthidae	T (Eo-l)	— R	(335)
Scaridae	T (Eo)	— R	
Scatophagidae	T (Eo)	— R	
Sciaenidae	K (Maes)	— R	
Scombridae	T (Pale)	— R	
Serranidae	T (Dani)	— R	(132)
Siganidae	T (Ol)	— R	
Sillaginidae	T (Ol)	— R	
Sparidae	T (Pale)	— R	
Sphyraenidae	T (Eo-l)	— R	(132)
Toxotidae	T (Eo)	— R	
Trachinidae	T (Eo-l)	— R	(132)
Trichiuridae	T (Pale)	— R	
Tripterygiidae	T (Plio)	— R	
Uranoscopidae	T (Eo-l)	— R	(132)
Xiphiidae	K (u)	— R	
 Or. Gobiesociformes			
Gobiesocidae	T (Mi-u)	— R	(132)
 Or. Pleuronectiformes			
Bothidae	T (Eo-l)	— R	(50,132)
Citharidae	T (Eo-l)	— R	(132)
Cynoglossidae	T (Mi)	— R	
Pleuronectidae	T (Eo-l)	— R	(50)
Psettodidae	T (Eo-m)	— R	(132)
Scophthalmidae	T (Eo-l)	— R	(132)
Soleidae	T (Pale)	— R	
 Or. Tetraodontiformes			
Balistidae	T (Eo-l)	— R	(132)
Diodontidae	T (Eo)	— R	
Eotrigonodontidae	K (Ceno)	— T (Eo-m)	(132)
Molidae	T (Mi)	— R	
Ostraciontidae	T (Eo)	— R	

Tetraodontidae	T (Eo)	— R	
Triacanthidae	K (u)	— R	
Triacanthodidae	T (Eo)	— R	
Trigonodontidae	K (Ceno)	— T (Plio)	(132)
Triodontidae	T (Eo-l)	— R	(132)

## Cl. Reptilia

[Marine families only. Classification is from Romer (1966), with stratigraphic ranges from Romer (1966) and *The Fossil Record*.]

Or. Chelonia			
Cheloniidae	K (l)	— R	(354a)
Dermochelyidae	T (Eo-l)	— R	
Protostegidae	K (u)	— T (Ol)	
Toxochelyidae	K (Albi)	— T (Eo)	
Or. Squamata			
Aigialosauridae	J (Tith)	— K (l)	
Dolichosauridae	K (l)	— K (u)	
Mosasauridae	K (u)		
Palaeophidae	T (Eo)		
Simoliophidae	K (Albi)	— K (Ceno)	
Or. Nothosauria			
Nothosauridae	P (Guad)	— Tr (Carn)	
Pachypleurosauridae	Tr (m)	— Tr (Carn)	
Simosauridae	Tr (m)	— Tr (Carn)	
Or. Plesiosauria			
Cymatosauridae	Tr (m)	— Tr (u)	
Elasmosauridae	K (Berr)	— K (Maes)	
Leptocleididae	J (Hett)	— K (Haut)	
Pistosauridae	Tr (Ladi)		
Plesiosauridae	Tr (Ladi)	— K (Apti)	
Pliosauridae	J (Bath)	— K (Maes)	
Polycotylidae	K (Apti)	— K (Maes)	
Thaumatosauridae	J (l)	— J (u)	
Or. Placodontia			
Helveticosauridae	Tr (Anis)	— Tr (Ladi)	
Henodontidae	Tr (Carn)		
Placochelyidae	Tr (Anis)	— Tr (Rhae)	
Placodontidae	Tr (Olen)	— Tr (Ladi)	
Or. Ichthyosauria			
Ichthyosauridae	J (Hett)	— K (Maes)	
Mixosauridae	Tr (Anis)	— Tr (Nori)	
Omphalosauridae	Tr (Olen)	— Tr (Ladi)	
Shastasauridae	Tr (Ladi)	— Tr (Nori)	
Stenopterygiidae	J (Plie)	— K (Apti)	

Or. Crocodilia		
Metriorhynchidae	J (Call)	— K (Haut)
Teleosauridae	J (Toar)	— K (Neoc)

## Cl. Mammalia

[Marine or largely marine families only. Classification is from Romer (1966), and stratigraphic ranges are from *The Fossil Record*, Lipps and Mitchell (1976), and Romer (1966).]

Or. Cetacea		
Acrodelphidae	T (Mi-l)	— T (Plio)
Agorophiidae	T (Ol-u)	
Balaenidae	T (Mi-l)	— R
Balaenopteridae	T (Mi-u)	— R
Basilosauridae	T (Eo-u)	— T (Ol-l)
Cetotheriidae	T (Ol-u)	— T (Plio)
Delphinidae	T (Plio)	— R
Dorudontidae	T (Eo-u)	
Eurhinodelphidae	T (Mi-l)	— T (Mi-u)
Hemisyntrachelidae	T (Mi-m)	— T (Plio)
Monodontidae	Q (Plei)	— R
Patriocetidae	T (Eo-u)	— T (Ol-u)
Phoncaenidae	T (Mi-u)	— R
Physeteridae	T (Mi-l)	— R
Platanisidae	T (Mi-m)	— R
Protocetidae	T (Eo-m)	
Squalodontidae	T (Ol-l)	— T (Mi-u)
Ziphiidae	T (Mi-l)	— R

## Or. Sirenia

Dugongidae	T (Eo-m)	— R
Manatidae	T (Mi-l)	— R

## Or. Carnivora (Pinnipedia)

Enaliarctidae	T (Mi-l)	— T (Mi-m)	(258a)
Odobenidae	T (Mi-u)	— R	
Otariidae	T (Mi-m)	— R	
Phocidae	T (Mi-m)	— R	

## BIBLIOGRAPHY

- 1 Adegoke, O.S. 1977. Stratigraphy and paleontology of the Ewekoro Formation (Paleocene) of southwestern Nigeria. *Bull. Amer. Paleontol.* 71:1-379.
- 2 Alberti, H. 1979. Devonian trilobite biostratigraphy. Pp. 313-324. In: M.R. House, C.T. Scrutton, and M.G. Bassett, eds. *The Devonian System*. Palaeontol. Assoc. Lond., Spec. Pap. Palaeontol. No. 23.
- 3 Allison, E.C. 1955. Middle Cretaceous Gastropoda from Punta China, Baja California, Mexico. *Jour. Paleontol.* 29:400-432.
- 4 Andreev, Yu.N., and M.I. Mandelstam. 1971. Biogeographical associations of Cretaceous ostracods in the USSR. Pp. 615-630. In: H.J. Oertli, ed. *Paléoécologi des Ostracodes*. Bull. Centre Rech. Pau-SNPA, v. 5 (Suppl.).
- 5 Apollonov, M.K. 1974. *Ashgill'skie trilobiti Kazakhstana*. Nauka; Moscow, 136 pp.
- 6 Bardack, D., and R. Zangerl. 1968. First fossil lamprey — a record from the Pennsylvanian of Illinois. *Science* 162:1265-1267.
- 7 Batten, R.L. 1958. Permian Gastropoda of the southwestern United States: 2. Pleurotomariacea: Portlockiellidae, Phymatopleuridae, and Eotomariidae. *Amer. Mus. Nat. Hist. Bull.* 114:153-246.
- 8 \_\_\_\_\_ 1973. The vicissitudes of the gastropods during the interval of Guadalupian-Ladinian time. Pp. 596-607. In: A. Logan and L.V. Hills, eds. *The Permian and Triassic Systems and their Mutual Boundary*. Can. Soc. Pet. Geol.; Calgary, Alberta.
- 9 Beikirch, D.W., and R.M. Feldmann. 1980. Decapod crustaceans from the Pflugerville Member, Austin Formation (Late Cretaceous: Campanian) of Texas. *Jour. Paleontol.* 54:309-324.
- 10 Bell, B.M. 1976. A study of North American Edrioasteroidea. N.Y. State Mus. Mem. 21, 446 pp.
- 11 \_\_\_\_\_, and J. Sprinkle. 1978. *Totiglobus*, an unusual new edrioasteroid from the Middle Cambrian of Nevada. *Jour. Paleontol.* 52:234-242.
- 12 Bengtson, S. 1968. The problematic genus *Mobergella* from the Lower Cambrian of the Baltic area. *Lethaia* 1:325-351.
- 13 \_\_\_\_\_, 1970. The Lower Cambrian fossil *Tommotia*. *Lethaia* 3:363-392.
- 14 \_\_\_\_\_, 1976. The structure of some Middle Cambrian conodonts, and the early evolution of conodont structure and function. *Lethaia* 9:185-206.
- 15 \_\_\_\_\_, 1977a. Aspects of problematic fossils in the early Paleozoic. *Acta Universitatis Upsaliensis*, No. 415, 71 pp.
- 16 \_\_\_\_\_, 1977b. Early Cambrian button-shaped phosphatic microfossils from the Siberian Platform. *Palaeontology* 20:751-762.
- 17 \_\_\_\_\_, 1981. *Atractosella*, a Silurian alcyonacean octocoral. *Jour. Paleontol.* 55:281-294.
- 18 \_\_\_\_\_, and V.V. Missarzhevsky. 1981. Coeloscleritophora — a major group of enigmatic Cambrian metazoans. Pp. 19-21. In: M.E. Taylor, ed. *Short Papers for the Second International Symposium on the Cambrian System*. U.S. Geol. Surv. Open-File Rpt. 81-743.

- 19 Berg-Madsen, V., and J.S. Peel. 1978. Middle Cambrian monoplacophorans from Bornholm and Australia, and the systematic position of the bellerophontiform molluscs. *Lethaia* 11:113-126.
- 20 Bergström, J. 1968. *Eolimulus*, a Lower Cambrian xiphosurid from Sweden. *Geol. Fören. Stockholm Förh.* 90:489-503.
- 21 \_\_\_\_\_, 1971. *Paleomerus* — merostome or merostomoid. *Lethaia* 4:393-401.
- 22 \_\_\_\_\_, 1975. Functional morphology and evolution of xiphosurids. *Fossils and Strata*, No. 4, pp. 291-306.
- 22a \_\_\_\_\_, 1980. Morphology and systematics of early arthropods. *Abh. Verh. naturwiss. Ver. Hamburg* 23:7-42.
- 23 \_\_\_\_\_, W. Stürmer, and G. Winter. 1980. *Palaeoisopus*, *Palaeopantopus*, and *Palaeothea*, pycnogonid arthropods from the Lower Devonian Hunsrück Slate, West Germany. *Paläontol. Z.* 54:7-54.
- 23a Bergström, S.M. 1979. First report of the enigmatic Ordovician microfossil *Konyrium* in North America. *Jour. Paleontol.* 53:320-327.
- 23b Bigey, F. 1981. Overgrowths in Palaeozoic Bryozoa: Examples from Devonian forms. Pp. 7-18. In: G.P. Larwood and C. Nielsen, eds. *Recent and Fossil Bryozoa*. Olsen and Olsen; Fredensborg, Denmark.
- 24 Bischoff, G.C.O. 1978. *Septodaeum silicum*, a representative of a new subclass Setodaearia of the Anthozoa, with partial preservation of soft parts. *Senckenbergiana leth.* 59:229-273.
- 25 \_\_\_\_\_, 1978. Internal structures of conulariid tests and their functional significance, with special reference to Circonculariina n. suborder. *Senckenbergiana leth.* 59:275-327.
- 26 Bless, M.J.M., and H. Jordan. 1971. The new genus *Copelandella* from the Carboniferous; the youngest known beyrichiacean ostracodes. *Lethaia* 4:185-190.
- 27 \_\_\_\_\_, and \_\_\_\_\_. 1971. Classification of paleocapid ostracodes belonging to the families Ctenoloculinidae, Hollinidae, and Hollinellidae. Pp. 869-890. In: H.J. Oertli, ed. *Paléoécologie des Ostracodes*. Bull. Centre Rech. Pau-SNPA, v. 5 (Suppl.).
- 28 Bloeser, B., J.W. Schopf, R.J. Horodyski, and W.J. Breed. 1977. Chitinozoans from the late Precambrian Chuar Group of the Grand Canyon, Arizona. *Science* 195:676-679.
- 29 Bogoyavlenskaya, O.V. 1974. Printsipi sistematizatsii Stromatoporoidei. Pp. 20-27. In: B.S. Sokolov, ed. *Ancient Cnidaria*, v. 1. Nauka: Moscow.
- 30 Bouček, B. 1964. The Tentaculites of Bohemia. Czech. Acad. Sci., Prague, 215 pp.
- 31 Boucot, A.J. 1975. Evolution and Extinction Rate Controls. Elsevier; Amsterdam, 427 pp.
- 32 Brazier, M.D. 1979. The Cambrian radiation event. Pp. 103-159. In: M.R. House, ed. *The Origin of Major Intertebrate Groups*. Academic Press; New York.
- 33 \_\_\_\_\_, and R.A. Hewitt. 1981. Faunal sequence within the Lower Cambrian "non-trilobite zone" (s.l.) of central England and correlated regions. Pp. 23-33. In: M.E. Taylor, ed. *Short Papers for the Second International Symposium on the Cambrian System*. U.S. Geol. Surv. Open-File Rpt. 81-743.
- 33a Breimer, A., and D.B. Macurda, Jr. 1972. The phylogeny of fissiculate blastoids. *Verh. K. Ned. Akad. Wet.* 26:1-390.

- 34 Brett, C.E. 1980. *Paracolocrinus*, a new inadunate crinoid genus from the Rochester Shale (Silurian, Wenlockian) of New York. Jour. Paleontol. 54:913-922.
- 35 Briggs, D.E.G. 1976. The arthropod *Branchiocaris* n. gen., Middle Cambrian Burgess Shale, British Columbia. Geol. Surv. Can. Bull. 264:1-29.
- 36 \_\_\_\_\_. 1977. Evolutionary significance of *Canadaspis*, the earliest positively identified crustacean. Jour. Paleontol. 51(Suppl.):4.
- 37 \_\_\_\_\_. 1977. Bivalved arthropods from the Cambrian Burgess Shale of British Columbia. Palaeontology 20:595-622.
- 38 \_\_\_\_\_. 1978. A new trilobite-like arthropod from the Lower Cambrian Kinzers Formation, Pennsylvania. Jour. Paleontol. 52:132-140.
- 39 \_\_\_\_\_. 1978. The morphology, mode of life, and affinities of *Canadaspis perfecta* (Crustacea: Phyllocarida) Middle Cambrian, Burgess Shale, British Columbia. Phil. Trans. R. Soc. Lond. ser. B. 281:439-487.
- 39a Broadhead, T.W. 1980. Range extension — Middle Ordovician blastoids (Echinodermata) from the Southern Appalachians. Geol. Soc. Amer. Abstr. Program 12:172.
- 39b \_\_\_\_\_, and F.C. Breland, Jr. 1980. Palaeocystitidae — Middle Ordovician echinoderms with eocrinoid and paracrinoid affinities. Geol. Soc. Amer. Abstr. Program 12:172.
- 40 Brooks, H.K., and K.E. Caster. 1956. *Pseudoarctolepis sharpi* n. gen., n. sp. (Phyllocarida), from the Wheeler Shale (Middle Cambrian) of Utah. Jour. Paleontol. 30:9-14.
- 41 Brunton, D.C. 1977. Appendages of *Sidneyia*, *Emeraldella*, and *Leancholia* and their bearing on trilobitoid classification. Jour. Paleontol. 51(Suppl.):4-5.
- 42 Chang, W.T. 1966. On the classification of Redlichiacea, with description of new families and new genera. Acta Paleontol. Sin. 14:135-184.
- 43 Cheetham, A.H. 1972. Cheiostome Bryozoa of late Eocene age from Eua, Tonga. U.S. Geol. Surv. Prof. Pap. 640-E, 26 pp.
- 44 Chen J.Y., Zou X.P., Chen T.E., and Qi D.L. 1979. Late Cambrian cephalopods of north China — Plectronocerida, Protactinocerida (ord. nov.) and Yanhecerida (ord. nov.). Acta Palaeontol. Sin. 18:1-24.
- 45 \_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_, and \_\_\_\_\_. 1979. Late Cambrian Ellesmerocerida (Cephalopoda) of north China. Acta Paleontol. Sin. 18:120-146.
- 46 Chlupáč, I. 1963. Report on the merostomes from the Ordovician of central Bohemia. Základní Otisk Věstnika Ústředního Ústavu Geologického 38:399-402.
- 47 Cisne, J.L. 1974. Evolution of the world fauna of aquatic free-living arthropods. Evolution 28:337-366.
- 48 Clark, D.L., and R.A. Robison. 1969. Oldest conodonts in North America. Jour. Paleontol. 43:1044-1046.
- 49 Clark, R.B. 1969. Systematics and phylogeny: Annelida, Echiura, Sipuncula. Pp. 1-68. In: M. Florkin and B.T. Scheer, eds. Chemical Zoology, v. 4. Academic Press; New York.
- 50 Clarke, M.R., and J.E. Fitch. 1980. Statoliths of Cenozoic teuthoid cephalopods from North America. Palaeontology 22:479-511.
- 51 Clarkson, E.N.K. 1979. Invertebrate Palaeontology and Evolution. George Allen &

- Unwin; London, 323 pp.
- 51a Collins, D., and D.M. Rudkin. 1981. *Priscansermarinus barnetti*, a probable lepadomorph barnacle from the Middle Cambrian Burgess Shale of British Columbia. Jour. Paleontol. 55:1006-1015.
- 52 Collins, J.S.H., and S.F. Morris. 1976. Tertiary and Pleistocene crabs from Barbados and Trinidad. Palaeontology 19:107-131.
- 53 Conway Morris, S. 1976. A new Cambrian lophophorate from the Burgess Shale of British Columbia. Palaeontology 19:199-222.
- 54 \_\_\_\_\_ 1977a. Fossil priapulid worms. Palaeontol. Assoc. Lond., Spec. Pap. Palaeontol. No. 20, 95 pp.
- 55 \_\_\_\_\_ 1977b. A redescription of the Middle Cambrian worm *Amiskwia sagittiformis* Walcott from the Burgess Shale of British Columbia. Paläontol. Z. 51:271-287.
- 56 \_\_\_\_\_ 1977c. A new entoproct-like organism from the Burgess Shale of British Columbia. Palaeontology 20:833-845.
- 57 \_\_\_\_\_ 1977d. A new metazoan from the Cambrian Burgess Shale of British Columbia. Palaeontology 20:623-640.
- 58 \_\_\_\_\_ 1977e. Aspects of the Burgess Shale fauna, with particular reference to the non-arthropod component. Jour. Paleontol. 51(Suppl.): 7-8.
- 59 \_\_\_\_\_ 1978. *Laggania cambria* Walcott: A composite fossil. Jour. Paleontol. 52:126-131.
- 60 \_\_\_\_\_ 1979. Middle Cambrian polychaetes from the Burgess Shale of British Columbia. Phil. Trans. R. Soc. Lond., ser. B, 285:227-274.
- 61 \_\_\_\_\_ 1979. The Burgess Shale (Middle Cambrian) fauna. Ann. Rev. Ecol. Syst. 10:327-349.
- 62 \_\_\_\_\_ 1981. Parasites and the fossil record. Parasitol. 82:489-509.
- 63a \_\_\_\_\_, and R.A. Robison. 1982. The enigmatic medusoid *Peytoia* and a comparison of some Cambrian biotas. Jour. Paleontol. 56:116-122.
- 63 Cooper, B.J. 1977. Toward a familial classification of Silurian conodonts. Jour. Paleontol. 51:1057-1071.
- 64 Cooper, G.A. 1976. Lower Cambrian brachiopods from the Rift Valley (Israel and Jordan). Jour. Paleontol. 50:269-289.
- 65 \_\_\_\_\_ 1979. Tertiary and Cretaceous Brachiopods from Cuba and the Caribbean. Smithson. Contrib. Paleobiology, No. 37, 45 pp.
- 66 \_\_\_\_\_, and R.E. Grant. 1972-1977. Permian brachiopods of west Texas, I-VI. Smithson. Contrib. Paleobiology, Nos. 14, 15, 19, 21, 24, 32.
- 67 Copper, P. 1973. New Siluro-Devonian atrypoid brachiopods. Jour. Paleontol. 47:484-500.
- 68 Cowie, J.W., and M.F. Glaessner. 1975. The Precambrian-Cambrian Boundary: A Symposium. Earth Sci. Rev. 11:209-251.
- 69 \_\_\_\_\_, and A. Yu. Rozanov. 1974. I.U.G.S. Precambrian/Cambrian Boundary Working Group in Siberia, 1973. Geol. Mag. 111:237-252.

- 70 \_\_\_\_\_, A.W.A. Rushton, and C.J. Stubblefield. 1972. A correlation of Cambrian rocks in the British Isles. Geol. Soc. London, Spec. Rept. 2, 42 pp.
- 71 Cox, L.R. 1965. Jurassic Bivalvia and Gastropoda from Tanganyika and Kenya. Br. Mus. (Nat. Hist.) Bull., Geol. Suppl. I, 213 pp.
- 72 Cressey, R., and C. Patterson. 1973. Fossil parasitic copepods from a Lower Cretaceous fish. Science 180:1283-1285.
- 73 Crick, R.E. 1981. Diversity and evolutionary rates of Cambro-Ordovician nautiloids. Paleobiology 7:200-215.
- 74 Daily, B. 1972. The base of the Cambrian and the first Cambrian faunas. Univ. Adelaide Centre for Precambrian Res., Spec. Pap. 1:13-42.
- 75 Davies, A.M. 1971. Tertiary Faunas: I. The Composition of Tertiary Faunas. Elsevier; Amsterdam, 571 pp.
- 76 Debrenne, F., and M. Debrenne. 1978. Archaeocyathid fauna of the lowest fossiliferous levels of Tiout (Lower Cambrian, southern Morocco). Geol. Mag. 115:101-120.
- 77 Denison, R.H. 1956. A review of the habitat of the earliest vertebrates. Fieldiana:Geol. 11:359-441.
- 78 Denison, R. 1978. Placodermi. Handbook of Paleoichthyology, v. 2. Gustav Fischer Verlag; Stuttgart, 128 pp.
- 79 \_\_\_\_\_, 1979. Acanthodii. Handbook of Paleoichthyology, v. 5. Gustav Fischer Verlag; Stuttgart, 62 pp.
- 80 Derstler, K. 1979. Biogeography of the stylophoran carpoids (Echinodermata). Pp. 91-104. In: J. Gray and A.J. Boucot, eds. Historical Biogeography, Plate Tectonics and the Changing Environment. Oregon State Univ. Press; Corvallis, Oregon.
- 81 Donovan, D.T., J.H. Callomon, and M.K. Howarth. 1981. Classification of the Jurassic Ammonitina. Pp. 101-155. In: M.R. House and J.R. Senior, eds. The Ammonoidea. Syst. Assoc. Spec. Vol. 18. Academic Press; New York.
- 82 Durham, J.W. 1974. Systematic position of *Eldonia ludwigi* Walcott. Jour. Paleontol. 48:750-755.
- 83 \_\_\_\_\_, 1978. A Lower Cambrian eocrinoid. Jour. Paleontol. 52:195-199.
- 84 Dzik, J. 1975. The origin and early phylogeny of the cheilostomatous Bryozoa. Acta Palaeontol. Pol. 20:395-420.
- 84a\_\_\_\_\_. 1981. Evolutionary relationships of the early Palaeozoic 'cyclostomatous' Bryozoa. Palaeontology 24:827-861.
- 85 Eldredge, N. 1974. Revision of the suborder Synziphosurina (Chelicerata, Merostomata), with remarks on merostome phylogeny. Amer. Mus. Nov., No. 2543, 41 pp.
- 86 Ellison, S.P., Jr. 1972. Conodont taxonomy in the Pennsylvanian. Geologica et Palaeontologica SB1:127-146.
- 87 Ergaliev, G.K. 1981. Upper Cambrian biostratigraphy of the Kyrshabakty Section, Maly Karatau, southern Kazakhstan. Pp. 82-88. In: M.E. Taylor, ed. Short Papers for the Second International Symposium on the Cambrian System. U.S. Geol. Surv. Open-File Rpt. 81-743.
- 88 Erina, M.V., and A.I. Kim. 1980. On some Ordovician Scleractinia-like corals from the

- south Tien-Shan. Acta Palaeontol. Pol. 25:375-380.
- 89 Ettensohn, F.R. 1981. *Crininicaminus haneyensis*, a new agglutinated worm tube from the Chesterian of east-central Kentucky. Jour. Paleontol. 55:479-482.
- 90 Fahraeus, L.E. 1976. Conodontophorid ecology and evolution related to global tectonics. Pp. 11-26. In: C.R. Barnes, ed. Conodont Paleoecology. Geol. Assoc. Canada, Spec. Pap. 15.
- 91 \_\_\_\_\_, and G.S. Nowlan. 1978. Franconian (Late Cambrian) to Early Champlainian (Middle Ordovician) conodonts from the Cow Head Group, western Newfoundland. Jour. Paleontol. 52:444-471.
- 92 Fedorov, A.B., L.I. Egorova, and V.E. Savitskiy. 1979. Pervaja nakhodka drevneishikh trilobitov v nizhiei chasti stratotipa Tommotskogo Jarusa nizhnego Kembrija (r. Aldan). Dokl. Akad. Nauk SSSR 249:1188-1190.
- 93 Feldmann, R.M., F.R. Schram, and N.J. Copeland. 1977. *Palaeopalaemon newberryi*, the earliest known (Late Devonian) decapod crustacean. Geol. Soc. Amer. Abstr. Programs 9:973.
- 94 Finks, R.M. 1960. Late Paleozoic sponge faunas of the Texas region. The siliceous sponges. Amer. Mus. Nat. Hist. Bull. 120:1-160.
- 95 \_\_\_\_\_, 1971. A new Permian eutaxicladine demosponge, mosaic evolution, and the origin of the Dicranocladina. Jour. Paleontol. 45:977-997.
- 96 \_\_\_\_\_, 1971. Sponge zonation in the west Texas Permian section. Pp. 285-300. In: J.T. Dutro, ed. Paleozoic Perspectives: A Paleontological Tribute to G. Arthur Cooper. Smithson. Contrib. Paleobiology 3.
- 97 Fisher, D.C. 1975. Evolution and Functional Morphology of the Xiphosurida. unpubl. Ph.D. dissertation; Harvard Univ.; Cambridge, Massachusetts, 396 pp.
- 98 Flower, R.H. 1964. The nautiloid order Ellesmeroceratida (Cephalopoda). N. Mex. Bur. Mines Miner. Resour. Mem. 12, 231 pp.
- 99 \_\_\_\_\_, 1976. ordovician cephalopod faunas and their role in correlation. Pp. 523-552. In: M.G. Bassett, ed. The Ordovician System. Palaeontol. Assoc., Univ. Wales Press and Nat. Mus. Wales.
- 0 Flügel, von H.W. 1970. Die Entwicklung der rugosen Korallen im hohen Perm. Verh. Geol. B.-A. 1:146-161.
- 1 \_\_\_\_\_, 1973. *Peroniella baloghi*, a new Inozoa from the Upper Permian of the Bükk-Mountains (Hungary). Acta Mineral. Petrogr., Szeged 21:49-53.
- 1a Forest, J., M. de St. Laurent, and F.A. Chace, Jr. 1976. *Neoglypheha inopinata*: A crustacean "living fossil" from the Philippines. Science 192:884.
- 1b Forey, P.L. 1973. A revision of the elopiform fishes, fossil and recent. Br. Mus. (Nat. Hist.) Bull., Geol. Suppl. 10, 222 pp.
- 1c Fortey, R.A., and R.M. Owens. 1975. Proetida — a new order of trilobites. Fossils and Strata, No. 4:227-240.
- 1d \_\_\_\_\_, and J.E. Whittaker. 1976. *Janospira* — an Ordovician fossil in search of a phylum. Lethaia 9:397-403.
- 1e Foster, M.W. 1979. Soft-bodied coelenterates in the Pennsylvanian of Illinois. Pp. 191-268. In: M.H. Nitecki, ed. Mazon Creek Fossils. Academic Press, New York.

- 106 Frest, T.J., and H.L. Strimple. 1980. New diminutive camerate crinoids from the Ludlow of Oklahoma. *Geol. Mag.* 117:255-266.
- 107 \_\_\_\_\_, and \_\_\_\_\_. 1981. New camerate crinoids from the Silurian of North America. *Jour. Paleontol.* 55:639-655.
- 108 \_\_\_\_\_, \_\_\_\_\_, and C.C. Coney. 1979. Paracrinoids (Platycystitidae) from the Benbolt Formation (Blackriverian) of Virginia. *Jour. Paleontol.* 53:380-398.
- 109 \_\_\_\_\_, \_\_\_\_\_, and B.J. Witzke. 1980. New Comarocystitida (Echinodermata: Paracrinoidae) from the Silurian of Iowa and Ordovician of Oklahoma. *Jour. Paleontol.* 54:217-228.
- 110 Fritz, W.H. 1970. Cambrian faunas. P. 593-600. In: R.J. Douglas, ed. *Geology and Economic Minerals of Canada*. Dept. Energy, Mines Res., Canada.
- 111 Furnish, W.M. 1973. Permian stage names. Pp. 522-548. In: A. Logan and L.V. Hill, eds. *The Permian and Triassic Systems and their Mutual Boundary*. Can. Soc. Pet. Geol.; Calgary, Alberta.
- 112 Fürsich, F.T., and J. Wendt. 1977. Biostratinomy and palaeoecology of the Cassian Formation (Triassic) of the southern Alps. *Palaeogeogr., Palaeoclimatol., Palaeoecol.* 22:257-323.
- 113 Gardiner, B.C., and R.S. Miles. 1975. Devonian fish of the Gogo Formation, Western Australia. *Colloque Int. C.N.R.S. No. 218*, pp. 78-79.
- 114 Germs, G.J.B. 1972. Trace fossils from the Nama Group, South-West Africa. *Jour. Paleontol.* 46:864-870.
- 115 \_\_\_\_\_. 1972. New shelly fossils from the Nama Group, South West Africa. *Am. J. Sci.* 272:752-761.
- 115a Ghiurca, V., and N. Mongereau. 1981. La faune de bryozoaires éocènes d'Europe orientale. Pp. 93-100. In: G.P. Larwood and C. Nielson, eds. *Recent and Fossil Bryozoa*. Olsen and Olsen; Fredensborg, Denmark.
- 116 Giammona, C.P., and R.J. Stanton, Jr. 1980. Octocorals from the Middle Eocene Stony City Formation, Texas. *Jour. Paleontol.* 54:71-80.
- 117 Glaessner, M.F. 1958. New fossils from the base of the Cambrian in South Australia. *Trans. Roy. Soc. S. Aust.* 81:185-188.
- 118 \_\_\_\_\_. 1971. The genus *Conomedusites* Glaessner & Wade and the diversifications of the Cnidaria. *Paläontol. Z.* 45:7-17.
- 119 \_\_\_\_\_. 1976. Early Phanerozoic annelid worms and their geological and biological significance. *Jour. Geol. Soc. London* 132:259-275.
- 120 \_\_\_\_\_. 1979. An echiurid worm from the late Precambrian. *Lethaia* 12:121-124.
- 121 \_\_\_\_\_. 1979. Lower Cambrian Crustacea and annelid worms from Kangaroo Island, South Australia. *Alcheringa* 3:21-32.
- 122 \_\_\_\_\_ and M. Wade. 1966. The late Precambrian fossils from Ediacara, South Australia. *Palaeontology* 9:599-628.
- 123 \_\_\_\_\_, and \_\_\_\_\_. 1971. *Praecambridium* — a primitive arthropod. *Lethaia* 4:71-77.

- 124 Glaessner, M.F., and M.R. Walter. 1975. New Precambrian fossils from the Arumbera Sandstone, Northern Territory, Australia. *Alcheringa* 1:59-69.
- 125 Gooday, A.J., and G. Becker. 1979. Ostracods in Devonian biostratigraphy. Pp. 193-197. In: M.R. House, C.T. Scrutton, and M.G. Bassett, eds. *The Devonian System*. Palaeontol. Assoc. Lond., Spec. Pap. Palaeontol. 23.
- 126 Goody, P.C. 1969. The relationships of certain Upper Cretaceous teleosts with special reference to the myctophoids. *Br. Mus. (Nat. Hist.) Bull., Geol. Suppl.* 7, 225 pp.
- 127 Gramm, M.N. 1977. A new family of Paleozoic ostracods. *Palaeontology* 20:475-482.
- 128 Grant, R.E. 1970. Brachiopods from Permian-Triassic Boundary beds and age of Chhidru Formation, West Pakistan. Pp. 117-153. In: B. Kummel and C. Teichert, eds. *Stratigraphic Boundary Problems*. Univ. Kansas Press; Lawrence, Kansas.
- 128a Gutschick, R.C., W.F. Canis, and K.G. Brill, Jr. 1967. Kinderhook (Mississippian) holothurian sclerites from Montana and Missouri. *Jour. Paleontol.* 41:1461-1480.
- 129 Hallam, A. 1977. Jurassic bivalve biogeography. *Paleobiology* 3:58-73.
- 129a\_\_\_\_\_. 1981. The end-Triassic bivalve extinction event. *Palaeogeogr., Palaeoclimat., Palaeoecol.* 35:1-44.
- 30 Halstead, L.B., and S. Turner. 1973. Silurian and Devonian ostracoderms. Pp. 67-80. In: A. Hallam, ed. *Atlas of Paleobiogeography*. Elsevier; New York.
- 31 Harland, W.B. 1974. The Pre-Cambrian-Cambrian Boundary. Pp. 15-42. In: C.H. Holland, ed. *Cambrian of the British Isles, Norden, and Spitsbergen*. Wiley-Interscience; New York.
- 32 \_\_\_\_\_, et al., eds. 1967. *The Fossil Record*. Geol. Soc. London; London, 828 pp.
- 33 Hartman, W.D., J.W. Wendt, and F. Wiedenmayer. 1980. *Fossil and Living Sponges. Sedimenta VIII*. Univ. Miami; Miami, Florida, 274 pp.
- 34 Haude, R., and F. Langenstrassen. 1976. *Rotasaccus dentifer* n. gen. n. sp., ein devonischer Ophiocistiode (Echinodermata) mit "holothuroiden" Wandskleriten und "echinoiden" Kauapparat. *Paläontol. Z.* 50:130-150.
- 35 Hayami, I. 1975. A Systematic Survey of the Mesozoic Bivalvia from Japan. Univ. Mus., Univ. Tokyo, Bull. No. 10, 249 pp.
- 6 \_\_\_\_\_, and T. Kase. 1977. A Systematic Survey of the Paleozoic and Mesozoic Gastropoda and Paleozoic Bivalvia from Japan. Univ. Mus., Univ. Tokyo, Bull. No. 13, 156 pp.
- 7 Henderson, R.A., and J.H. Shergold. 1971. *Cyclocystoides* from early Middle Cambrian rocks of northwestern Queensland, Australia. *Palaeontology* 14:704-710.
- 7a Hoare, R.D., R.H. Mapes, and C.J. Brown. 1982. Some Mississippian and Pennsylvanian rostroconchs from the midcontinent region. *Jour. Paleontol.* 56:123-131.
- 3 Holloway, D.J. 1980. Middle Silurian trilobites from Arkansas and Oklahoma U.S.A. — Part 1. *Palaeontographica* 170:1-85.
- 9 House, M.R. 1979. Biostratigraphy of the early Ammonoidea. Pp. 263-280. In: M.R. House, C.T. Scrutton, and M.G. Bassett, eds. *The Devonian System*. Palaeontol. Assoc. Lond., Spec. Pap. Palaeontol. 23.
- 1\_\_\_\_\_ 1981. On the origin, classification and evolution of the early Ammonoidea.

- Pp. 3-36. In: M.R. House and J.R. Senior, eds. *The Ammonoidea. Syst. Assoc. Spec.* Vol. 18. Academic Press; New York.
- 141 Hupé, P. 1960. Sur le cambrien inférieur du Maroc. 21st Int. Geol. Cong., Copenhagen Rep. Pt. 8, p. 75-85.
- 142 Hurcewicz, H. 1968. Siliceous sponges from the Upper Cretaceous of Poland: Part II. Monazonia and Triaxonnia. *Acta Paleontol. Pol.* 8:3-8.
- 143 Ivanovskij, A.B. 1978. The system of the Anthozoa. *Paleontol. Jour.* 12:23-27.
- 144 Jaanusson, V., S. Laufeld, and R. Skoglund, eds. 1979. Lower Wenlock faunal and floral dynamics — Vattenfallet section, Gotland. *Sveriges Geologiska Undersökning*; Uppsala, 294 pp.
- 145 Jaeger, H., and A. Martinsson. 1967. Remarks on the problematic fossil *Xenusion auerswaldae*. *Geol. Fören. Stockholm Förh.* 88:435-452.
- 146 Janjauskas, T.V. 1969. Pterociatidi nizhnego Kembriya Krasnojarskogo (Vostochnii Sajan). Pp. 114-157. In: I.T. Zhuravleva, ed. *Biostratigrafiya i Paleontologiya Nizhnego Kembriya Sibiri i Dalbiego Vostoka*. Nauka; Moscow.
- 147 Jeletzky, J.A. 1965. Taxonomy and phylogeny of fossil Coleoidea (= Dibranchiata). *Geol. Surv. Can. Pap.* 65-2:72-76.
- 148 Jell, P.A. 1974. Faunal provinces and possible planetary reconstruction of the Middle Cambrian. *Jour. Geol.* 82:319-350.
- 149 \_\_\_\_\_. 1979. *Plumulites* and the machaeridian problem. *Alcheringa* 3:253-260.
- 150 \_\_\_\_\_. 1980. Earliest known pelecypod on Earth — a new Early Cambrian genus from South Australia. *Alcheringa* 4:233-239.
- 151 \_\_\_\_\_. 1980. Two arthropods from the Lancefieldian (Lal) of central Victoria *Alcheringa* 4:37-46.
- 152 \_\_\_\_\_. 1981. *Thambetolepis delicata* gen. et sp. nov., an enigmatic fossil from the Early Cambrian of South Australia. *Alcheringa* 5:85-93.
- 153 \_\_\_\_\_, and J.S. Jell. 1976. Early Middle Cambrian corals from western New South Wales. *Alcheringa* 1:181-195.
- 154 Johnson, G.D., and J. Zidek. 1981. Late Paleozoic phyllodont tooth plates. *Jour. Paleontol.* 55:524-536.
- 155 Johnson, H., and S.K. Fox, Jr. 1968. Diplopoda from Lower Silurian of North America. *Science* 162:119-120.
- 156 Johnson, R.G., and E.S. Richardson, Jr. 1966. A remarkable Pennsylvanian fauna from the Mazon Creek area, Illinois. *Jour. Geol.* 74:626-631.
- 157 \_\_\_\_\_, and \_\_\_\_\_. 1968. The Essex Fauna and medusae. *Fieldiana: Geol.* 12:109-115.
- 158 Jones, D., and I. Thompson. 1977. Echiura from the Pennsylvanian Essex Fauna of northern Illinois. *Lethaia* 10:317-325.
- 159 Jones, P.J. 1968. Upper Devonian Ostracoda and Eridostraca from the Bonaparte Gulf Basin, northwestern Australia. *Aust. Bur. Miner. Res., Geol. Geophys. Bull.* 99:1-108.
- 160 \_\_\_\_\_, and K.G. McKenzie. 1980. Queensland Middle Cambrian Bradoriida

- (Crustacea): New taxa, palaeobiogeography and biological affinities. *Alcheringa* 4:203-226.
- 161 Kaźmierczak, J., and G. Hillmer. 1974. Schlerospunge nature of the lower Haute-Rivian "bryozoan" *Neuropora pustulosa* (Roemer 1839) from western Germany. *Acta Palaeontol. Pol.* 19:443-453.
- 162 Keller, B.M., and M.A. Fendonkin. 1976. Novye nahodki okamenelastej v Valdaiksej serii Dokembria po R. Suzme. *Izvestie Akad. Nauk SSSR. ser. geolog* for 1976, No. 3, Pp. 38-44.
- 163 \_\_\_\_\_, and A. Yu. Rozanov, eds. 1979. Upper Precambrian and Cambrian Paleontology of East-European Platform (in Russian). Nauka; Moscow, 211 pp.
- 64 Kelly, S.M., and J.K. Pope. 1979. A new camerata crinoid from the Upper Ordovician of Indiana. *Jour. Paleontol.* 53:416-420.
- 65 Kennedy, W.J., and O.H. Bayliss. 1977. The earliest tissotiid ammonite. *Paleontology* 20:901-906.
- 65a Kesling, R.U., and C.R.C. Paul. 1971. *Agostocrinus* and *Acolocrinus*, two new Ordovician crinoids with peculiar ray and respiratory structures. *Univ. Michigan Contrib. Museum Paleontol.* 23:221-237.
- 36 Kielan-Jaworowska, Z. 1966. Polychaete jaw apparatuses from the Ordovician and Silurian of Poland and a comparison with modern forms. *Palaeontol. Polonica* 16:1-152.
- 37 Kier, P.M. 1974. Evolutionary trends and their functional significance in the post-Paleozoic echinoids. *Jour. Paleontol.* 48(Supplement):1-95.
- 7a \_\_\_\_\_. 1982. Rapid evolution in echinoids. *Paleontology* 25:1-9.
- 8 Kling, S.A. 1978. Radiolaria. Pp. 203-244. In: B.U. Haq and A. Boersma, eds. *Introduction to Marine Micropaleontology*. Elsevier; New York.
- 3 \_\_\_\_\_, and W.E. Reif. 1969. The Paleozoic history of amphidisc and hemidisc sponges: New evidence from the Carboniferous of Uruguay. *Jour. Paleontol.* 43:1429-1434.
- 1 Kobayashi, T. 1956. The Cambrian of Korea and its relation to the other Cambrian territories. Pp. 343-362. In: J. Rogers, ed. *El Sistema Cámbrico*. 20th Int. Geol. Cong., Mexico.
- \_\_\_\_\_. 1966. The Cambro-Ordovician formations and faunas of South Korea. Part X. Stratigraphy of the Chosen Group in Korea and South Manchuria. Section B. The Chosen Group of North Korea and northeast China. *Jour. Fac. Sci., Univ. Tokyo, Sec. II*, 16:209-311.
- \_\_\_\_\_. 1967. The Cambro-Ordovician formations and faunas of South Korea. Part X. Stratigraphy of the Chosen Group in Korea and South Manchuria. Section C. The Cambrian of eastern Asia and other parts of the continent. *Jour. Fac. Sci., Univ. Tokyo, Sec. II*, 16:381-534.
- \_\_\_\_\_. 1971. The Cambro-Ordovician faunal provinces and the interprovincial correlation. *Jour. Fac. Sci., Univ. Tokyo, Sec. II*, 18:129-299.
- Kolata, D.R. 1975. Middle Ordovician echinoderms from northern Illinois and southern Wisconsin. *Paleontol. Soc. Mem.* 7, 74 pp.

- 175 \_\_\_\_\_, H.L. Strimple, and C.O. Levorsen. 1977. Revision of the Ordovician carpoid family Iowacystidae. *Palaeontology* 20:529-557.
- 176 Kollárova-Andrusovová, V., and M. Kochanová. 1973. Molluskenfauna des Bleskový Prameň bei Drnava. Bratislava, 239 pp.
- 177 Konevá, S.S. 1978. The first record of ostracodes in the Lower Cambrian of Kazakhstan. *Paleontol. Jour.* 12:137-138.
- 178 Konyashkov, K.N. 1972. K probleme yarusnogo deleniya nizhnego Kembriya. Pp. 7-14. In: I.T. Zhuravleva, ed. Problemy Biostratigrafi i Paleontologii Nizhnego Kembriya. Nauka; Moscow.
- 179 Kozur, H. 1970. Fossile Hirudinea aus dem Oberjura von Bayern. *Lethaia* 3:225-232.
- 180 Kruse, P.D. 1978. New Archaeocyatha from the Early Cambrian of the Mt. Wright area, New South Wales. *Aicheringa* 2:27-48.
- 181 Kukalová-Peck, J. 1973. A Phylogenetic Tree of the Animal Kingdom (Including Orders and Higher Categories). *Nat. Mus. Canada, Publ. Zool.* No. 8, 78 pp.
- 182 Kullmann, J. 1981. Carboniferous goniatites. Pp. 37-48. In: M.R. House and J.R. Senior, eds. *The Ammonoidea. Syst. Assoc. Spec. Vol. 18*. Academic Press; New York.
- 183 Landing, E. 1977. "Prooneotodus" *tenuis* (Müller, 1959) apparatuses from the Taconic allochthon, eastern New York: Construction, taphonomy and the protoconodont "super-tooth" model. *Jour. Paleontol.* 51:1072-1084.
- 183a Lane, N.G. 1970. Lower and Middle Ordovician crinoids from west-central Utah. *Brigham Young Univ. Geol. Studies* 17:3-17.
- 184 Larson, K. 1979. Silurian tentaculitids from Gotland and Scania. *Fossils and Strata*, No. 11, 180 pp.
- 185 Laub, R.S. 1979. The corals of the Brassfield Formation (mid-Llandovery; Lower Silurian) in the Cincinnati Arch region. *Bull. Amer. Paleontol.* 75:1-457.
- 186 Lendzion, K. 1975. Fauna of the *Mobergella* Zone in the Polish Lower Cambrian. *Kwartalnik geologiczny, Warszawa* 19(2):237-242.
- 187 Lessovaya, A.L. 1974. Filogeniya i sistematika semeistv Clathrodictyidae, Simplexodictyidae, Tienodictyidae, Geronostromatidae. Pp. 51-54. In: B.S. Sokolov, ed. *Ancient Cnidaria*, v. 1. Nauka; Moscow.
- 188 Lindström, M. 1970. A suprageneric taxonomy of the conodonts. *Lethaia* 3:427-445.
- 189 Linsley, R.M. 1979. Gastropods of the Devonian. Pp. 249-254. In: M.R. House, C.T. Scrutton, and M.G. Bassett, eds. *The Devonian System. Palaeontol. Assoc. Lond. Spec. Pap. Palaeontol.* 23.
- 190 Lipps, J.H., and E. Mitchell. 1976. Trophic model for the adaptive radiations and extinctions of pelagic marine mammals. *Paleobiology* 2:147-155.
- 191 Lochman, C., and D. Duncan. 1944. Early Upper Cambrian faunas of central Montana. *Geol. Soc. Amer. Spec. Pap.* 54, 181 pp.
- 192 Lochman-Balk, C. 1971. The Cambrian of the craton of the United States. Pp. 79-167. In: C.H. Holland, ed. *Cambrian of the New World*. Wiley, Interscience; New York.
- 193 \_\_\_\_\_, and J.L. Wilson. 1958. Cambrian biostratigraphy in North America. *Jour. Paleontol.* 32:312-350.

- 194 \_\_\_\_\_, and \_\_\_\_\_. 1967. Stratigraphy of Upper Cambrian-Lower Ordovician subsurface sequence in Williston Basin. Am. Assoc. Pet. Geol. Bull. 51:883-917.
- 195 Lu Y.H., Chu C.L., Chien Y.Y., Lin H.L., Chow T.Y., and Yuan K.S. 1974. Bioenvironmental control hypothesis and its application to the Cambrian biostratigraphy and paleozoogeography. Nanking Inst. Geol. Paleontol. Mem. 5:27-110.
- 196 Martinsson, A. 1974. The Cambrian of Norden. Pp. 185-283. In: C.H. Holland, ed. Cambrian of the British Isles, Norden, and Spitsbergen. Wiley, Interscience; New York.
- 197 Matthew, G.F. 1886. Illustrations of the fauna of the St. John Group, No. 3, descriptions of new genera and species. Trans. R. Soc. Canada 3(4):29-84.
- 198 Matthews, S.C., and V.V. Missarzhevsky. 1975. Small shelly fossils of late Precambrian and early Cambrian age: A review of recent work. Jour. Geol. Soc. Lond. 131:289-304.
- 199 McAlester, A.L. 1965. Systematics, affinities and life habits of *Babinka*. Palaeontology 8:231-241.
- 199a McIntosh, G.C. 1981. *Tagenocrinus* and *Lecocrinus*, two new flexible crinoid genera from the Silurian and Devonian of North America. Jour. Paleontol. 55:1103-1108.
- 199b MacKinnon, D.I. 1982. *Tuarangia paparua* n. gen. and n. sp., a late Middle Cambrian pelecypod from New Zealand. Jour. Paleontol. 56:589-598.
- 200 McLeod, J.D. 1978. The oldest bryozoans: New evidence from the Early Ordovician. Science 200:771-773.
- 201 Melton, W., and H.W. Scott. 1972 (1973). Conodont-bearing animals from the Bear Gulch Limestone, Montana. Geol. Soc. Amer. Spec. Pap. 141:31-65.
- 202 Meshkova, N.P. 1969. K voprosu o paleontologicheskoi kharakteristike nizhnetskobriiskikh otlozhennii Sibirskoi Platformi. Pp. 158-174. In: I.T. Zhuravleva, ed. Biostratigrafiya i Paleontologiya Nizhnego Kembriya Sibiri i Dalbiego Vostoka. Nauka; Moscow.
- 203 Mierzejewski, P., and G. Mierzejewska. 1975. Zenognath type of polychaete jaw apparatuses. Acta Palaeontol. Pol. 20:437-442.
- 204 Miller, J.F. 1975. Third day. Pp. 16-30. In: R.A. Robison, ed. Central and Western Utah: Cambrian Paleontology and Depositional Environments. Geol. Soc. Amer. 1975 Field Trip 9, Guidebook.
- 205 Missarzhevsky, V.V. 1974. Novie dannye o drevneishcikh okamelenostyakh rannego Kembriya Siberskoi Platformi. Pp. 179-189. In: I.T. Zhuravleva and A.Yu. Rozanov, eds. Biostratigrafiya i Paleontologiya Nizhnego Kembriya Evropy i Severnoy Azii. Nauka; Moscow.
- 206 Moore, R.C., C.G. Lalicker, and A.G. Fisher. 1952. Invertebrate Fossils. McGraw-Hill; New York, 766 pp.
- 207 \_\_\_\_\_, and H.L. Strimple. 1973. Lower Pennsylvanian (Morrowan) crinoids from Arkansas, Oklahoma, and Texas. Univ. Kansas Paleontol. Contrib. Art. 60, 84 pp.
- 208 \_\_\_\_\_, C. Teichert, and R.A. Robison, eds. 1953-1981. Treatise on Invertebrate Paleontology. Geol. Soc. Amer. and Univ. Kansas Press; Lawrence, Kansas.
- 209 Morozova, I.P. 1970. Mshanki pozonei Permi. Acad. Sci. USSR, Trans. Paleontol. Inst. 122:1-346.

- 210 Morris, N.J. 1979. On the origin of the Bivalvia. Pp. 381-414. In: M.R. House, ed. The Origin of Major Invertebrate Groups. Syst. Assoc. Spec. Vol. 12. Academic Press; New York.
- 211 \_\_\_\_\_, 1980. A new Lower Ordovician bivalve family, the Thoralidae (?Nuculoida), interpreted as actinodont deposit feeders. Br. Mus. (Nat. Hist.) Bull., Geol. Suppl. 34:265-272.
- 212 Moy-Thomas, J.A., and R.S. Miles. 1971. Palaeozoic Fishes, 2nd ed. Saunders; Philadelphia, 259 pp.
- 213 Müller, A.H. 1963-70. Lehrbuch der Paläozoologie. Veb Gustav Fischer Verlag; Jena.
- 214 Müller, J.K. 1977. *Palaeobotryllus* from the Upper Cambrian of Nevada — a probable ascidian. Lethaia 10:107-118.
- 215 Müller, K.J. 1981. Arthropods with phosphatized soft parts from the Upper Cambrian "Orsten" of Sweden. Pp. 147-151. In: M.E. Taylor, ed. Short Papers for the Second International Symposium on the Cambrian System. U.S. Geol. Surv. Open-File Rpt. 81-743.
- 216 Nakazawa, K., and B. Runnegar. 1973. The Permian-Triassic Boundary; A crisis for bivalves? Pp. 608-621. In: A. Logan and L.V. Hills, eds. The Permian and Triassic Systems and their Mutual Boundary. Canadian Soc. Pet. Geol.; Calgary, Alberta.
- 217 Nazarov, B.B. 1975. Lower and middle Paleozoic radiolarians of Kazakhstan (in Russian). Trudy Geol. Inst. Akad. Nauk. SSSR 275:1-202.
- 218 Nelson, J.S. 1976. Fishes of the World. Wiley-Interscience; New York, 416 pp.
- 219 Nitecki, M.H., and F.R. Schram. 1976. *Etacystis communis*, a fossil of uncertain affinities from the Mazon Creek Fauna (Pennsylvanian of Illinois). Jour. Paleontol. 50:1157-1161.
- 219a \_\_\_\_\_, and A. Solem. 1973. A problematic organism from the Mazon Creek (Pennsylvanian) of Illinois. Jour. Paleontol. 47:903-907.
- 220 North, F.K. 1971. The Cambrian of Canada and Alaska. Pp. 219-324. In: C.H. Holland, ed. Cambrian of the New World. Wiley-Interscience; New York.
- 221 Obut, A.M. 1974. New graptolites from the Middle Cambrian of the Siberian Platform. Palaeont. Assoc. Lond., Spec. Pap. Palaeontol. 13:9-13.
- 222 Oekentorp, Kl., C. Montenat, and H. Fontaine. 1978. Eine kleine Korallenfauna aus dem unteren Operperm von Saïq, Oman (Arabische Halbinsel). N. Jb. Geol. Paläontol. Abh. 155:374-397.
- 223 Okulitch, V.J., and W.G. Bell. 1955. *Gallatinospongia*, a new siliceous sponge from the Upper Cambrian of Wyoming. Jour. Paleontol. 29:460-461.
- 224 Olempska, E. 1979. Middle to Upper Devonian Ostracoda from the southern Holy Cross Mountains, Poland. Palaeontol. Polonica 40:55-62.
- 225 Oliver, W.A., and A.E.H. Pedder. 1979. Rugose corals in Devonian stratigraphical correlation. Pp. 233-248. In: M.R. House, C.T. Scrutton, and M.G. Bassett, eds. The Devonian System. Palaeont. Assoc. Lond., Spec. Pap. Palaeontol. 23.
- 226 Öpik, A.A. 1963. Early Upper Cambrian fossils from Queensland. Aust. Bur. Miner. Res., Geol. Geophys. Bull. 64:1-133.
- 227 \_\_\_\_\_, 1967. The Mindyallan fauna of north-western Queensland. Aust. Bur. Miner. Res., Geol. Geophys. Bull. 74:1-404.

- 228 \_\_\_\_\_. 1968. Ordian (Cambrian) Crustacea Bradoriida of Australia. Aust. Bur. Miner. Res., Geol. Geophys. Bull. 103:1-44.
- 229 Orlov, Yu. A., ed. 1958-64. Osnovy Paleontologii. Moscow.
- 230 Ossian, C.R. 1973. New Pennsylvanian scyphomedusan from western Iowa. Jour. Paleontol. 47:990-995.
- 231 Palij, V.M., E. Posti, and M.A. Fedonkin. 1979. Soft-bodied Metazoa and trace fossils of the Vendian and Lower Cambrian (in Russian). Pp. 49-82. In: B.M. Keller and A. Yu. Rozanov, eds. Upper Precambrian and Cambrian Paleontology of East-European Platform. Nauka; Moscow.
- 232 Palmer, A.R. 1971. The Cambrian of the Great Basin and adjacent areas, western United States. Pp. 1-78. In: C.H. Holland, ed. Cambrian of the New World. Wiley-Interscience; New York.
- 233 Palmer, C.P. 1974. A supraspecific classification of the scaphopod Mollusca. Veliger 17:115-123.
- 234 Parsley, R.L., and L.W. Mintz. 1975. North American Paracrinoida (Ordovician: Paracrinzoa, new, Echinodermata). Bull. Amer. Paleontol. 68:1-115.
- 235 Patterson, C. 1973. Interrelationships of holosteans. Pp. 233-306. In: P.H. Greenwood et al., eds. Interrelationships of Fishes. Academic Press; London.
- 236 \_\_\_\_\_, and D.E. Rosen. 1977. Review of ichthyodectiforms and other Mesozoic teleost fishes and the theory and practice of classifying fossils. Amer. Mus. Nat. Hist. Bull. 158:85-172.
- 237 Paul, C.R.C. 1968. Morphology and function of dichoporite pore-structures in cystoids. Palaeontology 11:697-730.
- 238 \_\_\_\_\_. 1976. Palaeogeography of primitive echinoderms in the Ordovician. Pp. 553-574. In: M.G. Bassett, ed. The Ordovician System. Palaeontol. Assoc., Univ. Wales Press and Nat. Mus. Wales.
- 239 Pawson, D.L. 1980. Holothuroidea. Pp. 175-189. In: T.W. Broadhead and J.A. Waters, eds. Echinoderms. Notes for a Short Course. Univ. Tenn. Dept. Geol. Sci., Stud. in Geol. 3.
- 239a Percival, I.G. 1979. Ordovician plectambonitacean brachiopods from New South Wales. Alcheringa 3:91-116.
- 40 Perry, D.G., and B.D.E. Chatterton. 1979. Late Early Triassic brachiopod and conodont fauna, Thaynes Formation, southeastern Idaho. Jour. Paleontol. 53:307-319.
- 41 Pflug, H.D. 1972. The Phanerozoic-Cryptozoic Boundary and the origin of Metazoa. 24th Int. Geol. Cong., Montreal. Sec. 1, pp. 58-67.
- 42 Pitrat, C.W. 1973. Vertebrates and the Permo-Triassic extinction. Palaeogeogr., Palaeoclimatol., Palaeoecol. 14:249-264.
- 43 Piveteau, J., ed. 1952-69. Traité de Paléontologie. Masson et Cie; Paris.
- 44 Plotnick, R. 1974. Middle Paleozoic scorpions. Unpubl. ms, 17 pp.
- 45 Pocock, K.J. 1970. The Emuellidae, a new family of trilobites from the Lower Cambrian of South Australia. Palaeontology 13:522-562.
- 46 Pohowsky, R.A. 1978. The boring ctenostomate Bryozoa: Taxonomy and paleobiology based on cavities in calcareous substrata. Bull. Amer. Paleontol. 73:1-192.

- 247 Pojeta, J., Jr. 1975. *Fordilla troyensis* Barrande and early pelecypod phylogeny. Bull. Amer. Paleontol. 67:363-384.
- 248 \_\_\_\_\_, and J. Gilbert-Tomlinson. 1977. Australian Ordovician pelecypod molluscs. Aust. Bur. Miner. Res., Geol. Geophys. Bull. 174:1-64.
- 249 \_\_\_\_\_, J. Gilbert-Tomlinson, and J.H. Shergold. 1977. Cambrian and Ordovician rostroconch molluscs from northern Australia. Aust. Bur. Miner. Res., Bull. 171:1-54.
- 250 \_\_\_\_\_, and B. Runnegar. 1976. The paleontology of rostroconch mollusks and the early history of the phylum Mollusca. U.S. Geol. Surv. Prof. Pap. 968, 88 pp.
- 251 Pope, J.K. 1975. Evidence for relating the Lepidocoleidae, machaeridian echinoderms, to the mitrate carpoids. Bull. Amer. Paleontol. 67:385-406.
- 252 Pospelov, A.G., A.S. Boyarinova, N.A. Aksarina, Yu. S. Nadler, and E.S. Fedyanova. 1972. Opornii razrez nizhnego Kembriya po r. Kie v Kuzneckom Alatau. Pp. 222-232. In: I.T. Zhuravleva, ed. Problemy Biostratigrafii i Paleontologii Nizhnego Kembriya Siberi. Nauka; Moscow.
- 253 Qian Y. 1977. *Hyolitha* and some problematica from the Lower Cambrian Meishucun Stage in central and southwestern China (in Chinese). Acta Palaeontol. Sin. 16:255-278.
- 254 Raasch, G.O. 1939. Cambrian Merostomata. Geol. Soc. Amer. Spec. Pap. 19, 146 pp.
- 255 Raup, D.M. 1962. The phylogeny of calcite crystallography in echinoids. Jour. Paleontol. 36:793-810.
- 256 Reid, R.E. 1968. Hexactinellid faunas in the Chalk of England and Ireland. Geol. Mag. 105:15-22.
- 257 Reif, W.-E. 1978. Tooth enameloid as a taxonomic criterion. 2. Is "Dalatias" barnstonensis Sykes, 1971 (Triassic, England) a squalomorphic shark? N. Jb. Geol. Paläontol. Mh. 1:42-58.
- 258 Remane, J. 1978. Calpionellids. Pp. 161-170. In: B.U. Haq and A. Boersma, eds. Introduction to Marine Micropaleontology. Elsevier; New York.
- 258a Repenning, C.A., C.E. Ray, and D. Grigorescu. 1979. Pinniped biogeography. Pp. 357-369. In: J. Gray and A.J. Boucot, eds. Historical Biogeography, Plate Tectonics and the Changing Environment. Oregon State Univ. Press; Corvallis, Oregon.
- 259 Repetski, J.E. 1978. A fish from the Upper Cambrian of North America. Science 200:529-531.
- 260 \_\_\_\_\_. 1981. An Ordovician occurrence of *Utahphospha* Müller & Miller. Jour. Paleontol. 55:395-400.
- 261 Repina, L.N. 1972. Biogeography of Early Cambrian of Siberia according to trilobites. 1968 Proc. Paleontol. Union, 23rd Int. Geol. Cong., Prague, pp. 289-300.
- 262 \_\_\_\_\_. 1972. Trilobiti tarinskogo gorizonta razrezov nizhnego Kembriya r. Sukharikhi (Igarskii raion). Pp. 184-216. In: I.T. Zhuravleva, ed. Problemy Biostratigrafii i Paleontologii Nizhnego Kembriya Siberi. Nauka; Moscow.
- 263 \_\_\_\_\_. 1974. K voprosy o granitse nizhnego i srednego Kembriya Sibirskoy Platformy i sopredel'nykh territory. Pp. 76-103. In: I.T. Zhuravleva and A. Yu. Rozanov, eds. Biostratigrafiya i Paleontologiya Nizhnego Kembriya Evropy i Severnoy Asii. Nauka; Moscow.

- 264 Resser, C.E., and B.F. Howell. 1938. Lower Cambrian *Olenellus* Zone of the Appalachians. Geol. Soc. Amer. Bull. 49:195-248.
- 265 Richards, R.B. 1977. Patterns of evolution in the graptolites. Pp. 333-358. In: A. Hallem, ed. Patterns of Evolution. Elsevier: Amsterdam.
- 266 Richardson, E.S., Jr. 1966. Wormlike fossil from the Pennsylvanian of Illinois. Science 151:75-76.
- 267 Rigby, J.K. Two new Upper Devonian hexactinellid sponges from Alberta. Jour. Paleontol. 44:7-16.
- 268 \_\_\_\_\_, 1974. *Vaurealspongia* and *Twenhofeella*, two new brachiospongid hexactinellid sponges from the Ordovician and Silurian of Anticosti Island, Quebec. Can. J. Earth Sci. 11:1343-1349.
- 269 \_\_\_\_\_, 1976. Some observations on occurrences of Cambrian Porifera in western North America and their evolution. Pp. 51-60. In: R.A. Robison and A.J. Rowell, eds. Paleontology and Depositional Environments: Cambrian of Western North America. Brigham Young Univ. Geol. Stud. 23(2).
- 270 \_\_\_\_\_, 1980. The new middle Cambrian sponge *Vauxia magna* from the Spence Shale of northern Utah and the taxonomic position of the Vauxiidae. Jour. Paleontol. 54:234-240.
- 271 \_\_\_\_\_, and A.W. Potter. 1980. Ordovician sphinctozoan sponges from the Klamath Mountains, California, the first known early Paleozoic occurrence. Geol. Soc. Amer. Abstr. Program 12:509.
- 272 \_\_\_\_\_, and A.J. Swenson. 1972. Pennsylvanian sponges from the Oquirrh Group of central Utah. Brigham Young Univ. Geol. Stud. 19(2):127-132.
- 273 \_\_\_\_\_, and F.M. Terrell. 1973. Permian sponges from western Ellesmere Island, Arctic Canada. Can. J. Earth Sci. 10:1431-1443.
- 274 Ritchie, A., and J. Gilbert-Tomlinson. 1977. First Ordovician vertebrates from the Southern Hemisphere. Alcheringa 1:351-368.
- 275 Robison, R.A. 1964. Late Middle Cambrian faunas from western Utah. Jour. Paleontol. 38:510-566.
- 276 \_\_\_\_\_, 1971. Additional Middle Cambrian trilobites from the Wheeler Shale of Utah. Jour. Paleontol. 45:796-804.
- 276a \_\_\_\_\_, and B.C. Richards. 1981. Larger bivalve arthropods from the Middle Cambrian of Utah. Univ. Kansas Paleontol. Contrib. Pap. 106, 19 p.
- 277 \_\_\_\_\_, and J. Sprinkle. 1969. Ctenocystoidea: New class of primitive echinoderms. Science 166:1512-1514.
- 278 Rolfe, W.D.I., and D.W. Brett. 1969. Fossilization processes. Pp. 123-244. In: G. Eglinton and M.T.J. Murphy, eds. Organic Geochemistry. Springer-Verlag; New York.
- 279 \_\_\_\_\_, and V.A. Edwards. 1979. Devonian Arthropoda (Trilobita and Ostracoda excluded). Pp. 325-329. In: M.R. House, C.T. Scrutton, and M.G. Bassett, eds. The Devonian System. Palaeontol. Assoc. Lond., Spec. Pap. Palaeontol. 23.
- 280 Romer, A.S. 1966. Vertebrate Paleontology, 3rd ed. Univ. Chicago Press; Chicago, 468 pp.

- 281 Rosen, D.E. 1973. Interrelationships of higher euteleostean fishes. Pp. 397-514. In: P.H. Greenwood et al., eds. *Interrelationships of Fishes*. Academic Press; London.
- 282 Rosman, Kh. S. 1976. Zoogeographical belt zonality of north Asia according to Ordovician brachiopods (in Russian). 25th Int. Geol. Session. Nauka; Moscow.
- 283 Rosova, A.V. 1968. Biostratigrafiya i trilobity verknego Kembriya i niznego Ordovika severo-zapada Sibirskoy Platformy. *Trudy Inst. Geol. Geof. Sib. Otd. Akad. Nauk SSSR* 36:1-196.
- 284 Ross, J.P. 1967. Fossil problematica from Upper Ordovician, Ohio. *Jour. Paleontol.* 41:37-42.
- 285 Ross, J.R.P. 1979. Permian ectoprocts in space and time. Pp. 259-276. In: J. Gray and A.J. Boucot, eds. *Historical Biogeography, Plate Tectonics, and the Changing Environment*. Oregon State Univ. Press; Corvallis, Oregon.
- 285a\_\_\_\_\_. 1981. Late Palaeozoic ectoprot biogeography. Pp. 213-220. In: G.P. Larwood and C. Nielsen, eds. *Recent and Fossil Bryozoa*. Olsen and Olsen; Fredensborg, Denmark.
- 286 Rowell, A.J. 1977. Early Cambrian brachiopods from the southwestern Great Basin of California and Nevada. *Jour. Paleontol.* 51:68-85.
- 287 Rozanov, A. Yu. 1967. The Cambrian lower boundary problem. *Geol. Mag.* 104:415-434.
- 288 \_\_\_\_\_, and F. Debrenne. 1974. Age of archaeocyathid assemblages. *Am. J. Sci.* 274:833-848.
- 289 \_\_\_\_\_, V.V. Missarzhevsky, N.A. Volkova, L.G. Voronova, I.N. Krylov, B.M. Keller, I.K. Korolyut, K. Lendzion, P. Mikhniak, N.G. Pykhova, and A.D. Sidorov. 1969. Tommotsky yarus i problema nizhneye granitsy Kembria. *Trans. Geol. Inst. Acad. Sci. USSR* 206:5-380.
- 290 Runnegar, B. 1974. Evolutionary history of the bivalve subclass Anomalodesmata. *Jour. Paleontol.* 48:904-940.
- 291 \_\_\_\_\_, and P.A. Jell. 1976. Australian Middle Cambrian molluscs and their bearing on early molluscan evolution. *Alcheringa* 1:109-138.
- 292 \_\_\_\_\_, and \_\_\_\_\_. 1980. Australian Middle Cambrian molluscs; Corrections and additions. *Alcheringa* 4:111-113.
- 293 \_\_\_\_\_, and J. Pojeta. 1974. Molluscan phylogeny: The paleontological viewpoint. *Science* 186:311-317.
- 294 \_\_\_\_\_, J. Pojeta, Jr., M.E. Taylor, and D. Collins. 1979. New species of the Cambrian and Ordovician chitons *Matthevia* and *Chelodes* from Wisconsin and Queensland: Evidence for the early history of polyplacophoran mollusks. *Jour. Paleontol.* 53:1374-1395.
- 295 Rushton, A.W.A. 1974. The Cambrian of Wales and England. Pp. 43-121. In: C.H. Holland, ed. *Cambrian of the British Isles, Norden, and Spitsbergen*. Wiley-Interscience; New York.
- 296 \_\_\_\_\_. 1978. Fossils from the Middle-Upper Cambrian transition in the Nuneaton district. *Palaeontology* 21:245-284.
- 297 Samsonowicz, J. 1956. Cambrian paleogeography and the base of the Cambrian System in Poland. Pp. 127-160. In: J. Rogers, ed. *El Sistema Cámbrico*. 20th Int. Geol. Cong., Mexico.

- 298 Sarv, L. 1972. Development of Ordovician ostracodes of the East Baltic. 1968 Proc. Int. Paleontol. Union, 23rd Int. Geol. Cong., Prague, pp. 203-210.
- 299 Schallreuter, R. 1968. Zur Taxonomie und Phylogenie der Eridostraca (Ostrocoidea). Paläontol. Z. 42:105-119.
- 299a\_\_\_\_\_. 1981. Ordovizische Problematika, III. *Ancienta Ross*, 1967 aus Europa. Paläontol. Z. 55:209-218.
- 300 Schram, F.R. 1969. The stratigraphic distribution of the Paleozoic Eumalacostraca. Fieldiana:Geology 12:213-234.
- 301 \_\_\_\_\_. 1971. A strange arthropod from the Mazon Creek of Illinois and the trans Permo-Triassic Merostomoidea (Trilobitoidea). Fieldiana:Geology 20:85-102.
- 302 \_\_\_\_\_. 1973. Pseudocoelomates and a nemertine from the Illinois Pennsylvanian. Jour. Paleontol. 47:985-989.
- 303 \_\_\_\_\_. 1979. The Mazon Creek biotas in the context of a Carboniferous faunal continuum. Pp. 159-190. In: M.H. Nitecki, ed. Mazon Creek Fossils. Academic Press; New York.
- 304 \_\_\_\_\_. 1979. Worms of the Mississippian Bear Gulch Limestone of central Montana, USA. Trans. San Diego Soc. Nat. Hist. 19:107-120.
- 305 \_\_\_\_\_. 1980. Miscellaneous late Paleozoic Malacostraca of the Soviet Union. Jour. Paleontol. 54:542-547.
- 306 \_\_\_\_\_. R.M. Feldmann, and M.J. Copeland. 1978. The Late Devonian Palaeopalaemonidae and the earliest decapod crustaceans. Jour. Paleontol. 52:1375-1388.
- 307 \_\_\_\_\_. and J. Horner. 1978. Crustacea of the Mississippian Bear Gulch Limestone of central Montana. Jour. Paleontol. 52:394-406.
- 308 \_\_\_\_\_. and J.M. Schram. 1979. Some shrimp of the Madera Formation (Pennsylvanian) Manzanita Mountains, New Mexico. Jour. Paleontol. 53:169-174.
- 309 Scott, H.W. 1973. New Conodontochordata from the Bear Gulch Limestone (Namurian), Montana. Michigan State Univ. Paleontol. Ser. 1(2):85-99.
- 310 Scrutton, C.T. 1979. Early fossil cnidarians. Pp. 161-208. In: M.R. House, ed. The Origin of Major Invertebrate Groups. Syst. Assoc. Spec. Vol. 12. Academic Press; New York.
- 311 Sdzuy, K. 1974. Mittelkambrische Graptolithen aus NW-Spanien. Paläontol. Z. 48:110-139.
- 312 Shaw, A.B. 1974. Simpson Group (Middle Ordovician) trilobites of Oklahoma. Jour. Paleontol. 48(Suppl.):1-54.
- 313 Shergold, J.H. 1975. Late Cambrian and Early Ordovician trilobites from the Burke River structural belt, western Queensland, Australia. Aust. Bur. Miner. Res., Geol. Geophys. Bull. 153:1-251.
- 314 \_\_\_\_\_. 1976. Biostratigraphical synopsis: Eastern Georgina Basin, QLD. (Appendix to International Geological Congress Excursion 4C Guidebook). Aust. Bur. Miner. Res., Geol. Geophys. Record 1975/69, 50 pp.
- 315 \_\_\_\_\_. R.A. Cooper, D.I. MacKinnon, and E.L. Yochelson. 1976. Late Cambrian Brachiopoda, Mollusca, and Trilobita from northern Victoria Land, Antarctica. Palaeontology 19:247-291.

- 316 Shimer, H.W., and R.R. Shrock. 1944. Index Fossils of North America. M.I.T. Press; Cambridge, Massachusetts. 837 pp.
- 316a Smith, A.B. 1981. Implications of lantern morphology for the phylogeny of post-Palaeozoic echinoids. *Palaeontology* 24:779-801.
- 316b \_\_\_\_\_. 1982. The affinities of the Middle Cambrian Haplozoa (Echinodermata). *Alcheringa* 6:93-100.
- 317 Smith, A.G., N.F. Sohl, and E.L. Yochelson. 1968. New Upper Cretaceous Amphineura (Mollusca). U.S. Geol. Surv. Prof. Pap. 593-G, pp. G1-G9.
- 318 Sohl, N.F. 1960. Archaeogastropoda, Mesogastropoda and stratigraphy of the Ripley, Owl Creek, and Prairie Bluff Formation. U.S. Geol. Surv. Prof. Pap. 331-A, pp. A1-A151.
- 319 \_\_\_\_\_. 1964. Neogastropoda, Opisthobranchia, and Basommatophora from the Ripley, Owl Creek, and Prairie Bluff Formations — Late Cretaceous gastropods in Tennessee and Mississippi. U.S. Geol. Surv. Prof. Pap. 331-B, pp. B153-B344.
- 320 \_\_\_\_\_. 1965. Marine Jurassic gastropods, central and southern Utah. U.S. Geol. Surv. Prof. Pap. 503D, pp. D1-D29.
- 321 \_\_\_\_\_. 1967. Upper Cretaceous gastropods from the Pierre Shale at Red Bird, Wyoming. U.S. Geol. Surev. Prof. Pap. 393B, pp. B1-B46.
- 322 \_\_\_\_\_. 1967. Upper Cretaceous gastropod assemblages of the western interior of the United States. Pp. 1-38. In: E.G. Kauffman and H.C. Kent, eds. Paleoenvironments of the Cretaceous Seaway in the Western Interior: A Symposium. Colorado School of Mines Spec. Publ; Golden, Colorado.
- 323 \_\_\_\_\_. 1969. The fossil record of shell boring by snails. *Amer. Zool.* 9:725-734.
- 324 \_\_\_\_\_. 1971. North American Cretaceous biotic provinces delineated by gastropods. *Proc. N. Am. Paleontol. Conv.*, Chicago 1969, Pt. L, pp. 1610-1638.
- 325 Sohn, I.G. 1970. Early Triassic marine ostracodes from the Salt Range and Surghar Range, West Pakistan. Pp. 193-206, In: B. Kummel and C. Teichert, eds. Stratigraphic Boundary Problems: Permian and Triassic of West Pakistan. Univ. Kansas Press; Lawrence, Kansas.
- 326 Sokolov, B.S. 1972. Vendian and Early Cambrian Sabelliditida (Pogonophora) of the USSR. 1968 Proc. Int. Paleontol. Union, 23rd Int. Geol. Cong., Prague, pp. 79-86.
- 327 \_\_\_\_\_. 1972. The Vendian Stage in Earth history. 24th Int. Geol. Cong., Montreal, Sec. 1., pp. 78-84.
- 328 \_\_\_\_\_. 1973. Vendian of northern Eurasia. Pp. 204-218, In: M.G. Pitcher, ed. Arctic Geology. Am. Assoc. Pet. Geol.; Tulsa, Oklahoma.
- 329 \_\_\_\_\_. 1976. Precambrian Metazoa and the Vendian-Cambrian Boundary. *Paleontol. Jour.* 10:1-13.
- 330 Sprinkle, J. 1973. Morphology and evolution of blastozoan echinoderms. *Spec. Publ. Mus. Comp. Zool.*; Cambridge, Massachusetts, 283 pp.
- 331 \_\_\_\_\_. 1976. Biostratigraphy and paleoecology of Cambrian echinoderms from the Rocky Mountains. Pp. 61-73. In: R.A. Robison and A.J. Rowell, eds. Paleontology and Depositional Environments: Cambrian of Western North America. Brigham Young Univ. Geol. Stud. 23(2).

- 332 Stearn, C.W. 1980. Classification of the Paleozoic stromatoporoids. *Jour. Paleontol.* 54:881-902.
- 333 Stevens, G.R. 1973. Jurassic belemnites. Pp. 257-274. In: A. Hallam, ed. *Atlas of Palaeobiogeography*. Elsevier; Amsterdam.
- 334 \_\_\_\_\_. 1973. Cretaceous belemnites. Pp. 385-401. In: A. Hallam, ed. *Atlas of Palaeobiogeography*. Elsevier; Amsterdam.
- 335 Stinton, F.C. 1975-1980. Fish otoliths from the English Eocene. *Palaeontogr. Soc. Monogr. Pts. 1, 2, 3, and 4*, 258 pp.
- 336 Størmer, L. 1963. *Gigantoscorpio willsi*, a New Scorpion from the Lower Carboniferous of Scotland and its Associated Preying Microorganisms. Oslo Univ. Press; Oslo, 124 pp.
- 337 \_\_\_\_\_. 1972. Arthropods from the Lower Devonian (lower Emsian) of Alken an der Mosel, Germany. Pt. 2: Xiphosura. *Senckenbergiana leth.* 53:1-29.
- 338 \_\_\_\_\_. 1974. Arthropods from the Lower Devonian (lower Emsian) of Alken an der Mosel, Germany. Pt. 4: Eurypterida, Drepanopteridae, and other groups. *Senckenbergiana leth.* 54:359-451.
- 339 Sweet, W.C., and S.M. Bergström. 1972. Multielement taxonomy and Ordovician conodonts. *Geologica et Palaeontologica SB1*:29-42.
- 340 Sysoev, V.A. 1972. *Biostratigrafiya i khilolity ortotetsimorfii nizhnego Kembriya Sibirskoi Platoformy*. Nauka; Moscow, 152 pp.
- 341 Szaniawski, H. 1968. Three new polychaete jaw apparatuses from the Upper Permian of Poland. *Acta Palaeontol. Pol.* 8:255-280.
- 342 \_\_\_\_\_. 1974. Some Mesozoic scolecodonts congeneric with Recent forms. *Acta Palaeontol. Pol.* 19:179-195.
- 343 \_\_\_\_\_. and R.M. Wrona. 1973. Polychaete jaw apparatuses and scolecodonts from the Upper Devonian of Poland. *Acta Palaeontol. Pol.* 18:223-267.
- 344 Tappan, H. 1966. Chitinozoan classification. *Jour. Paleontol.* 40:1394-1396.
- 345 \_\_\_\_\_. and A.R. Loeblich, Jr. 1968. Lorica composition of modern and fossil Tintinnida (ciliate Protozoa), systematics, geologic distribution, and some new Tertiary taxa. *Jour. Paleontol.* 42:1378-1394.
- 346 Taylor, D.W., and N.F. Sohl. 1962. An outline of gastropod classification. *Malacologia* 1:7-32.
- 347 Taylor, J.D., N.J. Morris, and C.N. Taylor. 1980. Food specialization and the evolution of predatory prosobranch gastropods. *Palaeontology* 23:375-410.
- 348 Taylor, M.E. 1966. Precambrian mollusc-like fossils from Inyo County, California. *Science* 153:198-201.
- 349 \_\_\_\_\_. and H.E. Cook. 1976. Continental shelf and slope facies in the Upper Cambrian and lowest Ordovician of Nevada. Pp. 181-214. In: R.A. Robison and A.J. Rowell, eds. *Paleontology and Depositional Environments: Cambrian of Western North America*. Brigham Young Univ. Geol. Stud. 23(2).
- 350 Tchernycheva, N.E. 1959. *Système Cambrien*. Pp. 185-244. In: A.P. Markovsky, ed. *Structure Géologique de l'U.R.S.S. Tome 1. Stratigraphie* (P. de Saint-Aubin and J. Rogers, trans.). Centre National de la Recherche Scientifique; Paris.

- 351 \_\_\_\_\_, ed. 1965. Kembriyskaya Sistem. Nedra; Moscow, 596 pp.
- 352 Teichert, C. 1967. Major features of cephalopod evolution. Pp. 162-210. In: C. Teichert and E.L. Yochelson, eds. Essays in Paleontology and Stratigraphy, Univ. Kansas Press; Lawrence, Kansas.
- 353 \_\_\_\_\_, B.F. Glenister, and R.E. Crick. 1979. Biostratigraphy of Devonian nautiloid cephalopods. Pp. 259-262. In: M.R. House, C.T. Scrutton, and M.G. Bassett, eds. The Devonian System. Palaeontol. Assoc. Lond., Spec. Pap. Palaeontol. 23.
- 353a Thomson, K.S. 1969. The environment and distribution of Paleozoic Sarcopterygian fishes. Am. J. Sci. 267:457-464.
- 354 Thompson, I. 1979. Errant polychaetes (Annelida) from the Pennsylvanian Essex Fauna of northern Illinois. Palaeontographica. Abt. A. 163:169-199.
- 354a Thurmond, J.T., and D.E. Jones. 1981. Fossil Vertebrates of Alabama. Univ. Alabama Press; University, Alabama. 244 p.
- 355 Tozer, E.T. 1981. Triassic Ammonoidea: Classification, evolution and relationship with Permian and Jurassic forms. Pp. 66-100. In: M.R. House and J.R. Senior eds. The Ammonoidea. Syst. Assoc. Spec. Vol. 18. Academic Press; New York.
- 356 Turner, S. 1973. Siluro-Devonian thelodonts from the Welsh Borderland. Geol. Soc. Lond. Q.J. 129:557-584.
- 357 \_\_\_\_\_, and R.S. Dring. 1981. Late Devonian thelodonts (Agnatha) from the Gneudna Formation, Carnarvon Basin, Western Australia. Alcheringa 5:39-48.
- 357a Tynan, M.C. 1981. Microfossils from the Lower Cambrian Campito and Poleta Formations. P. 231. In: M.E. Taylor, ed. Short Papers for the Second International Symposium on the Cambrian System. U.S. Geol. Surv. Open-File Rpt. 81-743.
- 358 Valkov, A.K. 1975. Biostratigrafiya i khiolity Kembriya severovostoka Sibirskoi Platformi. Nauka; Moscow, 137 pp.
- 359 Viskova, L.A. 1972. Late Cretaceous cyclostomatous Bryozoa of the Volga River region and the Crimea (in Russian). Acad. Sci. USSR, Trans. Paleontol. Inst. 96 pp.
- 360 Voigt, E. 1973. *Hydrallmania grapolithiformis* n. sp., eine durch Biomuration erhaltene Sertulariidae (Hydrozo.) aus der Maastrichter Tuffkreide. Paläontol. Z. 47:25-31.
- 361 \_\_\_\_\_, 1980. *Arachnidium longicairda* n. sp. (Bryozoa Ctenostomata) aus der Maastrichter Tuffkreide (Ob. Kreide, Maastrichtium). N. Jb. Palaontol. Mb. 12:738-746.
- 361a \_\_\_\_\_, 1981. Critical remarks on the discussion concerning the Cretaceous-Tertiary Boundary. Newsl. Stratigr. 10:92-114.
- 362 Vologdin, A.G. 1977. Monotskiaty Kembriya SSSR. Nauka; Moscow, 156 pp.
- 363 Wade, M. 1969. Medusae from uppermost Precambrian or Cambrian sandstones, central Australia. Palaeontology 12:351-365.
- 364 \_\_\_\_\_, 1971. Bilateral Precambrian chondrophores from the Ediacaran fauna, South Australia. Proc. Roy. Soc. Victoria 84:183-188.
- 365 \_\_\_\_\_, 1972. Hydrozoa and Scyphozoa and other medusoids from the Precambrian Ediacara fauna, South Australia. Palaeontology 15:197-225.
- 366 \_\_\_\_\_, 1972. *Dickinsonia*: Polychaete worms from the late Precambrian Ediacara fauna, South Australia. Queensland Mus. Mem. 16:171-190.

- 367 Warn, J., and H.L. Strimple. 1977. The disparid inadunate superfamilies Homocrinacea and Cincinnaticrinacea (Echinodermata: Crinoidea), Ordovician-Silurian, North America. Bull. Amer. Paleontol. 72:1-138.
- 368 Waterhouse, J.B., and G.F. Bonham-Carter. 1975. Global distribution and character of Permian biomes based on brachiopod assemblages. Can. J. Earth Sci. 12:1085-1146.
- 369 Waterston, C.D. 1979. Problems of functional morphology and classification in stylonuroid eurypterids (Chelicerta, Merostomata), with observations on the Scottish Silurian Styloinuroidea. Trans. Roy. Soc. Edinburgh 70:251-322.
- 370 Webby, B.D. 1969. Ordovician stromatoporoids from New South Wales. Palaeontology 12:637-662.
- 371 Welch, J.R. 1976. *Phosphannulus* on Paleozoic crinoid stems. Jour. Paleontol. 50:218-225.
- 372 Werner, B. 1967. *Stephanoscyphus* Allman (Scyphozoa Coronatae) ein recenter Vertreter der Conulata? Paläontol. Z. 41:137-153.
- 373 Whittington, H.B. 1975. The enigmatic animal *Opabinia regalis*, Middle Cambrian, Burgess Shale, British Columbia. Phil. Trans. R. Soc., Lond., ser. B, 271:1-43.
- 374 \_\_\_\_\_. 1977. The Middle Cambrian trilobite *Naraoia*, Burgess Shale, British Columbia. Phil. Trans. R. Soc., Lond., ser. B, 280:409-443.
- 375 \_\_\_\_\_. 1977. *Aysheaia* and arthropod relations. Jour. Paleontol. 51(Suppl.):31.
- 376 \_\_\_\_\_. 1978. The lobopod animal *Aysheaia pedunculata*, Middle Cambrian, Burgess Shale, British Columbia. Phil. Trans. R. Soc., Lond., ser. B, 284:165-197.
- 377 \_\_\_\_\_. 1979. Early arthropods, their appendages and relationships. Pp. 253-268. In: M.R. House, ed. The Origin of Major Invertebrate Groups. Syst. Assoc. Spec. Vol. 12; Academic Press, New York.
- 378 \_\_\_\_\_. and C.P. Hughes. 1972. Ordovician geography and faunal provinces deduced from trilobite distribution. Phil. Trans. R. Soc., Lond., ser. B, 263:235-278.
- 379 \_\_\_\_\_. and \_\_\_\_\_. 1974. Geography and faunal provinces in the Tremadoc Epoch. Pp. 203-218. In: C.A. Ross, ed. Paleogeographic Provinces and Provinciality. Soc. Econ. Paleontol. Mineral. Spec. Publ. 21.
- 380 Williams, A. 1974. Ordovician brachiopods from the Shelve District, Shropshire. Br. Mus. (Nat. Hist.) Bull., Geol. Suppl. 2, 163 pp.
- 381 Wilson, J.L. 1957. Geography of olenid trilobite distribution and its influence on Cambro-Ordovician correlation. Am. J. Sci. 255:321-340.
- 382 Wright, A.D. 1979. Brachiopod radiation. Pp. 235-252. In: M.R. House, ed. The Origin of Major Invertebrate Groups. Syst. Assoc. Spec. Vol. 12. Academic Press; New York.
- 383 Wright, C.W. 1981. Cretaceous Ammonoidea. Pp. 157-174. In: M.R. House and J.R. Senior, eds. The Ammonoidea. Syst. Assoc. Spec. Vol. 18. Academic Press; New York.
- 384 \_\_\_\_\_. and J.S.H. Collins. 1972. British Cretaceous crabs. Palaeontogr. Soc. Monogr., 114 pp.
- 385 Yancey, T.E. 1978. Brachiopods and molluscs of the lower Permian Arcturus Group, Nevada and Utah. Part 1: Brachiopods, scaphopods, rostroconchs, and bivalves. Bull. Amer. Paleontol. 74:257-367.
- 386 Yochelson, E.L., and B.W. Saunders. 1967. A bibliographic index of North American Late

- Paleozoic Hyolitha, Amphineura, Scaphopoda, and Gastropoda. U.S. Geol. Surv. Bull. 1210, 271 pp.
- 387 \_\_\_\_\_, 1969. Stenothecoida, a proposed new class of Cambrian Mollusca. Lethaia 2:49-62.
- 388 \_\_\_\_\_, 1975. Discussion of Early Cambrian "molluscs", Jour. Geol. Soc. Lond. 131:661-662.
- 389 \_\_\_\_\_, 1977. Agmata, a proposed extinct phylum of early Cambrian age. Jour. Paleontol. 51:437-454.
- 390 \_\_\_\_\_, J.F. McAllister, and A. Reso. 1965. Stratigraphic distribution of the Cambrian mollusk *Matthevia* Walcott 1885. U.S. Geol. Surv. Prof. Pap. 525B, pp. B73-B78.
- 391 \_\_\_\_\_, J.W. Pierce, and M.E. Taylor. 1970. *Salterella* from the Cambrian of central Nevada. U.S. Geol. Surv. Prof. Pap. 643H, pp. H1-H7.
- 391a \_\_\_\_\_, and G.D. Stanley, Jr. 1981. An early Ordovician patelliform gastropod, *Palaeophacmaea*, reinterpreted as a coelenterate. Lethaia 15:323-330.
- 392 Zangerl, R. 1981. Chondrichthyes I. Paleozoic Elasmobranchii. Handbook of Paleichthyology, v. 3A. Gustav Fischer Verlag; Stuttgart. 115 p.
- 393 Zhou Z. and Yuan J. 1981. The biostratigraphic distribution of Lower Cambrian trilobites in southwest China. Pp. 250-251. In: M.E. Taylor, ed. Short Papers for the Second International Symposium on the Cambrian System. U.S. Geol. Surv. Open-File Rpt. 81-743.
- 394 \_\_\_\_\_, and Zhang, J. 1978. Cambrian-Ordovician Boundary of the Tangshan area with descriptions of the related trilobite fauna. Acta Palaeontol. Sin. 17:1-28.
- 395 Zhuravleva, I.T. 1970. Marine faunas and Lower Cambrian stratigraphy. Am. J. Sci. 269:417-445.
- 396 Zinmeister, W.J. 1981. Middle to Late Eocene invertebrate fauna from the San Julian Formation at Punta Casamayor, Santa Cruz Province, southern Argentina. Jour. Paleontol. 55:1083-1102.

## ACKNOWLEDGMENTS

I thank Drs. R.L. Anstey, R.E. Grant, S. Kelly, G.V. Lauder, Jr., W.D.I. Rolfe, P.M. Sheehan and N.F. Sohl for their kind help in this project and especially D.B. Blake, R.E. Crick, R.E. Plotnick, and F.R. Schram for permitting me to use their unpublished compilations on bryozoans, nautiloids, eurypterids, and crustaceans, respectively. T.J.M. Schopf, C. Teichert, and A.M. Ziegler graciously gave me free access to their extensive private libraries. I take responsibility for any errors and omissions.

This research received partial support from the Systematic Biology Program, National Science Foundation, NSF Grant DEB81-08890.

## REFERENCES

- Harland, W.B., et al., eds. 1967. *The Fossil Record*. Geological Society of London; London. 828 p.
- Kukalová-Peck, J. 1973. A Phylogenetic Tree of the Animal Kingdom (Including Orders and Higher Categories). National Museums of Canada, Publications in Zoology, No. 8. 78 p.
- Moore, R.C., C. Teichert, and R.A. Robison, eds. 1953-1981. *Treatise on Invertebrate Paleontology*. Geological Society of America and University of Kansas Press; Lawrence, Kansas.
- Romer, A.S. 1966. *Vertebrate Paleontology*, 3rd ed. University of Chicago Press; Chicago. 468 p.
- Sepkoski, J.J., Jr. 1978. A kinetic model of Phanerozoic taxonomic diversity: I. Analysis of marine orders. *Paleobiology* 4:223-251.
- \_\_\_\_\_. 1979. A kinetic model of Phanerozoic taxonomic diversity: II. Early Phanerozoic families and multiple equilibria. *Paleobiology* 5:222-251.
- \_\_\_\_\_. 1981a. A factor analytic description of the marine fossil record. *Paleobiology* 7:36-53.
- \_\_\_\_\_. 1981b. The uniqueness of the Cambrian fauna. Pp. 203-207. In: M.E. Taylor, ed. *Short Papers for the Second International Symposium on the Cambrian System*. U.S. Geol. Surv. Open-File Rpt. 81-743.